Museum

7991 . 787 v. 96 . no. lb

## THE ZOOLOGICAL RECORD

VOLUME 96 SECTION 16 1959

# **AMPHIBIA**

(1958)

COMPILED BY

ALICE G. C. GRANDISON, B.Sc.,
MARCIA A. EDWARDS, B.Sc.,
and
PAULINE ARMITAGE, B.Sc.

LONDON
PUBLISHED BY
THE ZOOLOGICAL SOCIETY OF LONDON

PRICE TWELVE SHILLINGS



It will greatly help in the production of The Zoological Record and assist the Recorders of the individual sections, if authors will forward a copy of their paper or memoir to the Editor of the Zoological Record, The Zoological Society of London, Regent's Park, London, N.W.1. In the case of separately printed copies of articles so forwarded, the original pagination should be given.

All business correspondence concerning the Record should be addressed to the Scientific Director, The Zoological Society of London, Regent's Park, London, N.W.1.

### THE MUSEUMS JOURNAL

Published by the Museums Association
22 Fitzroy Street, Fitzroy Square, London, W.1.

This journal, which is published monthly, contains articles, reviews and technical notes on every Museum aspect of zoology and other sciences. It is the only English journal which deals with the installation and preservation of exhibits, and which reviews Museum work in all parts of the world.


It is published monthly price 4s., by the Association and can be obtained from the above address. It is distributed free to all members of the Museums Association (Subscription £3 3s. 0d. per annum), of which full particulars are obtainable from the Secretary at the above address.

## **AMPHIBIA**

COMPILED BY

ALICE G. C. GRANDISON, B.Sc., MARCIA A. EDWARDS, B.Sc., AND

PAULINE ARMITAGE, B.Sc.



# 16. AMPHIBIA (1958)

COMPILED BY

ALICE G. C. GRANDISON, B.Sc., MARCIA A. EDWARDS, B.Sc., AND

PAULINE ARMITAGE, B.Sc.

		CO	NTE	ENTS				
							PAGE	
I. TITLES	• • •		•••	***	***	•••	•••	4
II. SUBJECT IN	DEX			***	***	***	• • •	27
General		***		• • •	***		***	27
Structure	• • •		• • •	• • •		***	•••	28
Physiology	7		***	***	***		•••	30
Reproduct	ion a	and Sex	•••	•••			***	35
Developme	ent	***	***		***			35
Evolution	and	Genetics	***		• • •			38
Ecology an	nd H	abits	***	***	***		***	39
Geographical Distribution						***	***	41
Palaeontol	ogy	and Geol	ogica	al Distri	bution		***	42
III. SYSTEMATIC	IND	EX	***	•••	•••	***	***	42
General				***	***	***		42
Gymnophi	ona	***				***	***	43
Caudata	•••	•••			***	***	***	43
Anura				•••	•••	• • •		47

#### I.—TITLES

Opinion 520. Suppression under the Plenary Powers of the specific name tibiatrix Laurenti, 1768, as published in the combination Hyla tibiatrix, and of the generic name Acrodytes Fitzinger, 1843, and interpretation under the same Powers of the nominal species Rana venulosa Laurenti, 1768 (Class Amphibia). Opin. int. Comm. zool. Nom. 19 (7) 1958: 169-200.

Anon, [E. H. C.] [Karl Patterson Schmidt—obituary.] Curator 1 1958: 19.

- (1) Anon. Normal table of Xenopus laevis (Daudin) edited by P. D. Nieuwkoop and J. Faber. [Review.] Copeia 1958 1: 65.
- (2) Anon. Die Reptilien und Amphibien mitteleuropas by R. Sternfeld. [Review.] Copeia 1958 1: 65.
- (3) Anon. Tree frog [Hyla cinerea.] Turtox News 36 6, 1958: 149, 1 pl.
- (4) Anon. Giant tree frog [Hyla maxima]. Turtox News 36 1 1958: 46, 1 pl.
- (5) Anon. Joseph R. Slevin 1881–1957. Acad. News Letter. Cal. Acad. Sci. 207 1957: 2–3, photograph.
- Adams, M. S. & Clark, H. F. A herpetofaunal survey of Long Point, Ontario, Canada. Herpetologica 14 (1) 1958: 8-10.

Adler, J. Atelopus stelzneri. Aquar. Terrar. Z. 10 1957: 190-192.

(1) Adler, K. K. A Field Guide to Reptiles and Amphibians of Eastern North America by Roger Conant. [A review.] Newsletter Trimon. Rep. Ohio herpet. Soc. 1 4 1958: 2.

(2) Adler, K. K. A five-legged Rana from Ohio, Trimon. Rep. Ohio herpet. Soc. 1 4 1958: 21.

Aellen, W. Sur une nouvelle forme d'Hydromantes (Amphibia, Plethodontidae). Senck. biol. 39 1958: 155–163, 3 figs.

Aguggini, G. & Noseda, V. Depolarizing action of K-Strophantine and K-Strophantoside on isolated frog skin. Experientia 14 1958: 458-460, 3 figs.

Aguilera, S. E., see Cei, J. M.

- (1) Alcala, A. C. Philippine notes on the ecology of the giant marine toad. Silliman J. 4 1957: 90-96 1 fig.
- (2) Alcala, A. C. Amphibians of Negros Island. Including 2 new records. Silliman J. 5 1958: 171–174, 1 pl.

Alcala, A. C. see Rabor, D. S.

Alexander, D. R. & Diener, B. A. A collection of amphibians and reptiles from Japan. Copeia 1958 No. 3: 218-219.

Allen, B. M. Interpretation of some experiments upon the effect of ionizing irradiation on the tissues of amphibians. Ann. Zool., Agra 2 1958: 119-126 1 pl.

Allende, I. L. C. de & Astrada, J. J. Elimination des spermatozoïdes du Crapaud par action de la gonadotrophine chorionique. C.R. Soc. Biol. Paris 151 1958: 1964-1966.

Allison, R. M. The sensitivity of Bufo bufo to chorionic gonadotrophin throughout the year and after periods of enforced hibernation. J. Endocrinol. 14 1956: 275-278.

Amanuma, A., see Asayama, S.

Ambellan, E. Comparative effects of Moni, Di, and Triphosphorylated Nucleosides on amphibian morphogenesis, J. Embryol. exp. Morph. 6 1958: 86-93, 2 figs.

Ancona, U. D. Studi e ricerche sul differenziamento sessuale dei vertebrati, sulla biologia degli acridioidei e sulla gametogenesi degli insetti. Ric. sci. 28 1958: 1653–1658. [English, etc. summaries.]

Anderson, E. & Breemen, V. L. van. Electron microscopic observations on spinal ganglion cells of Rana pipiens after injection of Malononitrite. J. biophys. and biochem. Cytol. 4 1958: 83-86 4 pls.

Anderson, J., see Leaf, A.

Anderson, P. K. Inducted oviposition in Batrachoseps attenuatus, and incubation of eggs. Copeia 1958 No. 3: 221-222.

Anderson, S. C. & Volpe, E. P. Burnsi and Kandiyohi genes in the leopard frog Rana pipiens. Science 127 1958: 1048-1049 1 fig.

Andrews, T. F., see Clarke, R. F.

Aoki, K. Über die Wasserabsorption des Salamanderies. J. Fac. Sci. Hokkaido Univ. 7 1940: 381–398 1 fig.

Archdeacon, J. W., see Wilson, J. W.

Arima, S. Studies on the end products of metabolism in the early developmental stages of Amphibia. IV Ammonia-, Urea-, and iron-contents in the perivitelline fluid of Rhacophorus schlegelii var. arborea. Annot. Zool. jap. 31 1958: 136-140.

Arnoult, J. & Lamotte, M. Contribution à l'étude des Batraciens de l'Ouest africain. VI, Développement lavaire de deux espèces d'*Hyperolius*: *H. zonatus* Laurent et *H. lamottei* Laurent. Bull. Inst. franç. Afr. N. 20A 2 1958: 573-586 18 figs.

Asandei, A., see Degan, C.

Asano, T. & Hurlbut, W. P. Effects of potassium, sodium, and azide on the ionic movements that accompany activity in frog nerves. J. gen. Physiol. 41 1958: 1187-1203 7 figs.

Asayama, S. & Amanuma, A. Modification of sex development in *Hynobius nebulosus* by administration of methyl-testo-sterone. J. Inst. Polyt. Osaka City Univ. **3D** [1958]: 111–128 2 pls, 1 fig.

Asayama, 8. & Matsuzaki, H. Responses of the gonads of Triturus pyrrhogaster larvae to the administration of methyltestosterone. Zool. Mag., Tokyo 67 1958: 133–137, 10 figs. [In Japanese, English summary.]

Bi

[

goi of

de

. .

ve

20

an 3

at

12 gu 19

tr So

m

Session

1

a

L

•

.

a

d

t

n

- Asayama, S. & Miyamori, H. Responses of larval gonads in *Hymobius nebulosus* to the administration of estrogen. J. Inst. Polyt. Osaka City Univ. 8D [1958]: 129-144 pl. 2.
- Astrada, J. J. A. & Taleisnik, S. Action in vitro des solutions salines et hypophysaires sur l'imbibition testiculaire et le spermiation du Crapaud. C.R. Soc. Biol. Paris 151 1958: 1961–1963.
  - Astrada, J. J., see Allende, I. L. C., de.
- Auffenberg, W. A new species of Bufo from the Pliocene of Florida. Quart. Jour. Fla. Acad. Sci. 20 1 1957: 14-20.
  - Auffenberg, W., see Goin, C. J.
  - Bacq, Z. M., see Liébecq-Hutter, S.
- Baehner, R. L. see Peters, J. J.
- Baer, J. G. Cours d'anatomie comparée des vertébrés. Neuchatel (Griffon) 1958, pp. 207, 68 pls. col.
- ·Baffoni, G. M. Influenza dell' ormone tiroideo sull' attività mitotica del rombencefalo di un Anfibio anuro. R.C. Accad. Lincei (8) 23 1958: 495–503 3 figs.
- Baffoni, G. M. & Pinacci, R. L'andamento dell' attività mitotica nel midollo spinale di Anfibi anuri durante lo sviluppo. R.C. Accad, Lincei (8) 25 1958: 128-134 2 figs.
- (1) Bagnara, J. T. Hypophyseal control of guanophores in anuran larvae. J. exp. Zool. 137 1958: 265–283 10 figs.
- (2) Bagnara, J. T. Further observations on the tail melanophores of *Xenopus*. Anat. Rec. 131 1958: 529-530. [Abst.]
- Bagnara, J. T. & Neidleman, S. Effect of chromatotrophic hormone on pigments of anuran skin. Proc. Soc. exp. Biol. Med. 97 1958: 671-673 2 figs.
- (1) Baldauf, R. J. Contributions to the cranial morphology of Bufo valliceps Wiegmann. Texas J. Sci. 10 1958: 172-186 16 figs.
- (2) Baldauf, R. J. A procedure for the staining and sectioning of the heads of adult anurans. Texas J. Sci. 10 1958: 448-451.
- Baldisserotto, A. & Ugurgieri, B. I fenomeni di autolisi metamorfosi di *Bufo vulgaris*. R.C. Accad. Lincei (8) 25 1958: 81–86 2 figs.
- Balinsky, B. I. On the factors controlling the size of the brain and eyes in anuran embryos. J. exp. zool. 139 1958: 403–438, 13 figs., 1 pl.
- Bannikov, A. G. Data on the fauna and biology of Amphibia and Reptilia in Mongolia. Bull. Soc. nat. Moscow biol. 63 2 1958 71–92 8 figs. [In Russian with English summary.]
- Barbour, B. W. A deletion of distribution records of Hyla squirella. Copeia 1958 No. 1: 49-50.
- Barch, S. H. & Shaver, J. R. The effects of enzymes on prolongation of fertilizability of Rana pipiens eggs. Anat. Rec. 131 1958: 530. [Abst.]
- Barth, L. G. & L. J. Control of differentiation of cells from Rana pipiens gastrulae. Anat. Rec. 131 1958: 530-531. [Abst.].

- Barth, L. J., see Barth, L. G.
- Bašoglu, M. & Freytag, G. E. Über ein neues Exemplar von Mertensiella luschani (Steindachner). Zool. Anz. 160 1958: 20-25 2 figs.
  - Baylé, J. D., see Jullien, A.
- Bebak, B. Thermal stadia in the development of the grass frog (Rana temporaria L.). Bull. Acad. polon. Sci. Ser. Sci. biol. 6 1958: 367-369 3 figs.
- Becerra, B. G. Influences hormonales sur la diurèse du Crapaud. C.R. Soc. Biol. Paris 151 1958: 1966–1968.
- Bechtle, W. Die hochzeit der Grasfrösche. Kosmos, Stuttgart 54 1958: 130–132.
- Becker, U. Über die Korrelation zwischen der Linsenentwicklung und der Gesamtentwicklung des Keims bei *Triturus vulgaris*. Z. Naturf. **13B** 1958 : 208–213 2 figs.
- Bedate, M. A. Análisis experimental de la pigmentación dorsal de *Discoglossus pictus*. Bol, Soc. esp. Hist. nat. Sec. Biol. **55** 1958: 415-419.
- Beenakkers, A. M. Th., see Oordt, G. J. van.
- (1) Beetschen, J.-Cl. Élevage en parabiose d'individus hétéroploïdes hypomorphes chez le Triton, Pleurodeles walthi Michah. C.R. Acad. Sci. Paris 246 1958: 2672-2674 3 figs.
- (2) Beetschen, J. Cl. Étude cytologique de germes hétéroploïdes létaux chez le triton, *Pleurodeles walthii*, Michah. Bull. Soc. zool. Fr. **83** 1958 : 242–244.
- (3) Beetschen, J. Cl. Anomalies mitotiques liées à l'aneuploidie d'origine génétique, chez le Pleurodèle. Bull. Soc. zool. Fr. 83 1958 : 245.
- (1) Bellec, A. Contribution a l'étude expérimentale de l'apparition du glycogène hépatiqué chez Pleurodeles walthis Michah. Arch. Anat. micr. Morph. exp. 47 1958: 171-182 8 figs.
- (2) Bellec, A. Apparition du glycogène en culture in vitro de foie larvaire de Pleurodèle. C.R. Soc. Biol. Paris 151 1958: 1353-1355.
  - Bennett, S., see Simon, S. E.
- (1) Bentley, P. J. The effects of vasopressin on water uptake of the toad, Bufo marinus, while bathed different hypotonic solutions. J. Endocrin. 16 1957: 126-134.
- (2) Bentley, P. J. The effects of neurohypophysial extracts on water transfer across the wall of the isolated urinary bladder of the toad Bufo marinus. J. Endocrin. 17 1958: 201-209 1 fig.
- (3) Bentley, P. J. Loss of sodium from the skin of the dehydrated toad, *Bufo marinus*. Nature, Lond, 182 1958: 1810
- Bentley, P. J., Lee, A. K. & Main, A. R. Comparison of dehydration and hydration of two genera of frogs (*Heleioporus* and *Neobatrachus*) that live in areas of varying aridity. J. exp. Biol. 35 1958: 677-684 1 fig.
- Ber, A. & Zieleniewski, J. Studies on the side effects of antibiotics. Report 1. The effect of penicillin, streptomycin and penicillium mycelium on the growth and development of *Xenopus laevis* Daudin. Zool. polon., 8 1957 [1958]: 151-160 3 figs. In Polish, English summary.

[

her

fro

20

of

an

re

na

Te J.

di

m

th

13

0

n

e

la

F

Berezin, A., see Hadler, W. A.

Bern, H. A., see Wilson, L. D.

Berner, L. Amphibies et Reptiles des environs de Marseille, Bull, Soc. linn, Provence 20 1955 : 45-46.

**Beznák, A. B. L.** Effect of acetylcholinesterase on the surviving frog heart. Nature, Lond. **181** 1958: 1190–1192 3 figs.

Bhaduri, J. L. Himadri Kumar Mookerjee 1898–1955. Proc. zool. Soc. Calcutta, Mookerjee Memor. Vol., 1957: 1-11 photograph.

Bhattacharya, G. C. & Medda, A. K. Induced metamorphosis of tadpoles (*Bufo melanostictus*). Sci. and Cult. 23 1958: 380–382 2 figs.

Bijtel, J. H. The mode of growth of the tail in urodele larvae. J. Embryol. exp. Morph. 6 1958: 466-478 2 pls., 3 figs.

(1) Billett, F. β-Glucuronidase activity during the early development of Xenopus laevis embryos. Biochem. J. 67 1957: 463-466 2 figs.

(2) Billett, F. The temporary inactivation of newt larvae by benzimidazole and its alkyl derivatives. Experientia 14 1958: 99-100.

Biörck, G. & Johansson, B. Comparative studies on temperature effects and upon the electrocardiogram in some vertebrates. Acta physiol. scand. 34 1955: 257-272.

Birke, R., see Pilz, A.

Black, I. H. & Gosner, K. L. The barking tree frog, Hyla graticsa in New Jersey. Herpetologica 13 1958: 254–255.

Black, I. H., see Gosner, K. L.

Blackler, A. W. Contribution to the study of germ-cells in the Anura. J. Embryol. exp. Morph. 6 1958: 491-503 2 pls.

(1) Blair, W. F. Distribution patterns of vertebrates in the Southern United States in relation to past and present environments. Publ. Amer. Ass. Advanc. Sci. 51 (Zoogeography) 1958: 433-468 11 figs.

(2) Blair, W. F. Response of a green tree frog (Hyla cinerea) to the call of the male. Copeia 1958 No. 4: 333-334.

(3) Blair, W. F. Call difference as an isolation mechanism in Florida species of hylid frogs. Quart. J. Fla. Acad. Sci. 21 1958: 32-484 figs.

(4) Blair, W. F. Mating call in the speciation of anuran amphibians. Amer. Nat. 92 1958: 27-51 7 figs.

(5) Blair, W. F. Mating-call and stages of speciation of two allopatric populations of spadefoots (Scaphiopus). Texas J. Sci. 10 1958: 484–488.

Blair, W. F., see Hecht, M. K.

Bleakney, J. S. A zoogeographical study of the amphibians and reptiles of eastern Canada. Bull. nat. Mus. Canada No. 155 1958: 1-119 49 maps.

Bleakney, S. Cannibalism in Rana sylvatica tadpoles a well known phenomenon. Herpetologica 14 1958: 34.

Blount, R. W., see Flickinger, R. A.

Blum, H. F., see Butler, E. G.

Bodemer, C. W. The development of nerveinduced supernumerary limbs in the adult newt, Triturus viridescens. J. Morph. 102 1958: 555-581 19 figs.

Boettger, C. R. Haltung grosser Erdkröten. Natur u. Volk 88 1958 : 235–236.

Bogert, C. M. The first voices of spring. Frogs' calls are known to serve a mating purpose. Is that all? Nat. Hist. N.Y. 67 1958: 182–189 with figs.

Bogert, C. M., see Oliver, J. A.

(1) Bokermann, W. C. A. Una nueva especie del genero Sphoenohyla del Brasil. Neotropica 4 (14) 1958: 43–46 6 figs.

(2) Bokermann, W. C. A. Sobre una nueva especie de Dendrobates del Norte del Brasil. Neotropica 4 (15) 1958: 73-76, 9 figs.

(3) Bokermann, W. C. A. A preoccupied name of a neotropical frog, genus *Eleutherodactylus*. Herpetologica 14 1958: 95.

Bondi, C. Nuove ricerche sperimentali nei trapienti di abbozzi di *Triton* su *Rana esculenta*. Roux Arch, Entwick.-Mech. Organ. **150** 1958: 689-701 6 figs.

Bondi, C., see Spirito, A.

Borghgraef, R. & Carmeliet, E. Étude électrophysiologique du myocarde d'Amphibien en absence de potassium et sous l'influence de l'acétylcholine. C.R. Soc. Biol., Paris 151 1958: 1779–1781.

Boternbrood, E. C. Organization in aggregates of anterior neural plate cells of *Triturus alpestris*. Proc. Acad. Wet. Amst. **61C** 1958: 470-481 4 figs.

Bourguet, J. Action de la désoxycorticostérone sur le transport actif de sodium au niveau de la peau de Grenouille isolée. Influence du calcium. C.R. Soc. Biol., Paris 151 1958: 1782–1783.

Brachet, J. Effects of β-mercaptoethanol on morphogenesis in amphibian eggs. Nature, Lond. 181 1958: 1736–1737 1 fig.

(1) Bragg, A. N. On metamorphic and postmetamorphic aggregations in spadefoots. Herpetologica 13 1958: 273.

(2) Bragg, A. N. The egg of the dwarf American toad. Herpetologica 13 1958: 273-275.

(3) Bragg, A. N. Taxonomic status of the gray tree frog in Oklahoma. Herpetologica 14 1958: 79-84.

Bragg, A. N. & W. N. Parasitism of spadefoot tadpoles by Saprolegnia. Herpetologica 14 1958: 34.

Bragg, A. N. & Brooks, M. Social behaviour in juveniles of *Bufo cognatus* Say. Herpetologica 14 1958: 141-147.

Brahma, S. K. Experiments on the diffusibility of the amphibian evocator. J. Embryol. exp. Morph. 6 1958: 418-423 1 pl.

Brandon, A. M., see Cordier, D.

(1) Brattstrom, B. H. Additions to the herpetofauna of Cass County, Minnesota. Herpetologica 13 1958: 278, 1

1)

ti

h.

of

e.

18

n

y

n

4

6

(2) Brattstrom, B. H. Additions to the Pleistocene herpetofauna of Nevada. Herpetologica 14 1958: 36.

Breemen, V. L. van., see Anderson, E.

Breukelman, J., see Clarke, R. F.

Brice, M. C. A re-analysis of the consequences of frontal and sagittal constrictions of newt blastulae and gastrulae. Arch. Biol. Paris 69 1958: 371-439, 1 pl., 20 figs.

(1) Brindley, G. S. The passive electrical properties of the frog's retina, choroid and sclera for radial fields and currents. J. Physiol. 134 1956: 339–352.

(2) Brindley, G. S. The effect on the frog's electro retinogram of varying the amount of retina illuminated. J. Physiol. 134 1956: 353-359.

(3) Brindley, G. S. Response to illumination recorded by microelectrodes from the frog's retina. J. Physiol. 134 1956: 360-384.

Briner, G. P., Simon, S. E. & Shaw, F. H. The distribution of inorganic phosphate in amphibian muscle. J. gen. Physiol. 41: 755-766.

**Brode, W. E.** The occurrence of the pickerel frog, three salamanders and two snakes in Mississippi caves. Copeia **1958** (1): 47-48.

Brode, W. E. & Gunter, G. Egg clutches and prehensilism in the slimy salamander. Herpetologica 13 (4) 1958: 379–380.

Brodkorb, P., see Neill, W. T.

Brodsky, V. & Nechaeva, N. V. On the relations observed between the quantitative changes of ribonucleic acid, the intensity of function and the trophic condition of a neuron (exemplified by data of a cytochemical examination of the ganglionic cells of the retina). C.R. Acad. Sci. URSS 123 4 1958 756-759. [In Russian.]

Brongersma, L. D. The animal world of Netherlands New Guinea. Groningen (J. B. Wolters) 1958: pp. 71, 34 figs.

Brooks, M., see Bragg, A. N.

Brouant, J. F. & Cury, G. Effects du retournement de l'oeuf sur le développement embryonnaire de Pleurodeles walthi. C.R. Soc. Biol. Paris 151 1958 : 1413-1415.

Brunner, G. Nachtrag zur Breitenberghöhle bei Gössweinstein (ofr.). Neues Jb. Geol. Paläont. Monatshefte 1958: 500-517 figs.

Brunst, V. V. Effect of irradiation on some portions of head of young Axolotl (Siredon mexicanum). Proc. Soc. exp. Biol. Med. 99 1958: 571-574.

Bruyn, H. T. de., see Kappers, J. A.

Buchholz, K. F. Zum Nahrungserwerb von Salamandra salamandra (Amphibia). Bonn. zool. Beitr. 9 1958: 63–67.

Bücker, H. & Hanke, W. Comparison of the effect of ultra violet and x-rays on the capillaries of the frog's web. Nature, Lond. 182 1958: 1375-1376 2 figs.

Burgers, A. C. J. Investigations into the action of certain hormones and other substances on the melanophores of the South African Clawed Toad, Xenopus laevis. G. W. Van der Wiel & Co., Arnhem 1956 95 pp. 14 figs.

Burgers, A. C. J. & Mighorst, J. C. A. Is the melanophore-dispersing substance in human pregnancy urine a 3-4.-dihydroxybenzine derivative? Acta Endocrinol. 25 1957: 249-258.

Burgos, M. H. & Houssay, B. A. Action de LH et FSH sur la spermiation du *Bufo arenarum*. C.R. Soc. Biol. Paris **152** 1958: 197-198.

Burgos, M. H. & Ladman, A. J. The effects of purified gonadotrophins on the morphology of the testes and thumb pads of the normal and hypophysectomized autumn frog (Rana pipiens). Endocrinology 61 1957: 20–34 4 pls.

Burkhardt, D. Die Linnesorgane des Skeletmuskels und die nervöse Steuerung der Muskeltätigkeit. Ergebn. Biol. **20** 1958: 27–66 12 figs.

Butler, E. G., Blum, H. F. & Schmidt, S. E. The localized character of ultraviolet effects on the urodele forelimb. J. cell. comp. Physiol. 50 1957: 381–388.

Callan, H. G. & Maegregor, H. C. Action of deoxyribonuclease on lampbrush chromosomes. Nature, Lond. 181 1958: 1479-1480 1 fig.

Cambar, R. & Dechambe, C. Le matérial cellulaire néphrostomial du pronéphros ne participe pas à l'édification du pavilon génital femelle (ostium); observations chez la Grenouille agile (Rana dalmatina Bon.). C.R. Acad. Sci., Paris 246 1958: 174-176 figs.

Cambar, R. & Gipouloux, J.-D. Les différentes régions du blastème de l'uretère primaire possèdent initialement les mêmes potentialités morphogènes : démonstration expérimentale (Amphibiens anoures). C.R. Acad. Sci. 246 1958 : 312-315, 2 figs.

Campbell, H., see Holman, J. A.

Capuani, M. Considerazioni su alcuni risultati di trattamento con benzopirene in anfibi urodeli [Some results of benzopyrene treatment in urodele amphibians (*Triturus cristatus*).] Riv. Biol. Perugia 49 1957: 151–161.

(1) Capurro, L. Lista preliminar de los anfibios de Chile y breves apuntes sobre su distribución y biología. Invest. 2001. chil. 4 1958: 289–299.

(2) Capurro, L. F. Nota sobre la ovipostura y desarrollo de Eupsophus taeniatus (Girard). Invest. zool. chil. 4 1958: 208.

Capurro, L., see Cei, J. M.

Carasso, N. Ultra-structure des cellules visuelles de larves d'Amphibiens, C.R. Acad. Sci. Paris 247 1958: 527-531 4 pls.

Cardeza, A. F., see Penhos, J. C.

Cardot, J., see Jullien, A.

Carl, G. C. Salamander sallies. Victoria Nat. 15 1958: 30-32.

Carl, G. C. & Guiguet; C. J. Alien animals in British Columbia. Handb. B. C. Prov. Mus. No. 14 1958: 61-63 2 figs.

Carmeliet, E., see Borghgraef, R.

Carter, M. E., see Griffiths, I.

Carvalho, A. L. de., see Lutz, B.

Castro, V., see Sarzana, G.

- Cei, J. M. Polimorfismo y distribución geográfica en poblaciones chilenas de *Pleurodema* bibroni Tschudi. Invest. 2001. chil. 4 1958: 309–327
- (2) Cei, J. M. Geographical isolation and phenotypic constitution of populations of Lepidobatrachus asper, a neotropical batrachian inhabitant of arid regions. Evolution 12 1958: 166-172 5 figs.
- (3) Cei, J. M. Karl P. Schmidt 1890-1957. Invest. zool. chil. 4 1958: 129-131 photo.
- (4) Cei, J. M. Las laminas originales del suplemento a los Batraquios Chilenos de Philippi: primera impresión y comentario. Invest. zool. chil. 4 1958: 265-278 10 pls.
- Cei, J. M. & Aguilera, S. E. La vibración sexual preventiva ("warning vibration") en *Pleurodema* chilenas. Invest. zool. chil. 4 1958: 15-21 2 figs.
- Cei, J. M. & Capurro, L. Biologia y desarrollo de Eupsophus taeniatus Girard. Invest. zool. chil. 4 1958: 159–182 19 figs.

Cerf, J. A., see Koketsu, K.

Cerl, J. A. & Chacko, L. W. Retrograde reaction in motoneuron dendrites following ventral root section in the frog. J. comp. Neurol. 109 1958: 205-218 1 pl.

Chacko, L. W., see Cerf, J. A.

Chalfin, D., Cooperstein, I. L. & Hogben, C. A. M. Fluid and electrolyte movement across intestinal wall of bullfrog. Proc. Soc. exp. Biol. Med. 99 1958: 746-748.

Chang, C. & Witschi, E. Cortisone effect on ovulation in the frog. Endocrinology 61 1957 514-519 4 figs.

Chang, C. Y., see Witschi, E.

- Chapman, B. M. & R. F. A field study of a population of leopard toads (Bufo regularis regularis). J. Anim. Ecol. 27 1958: 265–286 10 figs.
- (1) Cherchi, M. A. Note su Rana obbiana (Calabresi). Natura, Milano 49 3 1958: 105-109.
- (2) Cherchi, M. A. Note su Chiromantis petersi kelleri Boettger e sui suoi nidi (Amphibia). Atti Soc. ital Sci. nat. 97 1958: 167-172 2 pls.
- (3) Cherchi, M. A. Note su Bufo gardoensis Scortecci. Boll. Mus. Ist. biol. Genova. 28 1958: 107-110.
- Chernov, S. A. [Systematic review of the animals of Crimea. III. Reptilia and Amphibia.] Zhivot. Mir 5 1958: 73-78. [In Russian.]

Chin, K.-C., see Chu, H.-W.

Chiuini, F. Azione della streptomicina sullo sviluppo e differenziazione di uova e larve di *Bufo vulgaris*. Riv. Biol. **50** 1958: 59-62 l fig. [English summary.]

Chrapliwy, P. S., see Smith, H. M.

Chu, H.-W. & Chin, K.-C. Histochemical studies on the distribution of phosphatases and mucopolysaccharides in the oviduct of the toad Bufo (B. gargarizane) after the laying of eggs. Acta zool. sinica 10 1958: 341-346 1 pl.

Claes, H., see Wessing, A.

Claracq, P., see Lapparent, A. F.

Clark, H. F., see Adams, M. S.

- Clarke, R. F., Breukelman, J. & Andrews, T. F. An annotated check list of the vertebrates of Lyon County, Kansas. Trans. Kansas Acad. Sci. 61 1958: 165–194.
- Cohen, N. W. & Howard, W. E. Bullfrog food and growth at the San Josquin Experimental Range, California. Copeia 1958 No. 3: 223-225 3 figs.

Collenot, A., see Gallien, L.

Colom, G. Biogeografia de las Baleares, Palma de Mallorca, 1957: 1–568 figs.

- (1) Conant, R. Notes on the herpetology of the Delmarva Peninsula. Copeia 1958 (1): 50-52.
- (2) Conant, R. A field guide to the reptiles and amphibians of the United States and Canada east of the 100th Meridian. Boston (Houghton Mifflin Co). 1958 pp. xv (ii) 366, 40 pls., text illust.

Conant, R., see Adler, K. K.

Concha, J., see Springmuller, V.

Condit, J. M. True type locality of the salamander Gyrinophilus perphyriticus inagnoscus. Copeia 1958 (1): 46-47.

Cooper, J. E. Some albino reptiles and polydactylous frogs. Herpetologica 14 1958: 54-56.

Cooperstein, I. L., see Chalfin, D.

- (1) Cordier, D. & Duveau, A. Action de l'anoxie et de l'asphyxie sur la phosphorolyse du glycogène musculaire chez la Grenouille (Rana esculenta). C.R. Soc. Biol., Paris 152 1958: 84-87.
- (2) Cordier, D. & Duveau, A. Troubles réversibles de la phosphorolyse du glycogène hépatique provoqués par l'asphyxie chez la Grenouille (Rana esculenta). C.R. Soc. Biol. Paris 152 1958: 1129-1131.
- Cordier, D., Duveau, A. & Brandon, A. M. Troubles de la phosphorolyse du glycogène hépatique chez la Grenouille (*Rana esculenta*) sous l'influence de l'anoxie et de l'asphyxie. C.R. Soc. Biol., Paris **151** 1958: 1524—1528.
- (1) Coromaldi, L. De C. Studio delle dipeptidasi negli embrioni di Anfibi trattati con raggi X. Arch. Sci. Biol. Bologna 43 1958: 205-212 3 figs.
- (2) Coromaldi, L. de C. Studio delle ribonucleasi nello sviluppo embrionale, larvale e nella metamorfosi di Bufo vulgaris. R. C. Ist. lombardo 92B 1958: 357-378 10 figs.

Crescitelli, F. The natural history of visual pigments. Ann. N.Y. Acad. Sci. 76 1958: 230–255 3 figs.

- Cruz, A. R. Sur l'existence de deux systemes porte dans l'hypophyse des amphibiens anoures. C.R. Acad. Sci., Paris 242 1956: 189–190.
- Csapo, A. & Suznki, T. The effectiveness of the longitudinal field, coupled with depolarization in activating frog twitch muscles. J. gen. Physiol. 41 1958: 1083-1098 3 figs.

Pad 97-1

[1

ther. Trit 818-

C: surf Zoo

C

visu J. P n'al

fem

la G

Biol D limil and 195

D Tar Pro

des 7 19 Lac

alpe Opt

effe viri Wo

J. 1

sen fro

act exp

1

Curry-Lindahl, K. Vertebratfauna i Sareks och Padjelantas fjällområden. Fauna och Flora 53 1958 : 97–149 2 figs. [English summary.]

Curtis, A. S. G. A ribonucleoprotein from amphibian gastrulae. Nature, Lond. 181 1958: 185.

Cury, G. & Dollander, A. Action d'un gradient thermique sur le développement embryonnaire de Triturus alpestris. C.R. Soc. Biol. Paris 152 1958 : 818-820

Cury, G., see Brouant, J. F.

Czopek, J. The vascularization of respiratory surfaces in *Ambystoma mexicanum* (Cope) in ontogeny. Zool. Polon., **8** 1957 [1958]: 131-149 6 figs.,

Darlington, P. J., see Savage, J. M.

Dartnall, H. J. A. Further observations on the visual pigments of the clawed toad Xenopus laevis. J. Physiol. 134 1956: 327-338.

Dechambe, C. L'ablation précoce du pronéphros n'altère pas le développement du pavilion génital femelle (ostium). Démonstration expérimentale chez la Grenouille agile (Rana dalmatina Bon.). C.R. Soc. Biol. Paris 152 1958: 1122-1124.

Dechambe, C., see Cambar, R.

Deck, J. D. & Riley, H. L. Regenerates on hindlimbs with reversed proximo-distal polarity in larval and metamorphosing urodeles. J. exp. Zool. 138 1958: 493-504 l fig.

Deck, J. D., see Malinin, T.

De Lanney, L. E. The development of the spleen of Taricha torosa and its experimental modification. Proc. Indiana Acad. Sci. 67 1958: 302.

Degan, C. & Asandei, A. La régénération de la musculature striée de *Triturus cristatus* sous l'action des vitamines A, B, C et PP. Stud. Cercet. Biol. Cluj 7 1958: 151-158 7 figs. [French summary.]

**Delleman, K.** Aantekeningen over de Heikikvors. Lacerta 15 11–12 1957: 92–93.

**Dely, O. G.** Les nouveaux habitats du Tritone alpestre (*Triturus alpestris* [Laurenti]) en Hongrie. Opusc. Zool. **2** 4 1958: 19-25 map.

Denis, H. Influence de la méthode d'élevage sur la différenciation nerveuse d'un fragment d'ectoblasti. J. Embryol. exp. Morph. 6 1958: 444-447.

Dent, J. N. & Lynn, W. G. A comparison of the effects of goitrogens on thyroid activity in *Triturus viridescens* and *Desmognathus fuscus*. Biol. Bull. Woods Hole 115 1958: 411-420 7 figs.

Denton, E. J. & Wyllie, J. H. Study of the photo sensitive pigments in the pink and green rods of the frog. J. Physiol. 127 1955: 81-89.

Detwiler, S. R., see Van Dyke, R. H.

Deuchar, E. M. Regional differences in catheptic activity in *Xenopus laevis* embryos. J. Embryol. exp. Morph. 6 1958: 223-237 5 figs.

Diener, R. A., see Alexander, D. R.

Dluhy, R. G., see Fankhauser, G.

Dobrowolski, J. A., Johnson, B. K. & Tansley, K. The spectral absorption of the photopigment of *Xenopus laevis* measured in single rods. J. Physiol. 130 1955: 533-542 6 figs.

**Dollander, A.** Effets du retournement de l'oeuf de *Triturus alpestris* sur le gradient de perméabilité corticale. C.R. Soc. Biol. Paris **152** 1958 : 982–984.

Dollander, A., see Cury, G.

Doneff, D., see Locker, A.

Doss, N. S., see Huf, E. G.

Dowling, H. G. Geographic relations of Ozarkian amphibians and reptiles. Southwest, Nat. 1 4 1956 [1958]: 174-189 3 maps 1 chart.

Dreyfus, E., see Viaud, G.

Droin, A., see Kiortsis, V.

Dubedout, C. Note sur une maladie regionale des Batraciens Le 'red leg ". Bull. Centr. Ét. Rech. sci. 2 1958: 7-8.

Dudich, E. Die Grundlagen der Fauna eines Karpaten-Flusses. Acta zool., Budapest 3 1958: 179-200.

(1) Duellman, W. E. A preliminary analysis of the herpetofauna of Colima, Mexico. Occ. Pap. Mus. Zool. Univ. Mich. No. 589 1958: 1–22 2 maps.

(2) Duellman, W. E. A review of the frogs of the genus Syrrhophus in Western Mexico. Occ. Pap. Mus. Zool. Univ. Mich. No. 594 1958; 15 pp. 3 pls.

Duellman, W. E. & Schwartz, A. Amphibians and reptiles of Southern Florida. Bull. Fla St. Mus. 3 5 1958: 181–324 28 figs.

Duguy, R. & Philippe-Knoepffler, L. Reptiles et batraciens des environs de Banyuls. Vie et Milieu 9 1958: 129-131.

Durbin, R. P., Frank, H. & Solomon, A. K. Water flow through frog gastric mucosa. J. gen. Physiol. 39 1956: 535-551 4 figs.

Durbin, R. P. & Heinz, E. Electromotive chloride transport and gastric acid secretion in the frog. J. gen. Physiol. 41 1958: 1035-1047 7 figs.

Durbin, R. P., see Heinz, E.

Duveau, A., see Cordier, D.

Dzieduszycka, S., see Lamotte, M.

Eaton, T. H., Jr. An anatomical study of a neotropical tree frog, Centrolene prosoblepon (Salientia: Centrolenidae). Univ. Kans. sci. Bull. 39 1958: 459-472 20 figs.

Eckhardt, E. T. & Govier, W. M. Chlorpromazine blockade of water inhibition by frog gastroenemius. Proc. Soc. exp. Biol. & Med. 97 1958: 124-127.

Edds, M. V., Jr. Development of collagen in the frog embryo. Proc. nat. Acad. Sci. Wash. 44 1958: 296-305 3 figs.

Eiselt, J. Der Feuersalamander, Salamandra salamandra (L.), Beiträge zu einer taxonomischen Synthese. Abh. Ber. Naturk. u. Vorgesch. Magdeburg. 10 6 1958: 77–154 3 pls., 3 maps.

Eksaeva, V. A. The histological structure of the esophagus of certain cold-blooded vertebrates. Dokl. Akad. Nauk SSSR (Transl.) Biol. Sci. 118 1958: 42–46 3 figs. [See also: C.R. Acad. Sci. U.R.S.S. 118 1: 181–184 4 figs. In Russian.]

Elbel, R. E., see Taylor, E. H.

- (1) Elkan, E. and pharyngeal mucous membrane in urodeles. Proc. zool. Soc. Lond. 131 1958; 335–355 1 pl. 12 figs.
- (2) Elkan, E. The cave salamander (Hydromantes Gistel). Brit. J. Herpet. 2 1958: 98-104, 1 pl., 2 figs.
- (3) Elkan, E. The cave salamander (Hydromantes Gistel). Brit. J. Herpet. 2 1958: 133-134.

Elsdale, T. R., Fischberg, M. & Smith, S. A mutation that reduces nucleolar number in *Xenopus laevis*. Exp. Cell. Res. **14** 1958: 642-643.

Elsdale, T. R., see Fischberg, M.

Elsdale, T. R., see Gurdon, J. B.

England, M. C. & Mayer, D. T. The comparative nucleic acid content of liver nuclei, ova nuclei and spermatozoa of the frog. Exp. Cell. Res. 12 1957: 249-253.

**Erspamer, V.** Isolamento della leptodactylina (m-idrossifenil-etil-trimetilammonia) degli estratti di pelle di Leptodactylus. Ric. sci. **28** 1958 : 2065–2069. [English etc. summaries.]

**Eyal-Giladi, H.** The notochord as inductor of the orohypophysis in urodeles (*Pleurodeles walthii*). Proc. Akad. Wet. Amst. **61C** 1958: 224–234 2 pls., 1 fig.

Fabian, N., see Wittenberger, C.

Fankhauser, G. & Dluhy, R. G. Relative constancy of somatic chromosome numbers in larvae of the salamanders Triturus viridescens, Pleurodeles waltlii and Eurycea bislineata. Proc. Xth internat. Congr. Genetics 2 1958: 80.

Fendler, R. Eine neue flexible Abguss-Masse. Präparator 1 1955: 8-9 1 fig.

Ferguson, D. E., Payne, K. E. & Storm, R. M. Notes on the herpetology of Baker Co., Oregon. Gt. Basin Nat. 18 1958: 63-65.

Ferguson, D. E. & Rhodes, J. R. A new locality for the zigzag salamander in Mississippi. Herpetologica 14 1958: 129.

Feriancová, Z. Bemerkungen zu der Verbreitung und Ökologie von Amphibien auf Reisfeldern der Slowakei. Biológia, Bratislava 13 1958: 26–38. [German summary.]

Filatova, A. G., see Tokin, B. P.

Filippio, S. De. Osservazioni sulla citomorfosi delle Cellule di Rohon-Beard negli Anfibi anuri. Ric. sei. 28: 967–972 3 figs. [English etc. summaries.]

Finamore, F. J. A study of a contaminent in amphibian egg extracts which interferes with the diphenylamine reaction for deoxyribonucleic acid. Exper. Cell. Res. 12 1957: 356-362 6 figs.

Finamore, F. J. & Volkin, E. Nucleotide and nucleic acid metabolism in developing amphibian embryos II. Composition and metabolic activity of ovarian egg nucleic acids. Exper. Cell. Res. 15 1958: 405-411.

Finnegan, C. V. The influence of the hypomere on chromatoblast migration in salamanders. J. exp. Zool. 138 1958: 453-491 31 figs.

Fischberg, M. Experimental tetraploidy in newts. J. Embryol. exp. Morph. 6 1958: 393-401, 1 pl.

Fischberg, M., Gurdon, J. B. & Elsdale, T. R. Nuclear transplantation in *Xenopus laevis*. Nature, Lond. **181** 1958: 424.

Fischberg, M., see Elsdale, T. R.

Fischberg, M. see Gurdon, J. B.

Fitch, H. S. Home ranges, territories, and seasonal movements of vertebrates of the natural history reservation. Univ. Kans. Publ. Mus. nat. Hist. 11 1958: 63-226 6 pls., 24 figs.

Fitch, K. L. Induced ovulation in Necturus Copeia 1958 No. 2: 151-152.

Fitch, K. L. & Merrick, A. W. The glycogen content of ovarian, body cavity, and uterine eggs of the frog, Rana pipiens. Exper. Cell. Res. 14 1958: 644-646.

Fitzimons, V. On a small collection of reptiles and amphibians from the Inyanga district, S. Rhodesia. Occ. Pap. nat. Mus. S. Rhod. 3 1958: 204-214 1 fig.

Fleming, W. R. The role of the corium in ion transport across the isolated frog skin. J. cell. comp. Physiol. 51 1958: 189-198 2 figs.

Flickinger, R. A. Regional localization of neural and lens antigens in the frog embryo in relation to induction. Biol. Bull. Woods Hole 115 1958: 201-208.

Flickinger, R. A. & Blount, R. W. The relation of natural and imposed electrical potentials and respiratory gradients to morphogenesis. J. cell. comp. Physiol. 50 1957: 403-422.

Flickinger, R. A. & Schjeide, O. A. The localization of phosphorus and the site of calcium binding in the yolk protein of the frog's egg. Exper. Cell. Res. 13 1957: 312-316.

Flickinger, R. A., see Rounds, D. E.

- (1) Foote, C. L. & F. M. Maintenance of gonads of frog larvae in organ cultures. Trans. Ill. Acad. Sci. 50 1958: 243-246 1 fig.
- (2) Foote, C. L. & F. M. Maintenance of gonads of frog larvae on synthetic media. Proc. Soc. exp. Biol. Med. 98 1958: 750-753.
- (3) Foote, C. L. & F. M. In vitro cultivation of gonads of larval anurans. Anat. Rec. 130 1958: 553-561 2 pls.

Foote, C. L., see Witschi, E.

Foote, F. M., see Foote, C. L.

Fouquette, M. J. A new tree frog, genus Hyla, from the Canal Zone. Herpetologica 14 1958: 125-128 2 figs.

Fowler, J. A., see Stine, C. J.

Bioch Fra Fre

[19

Fra

Pseud logica Fre

Frè Sei, p

(Trite dem ]

(2)

verkr 108–1 (3)

Fre in eir 1958

> Fre Sci. 5

Ambl 117-1 Fue fauna

Ful Curai frog. 1 fig.

de ce Paris Gal

(1) porta e Inv (2)

Lepto 14 19 (1) chez Biol.,

(2) dével verté walth 1 fig.

(fémir fémir chez ment penda 1958 Frank, G. B. The effects of trematode infection on the physiological activity of the frog. Can. J. Biochem. Physiol. 35 1957: 347-356 2 figs.

Frank, H., see Durbin, R. P.

Freeman, J. R. Burrowing in the salamanders Pseudobranchus striatus and Siren lacertina. Herpetologica 14 1958: 130.

Freiberg, M. A., see Oliver, J. A. (1)

Frèrejacque, M. Venins de crapauds. Rev. gén. Sci. pur. appl. 65 1958 : 145–149.

- (1) Freytag, G. E. Ein weiterer Bergmolch (*Triturus alpestris alpestris*) mit Bauchzeichnung aus dem Herz, Zool, Anz, 161 1958: 85.
- (2) Freytag, G. E. Marmormolch mit Rückgratsverkrümmung. Zool. Garten (N. F.) 24 1958: 108–109 1 fig.
- (3) Freytag, G. E. Vom Temperaturhaushalt der Schwanzlurche. Aquar. u. Terrar. 5 1958: 140–142.
- Freytag, G. E. & Hübener, H. E. Drei Anomalien in einer Bergmolchzucht. Aquar. Terrar. Z. 11 3 1958: 82–83.

Freytag, G. E., see Bašoglu, M.

**Fromm, F.** Some constants of toad fat. Ohio J. Sci. **58** 1958: 161-162.

Frothingham, M. L., see Ruben, L. N.

Frye, B. E. Development of the pancreas in Amblystoma opacum. Amer. J. Anat. 102 1958: 117-139 2 pls. 2 figs.

Fudakowski, J. Remarks on the herpetological fauna of Poland. Acta zool. Cracov. 2 1958: 825–844.

Fulchiero, E., Turcotte, M. R. & Martin, S. J. Curarising effect of neomycin and its fractions in the frog. Proc. Soc. exp. Biol. Med. 99 1958: 537-539

Gabe, M. Caractères histochimiques des parasomes de certaines cellules glandulaires. C.R. Acad. Sci., Paris 247 1958: 1907–1909.

Gallagher, S. B., see Stone, L. S.

of

- (1) Gallardo, J. M. Observaciones sobre el comportamiento de algunos anfibios argentinos. Cienc. e Invest. 14 1958; 291-302 l fig.
- (2) Gallardo, J. M. Observaciones biologicas sobre Leptodactylus prognathus Boulenger. Cienc. c Invest. 14 1958: 460–465 3 figs.
- (1) Gallien, L. Parabiose de larves teratologiques chez le Triton *Pleurodeles wattii* Michah. C.R. Soc. Biol., Paris 151 1958: 1085–1087 7 figs.
- (2) Gallien, L. Production expérimentale et développement d'un Vertébré dépourvu de colonne vertébrale. Cas de l'Amphibien Urodèle Pleurodeles waltii. C.R. Acad. Sci. Paris 247 1958: 1394-1396 l fig.
- Gallien, L. & Collenot, A. Inversion totale (féminisation) du sexe chez Pelobates cultripes Cuv. et féminisation partielle des gonades (intersexualité) chez Pelobates punctatus Daud., à la suite d'un traitement gynogène par le benzoate d'oestradiol, administré pendant la vie larvaire. C.R. Acad. Sci., Paris 247 1958: 1042-1044.

- Gans, C. Terrarienkunde 2. Teil Lurche by W. Klingelhöffer. [Review.] Copeia 1958 1:63.
- Gaze, R. M. The representation of the retina on the optic lobe of the frog. Quart. J. exp. Physiol. 43 1958: 209-214 3 figs.
- Gerbes, L. Über den Einfluss der Temperatur auf die Entwicklung des Krallenfrosches (Xenopus laevis Daud). Zool. Jb. (Allg. Zool.) 67 1958: 373-406, 21 figs.
- Germaine, L. A. P. La faune des lacs, des étangs et des marais de l'Europe occidentale. 2me édition revisée...par E. Séguy. Encyclopédie pratique du Naturaliste, Tome XX. Paris (Editions Paul Lechevalier) 1957, pp. x, 549: 32 col. pls. test illust. 418-444.
- Getz, L. L. The winter activities of Rana clamitans tadpoles. Copeia 1958 No. 3: 219.
- Geyer, G. Histotopochemische Untersuchungen mit der Hale-PAS-Reaktion ander Niere von Rana esculenta. Acta histochem. 5 1958: 1-9 1 pl.
- (1) Geyer, H. Rana japonica (Guenther). Aquar. Terrar, Z. 11 1958: 25-26.
- (2) Geyer, H. Kleinlaubfrösche der Vereinigten Staaten. Aquar. Terrar. Z. 11 1958: 276–279.
- Giang, H.-C. & Li, J.-C. The study of the development of *Kaloula borealis* (Barbour). I. The origin, development and changes of the gills and lungs. Acta zool. Sinica 10 1958: 35–52 5 pls. [In Chinese: English summary.]
- Gianotti, F. S. Mortalità delle rane verdi (Rana esculenta L.) dopo i trattamente insetticide lungo le sponde del Lago Trasimeno. Riv. Biol. Perugia 50 1958: 269–278 4 pls. [Italian with English summary.]
- **Giebisch, G.** Electrical potential measurements on single nephrons of *Necturus*. J. Cell. comp. Physiol. **51** 1958: 221–239 2 figs.
- Gines, H. Representantes de la familia Pipidae (Amphibia, Salientia) en Venezuela. Mem. Soc. Cienc. nat. La Salle 18 49 1958 : 5–18 12 figs.

Gipouloux, J.-D., see Cambar, R.

Glauert, L., see Serventy, D. L.

- (1) Goin, C. J. Comments upon the origin of the herpetofauna of Florida. Quart. J. Fla. Acad. Sci. 21 1958: 61-70.
- (2) Goin, C. J. Notes on the maxillary dentition of some hylid frogs. Herpetologica 14 1958: 117-121
- Goin, C. J. & Auffenberg, W. New salamanders of the family Sirenidae from the Cretaceous of North America. Fieldiana Geol. 10 33 1958: 449-459 3 figs.
- Goin, C. J. Layne, J. N. Notes on a collection of frogs from Leticia, Colombia, Publ. Res. Div. Ross Allen Rept. Inst. 1 8 1958: 97-114 4 photos.
- Goin, O. B. A comparison of the nonbreeding habits of two treefrogs, Hyla squirelli and Hyla cinerea. Quart. J. Fla. Acad. Sci. 21 1958: 49-60.

Gonzales, R. B., see Rabor, D. S.

Gooch, B. Amphibians (in Report on Dorset Natural History for 1957). Proc. Dorset nat. Hist. arch. Soc. 79 1958: 31.

Goodman, J. D. Anuran records from the Upper Coastal Plain of western and central Georgia. Herpetologica 14 1958: 123-124.

- (1) Gosner, K. L. & Black, I. H. Notes on the life history of Brimley's chorus frog. Herpetologica 13 1958: 249-254 1 fig.
- (2) Gosner, K. L. & Black, I. H. Notes on larval toads in the Eastern United States with special reference to natural hybridization. Herpetologica 14 1958: 133-140, 2 figs.

Gosner, K. L., see Black, I. H.

- Goss, R. J. Skeletal regeneration in amphibians. J. Embryol. exp. Morph. 6 1958: 638-644 1 pl. 1 fig.
- (1) Goss, R. J. & Stagg, M. W. Regeneration in lower jaws of newts after excision of the intermandibular regions. J. exp. Zool. 137 1958: 1-11, 15 figs.
- (2) Goss, R. J. & Stagg, M. W. Regeneration of lower jaws in adult newts. J. Morph. 102 1958: 289-309, 19 figs.
- Goto, M. & Hama, T. Über die fluoreszierenden Stoffe aus der Haub eines Frosches Rana nigromaculata I. Isolierung und Eigenschaften des "Ranachrom. 1." Proc. Japan Acad. 34 1958: 724–729
- Gourley, D. R. H. Relation between structure of sulfomylurea compounds and their effect on frog muscle metabolism. Proc. Soc. exp. Biol. Med. 99 1958: 67-72.
- Goux, L. Contribution à l'étude écologique biologique et biogéographique de *Chioglossa lusitanica* Barb. (Urodela Salamandridae). Bull. Soc. zool. Fr. 82 1958 : 361–377.

Govier, W. M., see Eckhardt, E. T.

Grabie-Broda, C. Der Einfluss von methylthiouracil auf die Haematopoesis bei Anura. Zool. polon. 8 1958: 65-86 8 figs 3 pls.

- (1) Grandison, A. G. C. Dr. Malcolm A. Smith. [Obituary.] Nature Lond. 182 1958: 697.
- (2) Grandison, A. G. C. The Amphibia of Ceylon by P. Kirtisinghe. [Review.] Nature Lond. 181 1958: 1758.
- Grant, C. Irruption of young Batrachoseps attenuatus. Copeia 1958 No. 3: 222.

Grant, C., see Smith, H. M.

Grant, J. A., see Grant, W. C.

- Grant, W. C. & Grant, J. A. Water drive studies on hypophysectomized efts of Diemyctylus viridescens. Part I. The role of the lactogenic hormone. Biol. Bull. Woods Hole 114 1958: 1-9 1 fig.
- Gray, E. G. The structures of fast and slow muscle fibres in the frog. J. Anat., Lond. 92 1958: 559-562, 1 pl. 1 fig.
- Grenard, S. Life history of Gastrotheca marsupiata spp. Herpetologica 14 1958: 151-152, 1 fig.

- Gribenski, A. Sur le sens des reactions provoquées par la stimulation d'un canal semi-circulaire horizontal chez la Grenouille (*Rana viridis*). J. Physiol. Paris 46 1954: 377–379.
- Griffiths, I. & Carter, M. E. Sectioning refractory animal tissues. Stain Technology 33 1958: 209-214.
- (1) Gualtierotti, T. The contribution of spinal centres to the action of caffeine on frog's spinal reflexes. J. Physiol. 128 1955: 326-332 4 figs.
- (2) Gualtierotti, T. Variations in the frog's spinal reflexes caused by the action on the brain of large doses of caffeine. J. Physiol. 128 1955: 320–325 2 figs.
- (1) Guibé, J. & Lamotte, M. Les Ptychadena (Batraciens, Ranidés) du Cameroun. Bull. Inst. franç, Afr. noire 20A 1958: 1448–1463 5 figs.
- (2) Guibé, J. & Lamotte, M. Merphologie et reproduction par développement direct d'un Anoure du Mont Nimba, Arthroleptis crusculum Angel. Bull. Mus. Hist. nat. Paris (2) 30 1958: 125-133 4 figs.
- (3) Guibé, J. & Lamotte, M. Une espèce nouvelle de Batracien du Mont Nimba (Guinée française) appartenant au genre *Phrynobatrachus*: *Ph. macutiventris* n. sp. Bull. Mus. Hist. nat. Paris (2) 30 1958: 255-257 1 fig.
- (4) Guibé, J. & Lamotte, M. La réserve naturelle intégrale du Mont Nimba. Batraciens (auf Arthrolepites, Phrynobatrachus et Hyperolius). Mém. Inst. franç. Afr. N. No. 53 1958: 241-273 11 pl. 32 figs.

Guiguet, C. J., see Carl, G. C.

Gunter, G., see Brode, W. E.

Gupta, B. L., see Nath, V.

Gurdon, J. B., Elsdale, T. R. & Fischberg, M. Sexually mature individuals of Xenopus laevis from the transplantation of single somatic nuclei. Nature, Lond. 182 1958: 64-65.

Gurdon, J. B., see Fischberg, M.

Gut, H. J., see Neill, W. T.

- **Guyénot, E.** L'évolution des nucléoles dans les ovocytes de batraciens. Rev. suisse Zool. **65** 1958 : 879–889 5 figs.
- Gyévai, A. T. Herzstudien. I. Histologie und histochemie des Herzens von erwachsenen Triturus cristatus II. Untersuchungen von Überlebenden Herzabschnitten des Triturus cristatus. Acta biol., Budapest 8 1958; 333–346 5 figs. [English. etc, summaries.]
- Hadler, W. A., Berezin, A. & Lison, L. L'action de la pitressine et de l'adrénaline sur le bilan hydrique chez *Bufo paracnemis* L. pendant et après la métamorphose. J. Physiol. Paris **50** 1958: 785–797.
- Hahn, D. E. New county records from southern Michigan. Trimon. Rep. Ohio herpet. Soc. 1.4 1958: 19.
- Hahn, H. P. van. Die regionale Verteilung der Katalase-Aktivität im Schwanz von Rungernden und gefütterten Xenopus larven. Experientia 14 1959: 67-68

Hasubs dans Soc.

[1

fluor of to 182-

(1) upon J. F. 1 fig

propretari 413-

(2

genu 3–4 1 Hs spad

Ha frog 8 figs

H

Ha the p isolate 148– (1)

(2) and limb Cyto

60-6 (2) easte Sci.

(1)

halla porficione by I Cope

No.

Phys

a t.

m

1

le

6)

:

le

m re,

nd rus len

ol., m-

ion

ta-

ern

8 :

der

59 :

- Hama, T. & Horiuchi, S. Sur la localisation des substances fluorescentes ptériniques et flaviniques dans l'oeil de la Grenouille Rana nigromaculata. C.R. Soc. Biol., Paris 152 1958: 213-216.
- Hama, T. & Obika, M. On the nature of some fluorescent substances of pterin type in the adult skin of toad Bufo vulgaris formosus. Experientia 14 1958: 182-184 2 figs.

Hama, T., see Goto, T.

Hammen, C. S., see Hutchison, V. H.

- (1) Hanaoka, K. The effect of follicular hormone upon the sex differentiation in *Hynobius retardatus*. J. Fac. Sci. Hokkaido Univ. 7 1940: 399-412 2 pls. 1 fig.
- (2) Hanaoka, K. The effect of testosterone-propionate upon the sex differentiation in *Hynobius retardatus*. J. Fac. Sci. Hokkaido Univ. 7 1940: 413-419 2 pls.

Hanke, W., see Bücker, H.

- Hansen, A. M. & Tanner, W. W. A comparative osteological study of certain species belonging to the genus Bolitoglossa (Amphibia). Gr. Basin Nat. 18 3-4 1958: 85-100 30 figs.
- Hansen, K. L. Breeding pattern of the eastern spadefoot toad. Herpetologica 14 1958: 57-67.

Harell-Steinberg, A., see Lunenfeld, B.

- Harris, E. J. Permeation and diffusion of K ions in frog muscle. J. gen. Physiol. 41 1957: 169-195 8 figs.
  - Harvey, S. M., see Volpe, E. P.
- Hashimoto, K., Morita, Y. & Matsuyama, S. On the peculiarity of the carbohydrate metabolism of the isolated bullfrog's heart. Jap. J. Physiol. 8 1958: 148-154 5 figs.
- (1) Hay, E. D. Effects of thyroxine on limb regeneration in the newt, *Triturus viridescens*, John Hopk, Hosp. Bull. 99 1956: 262-285. [Not seen.]
- (2) Hay, E. D. The fine structure of blastema cells and differentiating cartilage cells in regenerating limbs of Amblystoma larvae. J. biophys. & biochem. Cytol. 4 1958: 583-592 7 pls. 1 fig.
- (1) Hecht, M. K. Vertebrates of the United States by W. F. Blair and others [Review.] Copeia 1958 1: 60-61.
- (2) Hecht, M. K. A synopsis of the mud puppies of eastern North America. Proc. Staten Id. Inst. Arts Sci. 21 1 1958: 5-38 1 fig. 2 maps.
- Hecht, M. K. & Ruibal, R. Breve notice sobre el hallazgo de anuros en el denominado "complejo porfirico de la Patagonia extraandina" con consideraciones acerca de la composicion geologica del mismo by P. N. Stipanincic and O. A. Reig. [Review.] Copeia 1958 3: 242.
- Heilingbrunner, F. Lurche und Kreichtiere in der Umgebung von Bamberg. Ber. naturf. Ges. Bamberg No. 36 1958: 42-50.
- Heinz, E. & Durbin, R. P. Studies on the chloride transport in the gastric mucosa of the frog. J. gen. Physiol. 41 1957: 101-117 8 figs.

- Heinz, E., see Durbin, R. P.
- Heller, F. Eine neue altquartäre Wirbeltierfauna von Erpfingen (Schwäbische Alb). Neues Jb. Geol. Paläont. Abh. 107 1958: 1–102 3 pls. figs.
- (1) Hellmich, W. Vom Vorkommen des Feuersalamanders in Tirol. Natura Tiroliensis Schlern Jehr. No. 188 1958: 167–170.
- (2) Hellmich, W. Nemrut Dag und Suphan Dag in Ostanatolien. Mitt. Deutsch. Alpenvereins 10 2: 24-26 photos.
- Findricks, L. J. & Kezer, J. An unusual population of a blind cave salamander and its fluctuation during one year. Herpetologica 14 1958: 41-43.
- Hensley, M. A long lived newt, Triturus pyrrhogaster. Herpetologica 14 1958: 202.
  - Hensley, M. M., see Smith, P. W.
- Hertwig, G., Weiss, I. & Ziemann, C. Unterschiedliche Ergebnisse reziproker Kreuzungen der drei europäischen Krötenarten unter besonderer Berücksichtigung ihrer Kern-und Zellgrössen. Biol. Zbl. 78 1958: 675–702 5 pls., 17 figs.,
- Hertwig, G., & Others. Die Kerngrössen triploider Krötenbastarde mit experimentell verdoppeltem mütterlichen Genom. Z. mikr.-anat. Forsch. 64 1958: 129-158 11 figs.
- Hertzler, E. C. The red and black pigments of the salamander, *Plethodon cinereus*. Physiol. Zoöl. 31 1958: 248-256 4 figs.
- (1) **Heusser, H.** Ueber die Beziehungen der Erdkröte (*Bufo bufo L.*) zu ihrem Laichplatz I. Behaviour **12** 1958 : 208–232 4 figs.
- (2) Heusser, H. Zum Häutungsverhatten von Amphibien. Rev. suisse Zool. 64 1958: 793-823.
- (3) Heusser, H. Zum geruchlichen Beutefinden und Gähnen der Kreuzkröte (Bufo calamita Laur.). Z. Tierpsychol. 15 1958: 94–98, 3 figs.
- (4) Heusser, H. Markierungen an Amphibien. Vjschr. naturf. Ges. Zurich 103 1958: 304-320 13 figs.
- Hobson, B. M. Polymely in Xenopus laevis. Nature, Lond. 181 1958: 862 2 figs.
- Hodler, F. Untersuchungen über den Crowd-Effekt an Kaulquappen von Rana temporaria L. Rev. suisse Zool. 65 1958: 350-359 3 figs.
- Hoffmann, J. Sur la présence de Pelodytes punctatus Daudin au Grand-Duché de Luxembourg. Arch. Inst. Grand-Duc. Luxemb. Sect. Sci. N.S. 25 1958: 235-237.
  - Hogben, C. A. M., see Chalfin, D.
- (1) Holman, J. A. The Pleistocene herpetofauna of Saber-tooth Cave, Citrus County, Florida. Copeia 1958 No. 4: 276-280.
- (2) Holman, J. A. Notes on reptiles and amphibians from Florida caves. Herpetologica 14 1958: 179–180.
- Holman, J. A. & Campbell, H. An interesting feeding activity of *Microhyla carolinensis*. Herpetologica 14 1958: 205.

Hopkins, A. L. The rehydration contraction of frozen dried muscle. J. cell. comp. Physiol. 51 1958: 67–80 7 figs.

Horiuchi, S., see Hama, T.

Hosmer, W. A note on the identity of Heleioporus sudelli Lamb (Amphibia—Leptodactylidae). N. Qd. Nat. 27 121 1958: 1-2.

Houssay, B. A., see Burgos, M. H.

Howard, W. E., see Cohen, N. W.

Hübener, H. H., see Freytag, G. E.

Hudson, R. G. Field trips to an unusual amphibian habitat. National Assoc. Biol. Teachers, Bryan, Ohio, 18 1956: 164–165. [Not seen.]

- (1) Huene, F. von. The first Chinese Labyrinthodonts. Vertebrata Palasiatica 2 1958: 101-105 figs.
- (2) Huene, F. von. Pre-Tertiary saurians of Asia. Vertebrata Palasiatica 2 1958: 201-207.
- Hut, E. G. & Doss, N. S. Effect of temperature on secretory activity and oxygen consumption of isolated frog skin. Va. J. Sci. N.S. 9 1958: 436.
- Huf, E. G., Doss, N. S. & Wills, J. P. Effects of metabolic inhibitors and drugs on ion transport and oxygen consumption in isolated frog skin. J. gen. Physiol. 41 1957: 397-417 4 figs.

Huf, E. G., see Skjelkvale, L.

Hughes, A. & Tschumi, P. A. The factors controlling the development of the dorsal root ganglia and ventral horn in *Xenopus laevis* (Daud.). J. Anat. Lond. 92 1958: 498-527 2 pls. 13 figs.

Humphrey, F. L., see Shannon, F. A.

Hurlbut, W. P. Effects of azide and chloretone on the sodium and potassium contents and the respiration of frog sciatic nerves. J. gen. Physiol. 41 1958: 959-988 12 figs.

Hurlbut, W. P., see Asano, T.

Hurley, M. P. The role of the thyroid in kidney development in the tadpole. Growth 22 1958: 125-166 5 pls.

Hutchison, V. H. The distribution and ecology of the cave salamander, *Eurycea lucifuga*. Ecol. Monogr. **28** 1958: 1–20 13 figs.

Hutchison, V. H. & Hammen, C. S. Oxygen utilization in the symbiosis of embryos of the salamander, Ambystoma maculatum and the alga, Oophila amblystomatis. Biol. Bull. Woods Hole 115 1958: 483-489 2 figs.

Hutter, O. F. & Loewenstein, W. R. Nature of neuromuscular facilitation by sympathetic stimulation in the frog. J. Physiol. 130 1955: 559-571 9 figs.

Ikushima, N. Kinetic properties of the marginal zone of the amphibian egg in relation to the histological differentiation. Mem. Coll. Sci. Univ. Kyoto 25B 1958: 145-160 11 figs.

(1) Inger, R. F. The vocal sac of the Colorado River toad (*Bufo alvarius* Girard). Texas J. Sci. 10 1958: 319-324 2 figs. (2) Inger, R. F. A new toad from Sarawak. Sarawak Mus. J. 8 1958: 476-479 1 fig.

(3) Inger, R. F. A note on the Philippine frogs related to Rana macrodon. Fieldiana Zool. 39 (23) 1958: 253-255.

Ioff, N. A. The process of wound healing and the problem of the integrated nature of the organism as exemplified by Amphibia. Dokl. Akad. Nauk SSSR (Transl.) Biol. Sci. 118 1958: 36-38 3 figs. [See also C.R. Acad. Sci. U.R.S.S. 118 6: 1200-1202. In Russian.]

Ishida, K. On the Schmidt-Lautermann's incisure of the peripheral medullated nerve fiber. Okajimas Folia anat. Jap. 31 1958: 285–296 4 figs.

- (1) Iwasawa, H. Différenciation anormale de la gonade du têtard provoquée par l'injection de la gonadtrophine hypophysaire. Zool. Mag., Tokyo 67 1958: 214–216 é figs. [In Japanese, French summerv.]
- (2) Iwasawa, H. Relation between the anterior lobe of hypophysis and the sex-reversal in thioureatreated frog larvae. Sci. Rep. Tôhoku Univ. (4) 24 1958: 55-58.

Jacob, J. & Pécot-Dechavassine, M. Hydrolyse enzymatique de la propionylcholine de l'acetylthiocholine par le rectus de grenouille. Experientia 14 1958 : 330–332.

Jacob, J., see Pantelouris, E. M.

Jacobson, A. G. The roles of neural and non-neural tissues in lens induction. J. exp. Zool. 139 1958: 525-553 4 figs. 2 pls.

Jakowska, S. & others. Radiobiology of the newt, Diemictylus viridescens. Hematological and histological effects of whole-body X irradiation. Zoologica, N.Y. 43 1958: 155-160 3 pls.

Jard, S., see Maetz, J.

Jenden, D. J. A quantitative interpretation of eserine-acetylcholine interaction in the frog rectus abdominis. J. cell. comp. Physiol. 51 1958: 309-323 3 figs.

Jenerick, H.P. The relations between prepotential resting potential and latent period in frog muscle fibres. J. gen. Physiol. 39 1956 773-787 8 figs.

Jewell, B. R. & Wilkie, D. R. An analysis of the mechanical components in frog's striated muscle. J. Physiol. 143 1958: 515-540 14 figs.

Jimbo, G. & Nagasawa, M. On the skin disease of frogs. Bull. exp. Biol. Japan 6 2 1956: 113-114 1 fig.

Johansson, B., see Biörck, G.

Johnson, B. K., see Dobrowolski, J. A.

**Joly, J.** Influence des basses températures sur le cycle alimentaire de quelques tritons français. Bull. Soc. 200l. Fr. **83** 1958 : 128–131.

Jordan, M. Regeneration of the endbrain in postmetamorphic Xenopus laevis. Folia biol., Kraków 6 1958: 103–116 7 figs.

Jergensen, C. B. & Nielsen, L. Effect of synthetic lysine-vasopressin on adenohypophysial activity in toads. Proc. Soc. exp. Biol. Med. 98 1958: 393-395.

Ju circu terre la ca Biol. Ju évide

[19

Jo

elect

23-3

Jul Supp artéri

Zool.

etc. s

Jul Tisser pulm Grend circul 1958

Modi

larva

Jul Actionic circu de la auxa

Ann. 1958

Jul Thom dans soluti alcali pulm 1132-Ka Laby

> Ka the l Inst. plts.

> > Ka

laevie

Ka acina Expe

Ka On r (Amb 10 fig

turtle

- Joynt, R. J. Micro-electrode studies of cerebellar electrical activity in the frog. J. Physiol. 144 1958: 23-37 9 figs.
- Jullien, A., Cardot, J. & Moulin, J. Effets sur la circulation pulmonaire des ions alcalins et alcalinoterreux appliques en solutions monocationiques dans la cavité du poumon chez la Grenouille. C.R. Soc. Biol., Paris 152 1958; 631-633.
- Jullien, A., Ripplinger, J. &. Cardot, J. Mise en évidence, chez la Grenouille, d'une substance cardiomotrice d'origine nerveuse. C.R. Soc. Biol., Paris 151 1958: 1926–1929.
- Jullien, A., Ripplinger, J., Cardot, J. & Baylé, J. D. Suppressions partielles de la circulation par ligatures artérielles chez la grenouille et conséquences sur la survie de l'animal. Ann. Sci. Univ. Besançon (2) Zool. et Phys. No. 10 1958: 91–165 38 figs. [English etc. summaries.]
- Jullien, A., Ripplinger, J., Cardot. J., Baylé, J. & Tisserand, M. Influence de la ligature des troncs pulmo-cutanés et de leurs branches sur la survie de la Grenouille et possibilité de rétablissement de la circulation après arrêt. C.R. Soc. Biol., Paris. 152 1958 : 1161-1163.
- Jullien, A., Ripplinger, J., Cardot. J. & Marcot, M. Modifications de l'automatisme cardiaque chez des larves de *Triton alpestris* placées dans des solutions sodiques, potassiques ou calciques. C.R. Soc. Biol., Paris 151 1958: 1556–1558.
- Jullien, A., Ripplinger, J., Cardot, J. & Moulin, J. Action des solutions alcalines (Na<sup>+</sup> et K<sup>+</sup>) et alcalino-terreuses (Ca<sup>++</sup> et Mg<sup>++</sup>) mono-et bicationiques, en application interne et externe sur la circulation pulmonaire chez la Grenouille. Influence de la variation du rapport des alcalins et des alcalines auxalcalino-terreux. Action du vague et des droques Ann. Sci. Univ. Besançon (2) Zool. et Phys. No. 10 1958: 3-72 19 figs. [English etc. summaries.]
- Jullien, A., Ripplinger, J., Cardot, J., Moulin, J. & Thomas, R. Sur les modifications de la circulation dans le poumon de Grenouille provoquées par les solutions bicationiques d'ions alcalins (Na\* et K\*) et alcalino-terreux (Ca\*\*) introduites dans la cavité pulmonaire. C.R. Soc. Biol., Paris 152 1958: 1132-1134.
- Kahmann, H. & Frettlöh, U. Untersuchung am Labyrinth des Afrikanischen Krallenfrosches *Xenopus laevis* (Daudin, 1803). Zool. Beitr., Berl., 3 (2) 1958: 195–226 12 figs.
- Kankave, V. L. [Investigation into the capacity of the lens forming epithelium in amphibia.] Trud. Inst. Zool. Akad. Nauk Ghruzinsk. SSR 16 157-208 4 plts.
  - Kaur, R., see Nath, V.

of

13

al

le

e.

of

g.

le

11.

in

tio

in

15.

- Kanwar, K. C. What constitutes a network in the acinae cells of Amphibian and Mammalian Pancreas? Experientia 14 1958: 403–404.
- Kaplan, H. M. Marking and banding frogs and turtles. Herpetologia 14 1958: 131-132.
- Kappers, J. A., Ten Kate, I. B. & Bruyn, H. T. de. On mast cells in the choroid plexus of the axolotl (*Ambystoma mex.*). Z. Zellforsch. **48** 1958: 617-634 10 figs.

- Karasaki, S. & Komoda, T. Electron micrographs of a crystalline lattice structure in yolk platelets of the amphibian embryo. Nature, Lond. 181 1958: 407–408 2 figs.
- Karlstrom, E. L. Sympatry of the Yosemite Western toads in California. Copeia 1958 No. 2: 152-153.
- Kato, K. Ecological notes on the green frogs during the breeding season, I. Jap. J. Ecol. 5 2 1955: 70-73 5 figs.
- (2) Kato, K. Ecological notes on the green frogs during the breeding season, II. Breeding habit and others. Jap. J. Ecol. 6 2 1956: 57-61 4 figs.
- Kato, K. Studies on the differentiating potencies of the dorsal part of the blastopore lip in Triturus-gastruls.
   1. Differentiation of mesodermal tissues in relation to the neural tissue. Mem. Coll. Sci. Kyoto Univ. 25B 1958: 1-10.
- Kawasaki, K. Application of the indirect peroxidase reaction to tissue staining with special reference to cartilage tissue. The 28th report of histochemical study of peroxidase. Okajimas Folia Anat. Jap. 32 1958: 11–23.
- Kaye, N. W. & Le Bourhis, E. E. Uptake and turnover of a single injected dose of I<sup>131</sup> in tadpoles of Rana clamitans. Zoologica N.Y. 43 1958: 73-76 3 figs.
- Kelly, D. E. Embryonic and larval epiphysectomy in the salamander, *Taricha torosa*, and observations on scotiosis. J. Morph. **103** 1958: 503-537 4 figs.
- Kemp, N. E. Differentiation of the cortical cytoplasm and inclusions in cocytes of the frog. J. Biophys. & Biochem. Cytol. 2 No. 4 pt. 2 suppl. 1956: 187-190.
- Kennedy, J. P. Notes on a breeding congress of Pseudacris clarki and Pseudacris nigrita in Harris County, Texas. Herpetologica 14 1958: 192.
  - Kezer, J., see Hendricks, L. J.
- Khristoliubova, N. B. [An investigation of the myocardium with the electron microscope.] C.R. Acad. Sci. U.R.S.S. 119 1 1958: 168-170, pl. [In Russian.]
- (1) Kiortsis, V. Action combinée de l'ovariotomie, de la méthyl-testostérone et du nerf dévié sur le développement de la crête dorsale chez la femelle de Triturus cristatus Laur. Verh. schweiz. naturf. Ges. 138 1958: 144–145.
- (2) Kiortsis, V. L'induction d'extremités surnuméraires par la placode olfactive est-elle efficace sur tout le territoire du membre antérieur chez les embryons d'Urodèles? Arch. Sci. Genève 11 1958 : 159-164 3 figs.
- Kiortsis, V. & Droin, A. Importance de la moelle épinière dans la régénération caudale des larves de Salamandra salamandra Laur. Arch. Sci. Genève 11 1958: 235-245 3 figs.
- Kirtisinghe, P. Some hitherto undescribed anuran tadpoles. Ceylon J. Sci. Biol. Sci. 1 1958: 171–176 7 figs.
  - Kirtisinghe, P., see Grandison, A. G. C. (2).

Kitching, J. W. A new small Stereospondylous labyrinthodont from the Triassic beds of South Africa. Palaeont. africana 5 1958: 67-82 figs.

Kleerekoper, H. & Sibabin, K. A study on hearing in frogs (Rana pipiens and Rana clamitans). Z. vergl. Physiol. 41 1958: 490–499 9 figs.

Klekowski, R. The natterjack (Bufo calamita Laur.) in the neighbourhood of Warsaw. Przegl. Zool. 2 1958: 41-44 1 fig.

Klingelhöffer, W., see Gans, C.

Kloot, W. G. Van der. The effect of enzyme inhibitors on the resting potential and on the ion distribution of the sartorius muscle of the frog. J. gen. Physiol. 41 1958: 879-900 4 figs.

Koba, K. Herpetofauna of the Amami group of the Loo Choo Islands (II). Mem. Fac. Educ. Kumamoto Univ. 6: 173–185 photos.

Kobayashi, H. Effect of desoxycorticosteroneacetate on metamorphosis induced by thyroxine in anuran tadpoles. Endocrinology 62 1958: 371–377 3 firs.

Koblick, D. C. The characterization and localization of frog skin cholinesterase, J. gen. Physiol. 41 1958: 1129-1134.

Koketsu, K., Cerf., J. A. & Nishi, S. Electrical responses of frog spinal ganglion cells in sodium free media. Nature, Lond. 181 1958: 703-704 1 fig.

Kolatat, T., Kramer, K. & Mühl, N. Über die Aktivität sensibler Herznerven des Frosches und ihre Beziehungen zur Herzdynamik. Pflüg. Arch. Physiol. 264 1957: 127-144 9 figs.

Koloss, E. I. [One type of evolution of the striated muscle tissue.] J. Acad. Sci. U.R.S.S. biol. 4 1958 408-415 5 figs. [In Russian, English summary.]

Komoda, T., see Karasaki, S.

Kondo, E., see Tamura, J.

Konieczna-Marczyńska, B. & Skowron-Cendrzak, A. The effect of the augmented nerve supply on the regeneration in postmetamorphic Xenopus laevis. Folia biol., Kraków 1 1958: 37-46 13 figs.

König, D. Förekomst av Rana dalmatina Bonap. i Göteborgs omnejd. Fauna och Flora 53 1958: 94–96 1 pl.

Kosciuszko, H. The influence of the olfactory nerves on the regeneration of the endbrain in *Xenopus laevis* tadpoles. Folia biol., Kraków 6 1958: 117–130 8 figs.

(1) Koshida, Y. Autoradiographic study on iodine accumulation in pigmented tissues of amphibian larvae. Annot. Zool. jap. 31 1958: 141-146 l pl.

(2) Koshida, Y. On the fractionation of radioiodine in the skin of *Rhacophorus* larvae. Zool. Mag., Tokyo 69 1958: 261–263.

Koshtoyants Ch. S. [On a possible rôle of nucleic soids in the transmission of the nervous irritation, and on the effect of acetylcholine.] C.R. Acad. Sci. U.R.S.S. 120 4 1958: 926-928 figs. [In Russian.]

Kotani, M. On the formation of the primordial germ cells from the presumptive ectoderm of *Triturus* gastrulae. J. Inst. Polyt. Osaka City Univ. **8D** [1958] 145-160 1 pl. 3 figs.

Kozák, J., see Zelený, A.

Kramer, K., see Kolatat, T.

Krivosheev, V. G. [Occurrence of Bufo viridis in Northern Kyzyl Kum.] Priroda 3 1958: 114. [In Russian.]

(1) (Krolenko, S. A.). [Variation of the strength-duration curve of isolated muscle fibre treated with calcium chloride.] C.R. Acad. Sci. U.R.S.S. 120 3 1958 647-649. [In Russian.]

(2) (Krolenko, S. A.) [A study of the contractions of an isolated muscle fibre.] J. Acad. Sci. U.R.S.S. biol. 3 1958: 313–326 6 figs. In Russian with English summary.]

Kucias, J. Mehrfache Regeneration des Kaulquappenschwanzes des glatten Sporifrosches Xenopus laevis Daud. Naturwissenschaften 45 1958: 298–299 1 fig.

Kuczka, H. Verhaltensphysiologische Untersuchungen über die Wischhandlung der Erdkröte (*Bufo bufo* L.). Z. Tierpsychol. 13 1956: 185–207 figs. [English summary.]

Kumamoto, T. Studies of physical and chemical properties of neurosecretory substance II. Effect of ultra-centrifugation on neurosecretory system. Zoo. Mag., Tokyo 67 1958: 101-105.

Kumar, K. S. V. S., see Sampath Kumar, K. S. V.

(Kurentzov, A. I.) [Migration of grass frogs.] Priroda 2 1958: 119. [In Russian.]

Lác, J. Beitrag zur Nahrung der Bombina bombina L. Biológia, Bratislava 13 1958: 844-853 3 figs. [German summary.]

Lacquet, L. M. Potential diastolique et potentiel de repos du muscle ventriculaire cardiaque chez la Grenouille. C.R. Soc. Biol., Paris. 151 1958: 1993– 1995.

Ladman, A. J., see Burgos, M. H.

Lafontaine, J. G. & Ris, H. An electron microscope study of lampbrush chromosomes. J. biophys. & biochem. Cytol. 4: 1958: 99-106 4 pls.

Lakshman, A. B., see Ramaswami, L. S.

Lamotte, M. & Dzieduszycka, S. Contribution à l'étude des batraciens de l'Ouest africain. VII. Le développement larvaire de *Phrynobatrachus francisci*. Bull. Inst. franç. Afr. N. 29A 3 1958: 1071-1086 15 figs.

Lamotte, M. & Prum, P. Analyse quantitative du développement de la thyroïde au cours des métamorphoses de l'embryon de Nectophrynoïdes occidentalie Angel. C.R. Soc. Biol., Paris 151 1957 [1958]: 1187-1191 2 figs.

Lamotte, M. & Rey, P. Evolution de l'ovaire chez les femelles vierges de Nectophrynoides occidentalis Angel. C.R. Soc. Biol., Paris 151 6 1957 [1958] : 1191-1194 2 figs. U. 19. Sa. 19.

Bu

No con C.J

gro

et ]

151 du Mé: 3 p

11.

67 tra Ph

1

che

Ace I I Va.

see par Par

R.C [Er lipi Ric

ete I I la

glan tem 14

131

Lamotte, M., see Arnoult, J.

Lamotte, M., see Guibé, J.

(1) Lang, H. Eine hübsche Kröte aus Amerika: Bufo debilis. Aquar. Terrar. Z. 11 (2) 1958: 60.

2) Lang, H. Ein schöner Salamander aus dem U.S.A.: Ambystoma opacum. Aquar. Terrar. Z. 11 1958 : 118-119.

(3) Lang, H. Die nordamerikanischen "Wurm". Salamander Batrachoseps. Aquar. Terrar. Z. 11 1958 : 279-280, 1 fig.

Lapparent, A., F. de, Claracq, P. & Nougarède, F. Nouvelles découvertes de Vertébrés dans les séries continentales au Nord d'Edjelch (Sahara Central). C.R. Acad. Sci., Paris 247 1958: 2399-2402.

Lasansky, A. & Sabatani, D. D. Distribution des groupes sulfydrile et disulfure dans la neurohypophyse et l'hypothalmus du Crapaud. C.R. Soc. Biol., Paris 151 1958 : 1755.

(1) Laurent, R. F. La réserve naturelle intégrale du Mont Nimba. Les Rainettes du genre *Hyperolius*. Mém. Inst. franç. Afr. N. No. 53 1958 : 275–299

(2) Laurent, R. F. Notes herpétologiques africaines II. Rev. Zool. Bot. Afr. 58 1958: 115-128.

Layne, J. N., see Goin, C. J.

0.

.]

3-

86

ta-

ta-

3]:

lis

Le Bourhio, E. E., see Kaye, N. W.

Leach, A. A. The amino acid composition of amphibian, reptile and avian gelatins. Biochem. J. 67 1957: 83-87.

Leaf, A. Anderson, J. & Page, L. B. Active sodium transport by the isolated toad bladder. J. gen. Physiol. 41 1958: 657-669 5 figs.

Lecamp, M. Sur un cas singulier de régénération chez un Batracien anoure, Alytes obstetricans. C.R. Acad. Sci., Paris 246 1958: 1300-1301.

Lee, A. K., see Bentley, P. J.

Lee, A. K., see Main, A. R.

Leftwich, F. B. Blood oxygen capacity in frogs. Va. J. Sci. N.S. 9 1958: 397.

Legait, H. & Legait, E. Modifications cérébrales secondaires à l'ablation ou à la cautérisation de la paraphyse chez Rana esculenta. C.R. Soc. Biol., Paris 151 1958: 1418-1420.

Leone, V. Tumori da metilcolantrene in Tritoni. R.C. Ist. Lombardo 92B 1958: 220-240 5 pls. [English summary.]

Lernia, E. Di. Osservazioni sulla utilizzazione dei lipidi nello sviluppo embrionale di *Bufo vulgaris*. Ric. sci. **28** 1958: 2060–2064 2 figs. [English etc. summaries.]

Li, J.-C., see Giang, H.-C.

Liébecq-Hutter, S. & Bacq, Z. M. Disparition de la 5-hydroxytryptamine dans les granulations des glandes de la peau de Discoglossus pictus et de Rana temporaria après injection de réserpine. Experientia 14 1958 : 180-181.

Liedke, K. B. Experiments on the development of trunk muscles in Anura (Rana pipiens). Anat. Rec. 131 1958 : 97-118 19 figs.

Ling, K. An investigation on the breeding habits of Rana nigromaculata in the suburbs of Nanchang. Acta zool, sinica 10 1958: 421-430.

Liotti, F. S. Ricerche comparative sull'attivita catalasica (ossiforo di spirito) del sangue nelle varie classi dei vertebrati. Riv. Biol. 50 1958: 43-57. [English summary.]

Lison, L., see Hadler, W. A.

(1) Littlejohn, M. J. A new species of frog of the genus Crinia Tschudi from South Eastern Australia. Proc. linn. Soc. N.S.W. 83 1958: 222-226 1 fig.

(2) Littlejohn, M. J. Mating behaviour in the treefrog, Hyla versicolor. Copeia 1958 No. 3: 222-

Littlejohn, M. J., see Main, A. R.

**Livezey, R. L.** Procaine hydrochloride as a killing agent for reptiles and amphibians. Herpetologica 13 1958: 280.

Locker, A. Atmung, R. Q. und aerobe Glykolyse der Froschhaut unter dem Einfluss der Temperatur. Naturwissenschaften 45 1958: 398.

Locker, A. & Doneff, D. Der Einfluss der Tempera-tur auf die Abhängigkeit der Gewebsatmung der Froschleber vom Glykogengehalt. Z. vergl. Physiol. 41 1958 : 242-248 1 fig.

Loewenstein, W. R., see Hutter, O. F.

Lomont, H., see Petit, G.

Levtrup, S. A physiological interpretation of the mechanism involved in the determination of bilateral symmetry in amphibian embryos. J. Embryol. exp. Morph. 6 1958: 15-27.

Levtrup, 8. & Pigon, A. Inversion of the dorso-ventral axis in amphibian embryos by unilateral restriction of oxygen supply. J. Embryol. exp. Morph. 6 1958: 486-490.

Lucas, J. W. & Pickering, D. C. Direct absorption of dissolved strontium-90 and yttrium-90 by tadpoles of Rana temporaria. Nature, Lond. 182 1958: 1242-1243 1 fig.

Lunenfeld, B., Rabau, E., Harell-Steinberg, A. & Szejnberg, A. Effect of injection medium on quantal assay of chorionic gonadotrophin in male toads. Acta Endocrin. 24 1957: 113-118.

Luther, W., see Oberheim, K. W.

Lüttgas, H. C. & Niedergerke, R. The antagonism between Ca and Na ions on the frog's heart. J. Physiol. 143 1958: 486-505 11 figs.

Lutz, B. & Carvalho, A. L. de. Novos anfibios anuros das serras costeiras do Brasil. Mem. Inst. O. Cruz 56 1958 : 239-259 5 pls.

Lynn, W. G. Some amphibians from Haiti and a new subspecies of *Eleutherodactylus schmidti*. Herpetologica 14 1958: 153-157.

Lynn, W. G., see Dent, J. N.

Macgregor, H. C., see Callan, H. E.

Maetz, J., Jard, S. & Morel, F. Action de l'aldostérone sur le transport actif de sodium de la peau de grenouille. C.R. Acad. Sci. Paris 247 1958 : 516-518 1 fig.

rep Ce (B

5 1

tac

na

4 1

Ar

th

ric

Se

co

ch

63

19

WE

to

19

bic 27

m

E

Se

84

H

Main, A. R., Lee, A. K. & Littlejohn, M. J. Evolution in three genera of Australian frogs. Evolution 12 1958: 224-233 5 figs.

Main, A. R., see Bentley, P. J.

- (1) Malamed, S. Gastrular blockage of frog's eggs produced by oxgyen at atmospheric pressure. Exper. Cell Res. 13 1957: 391–394.
- (2) Malamed, 8. Gastrular blockage in frog's eggs produced by oxygen poisoning. Biol. Bull. Woods Hole 114 1958: 226-246 8 figs.
- Malhotra, S. K. The cytoplasmic inclusions of the ageing neurones of the frog Rana tigrina. Cellule 58 (3) 1957: 363-387, 2 pls. 38 figs.
- Malinin, T. & Deck, J. D. The effects of implantation of embryonic and tadpole tissues into adult frog limbs. J. exp. Zool. 139 1958: 307-324 1 fig. 1 pl.
- Manacas, S. Amphibians and reptiles of the islands of São Tomé and Principe and of Rolas Islet. Conf. int. Afr. occid. Sess. 6 4: 179–192 2 figs. [Not seen.]
- (1) Manelli, H. Ricerche sperimentali su l'origine e lo sviluppo del pronefro e del dotto di Wolff nella Rana esculenta. Arch. Sci. Biol. Bologna 41 1957 : 404-408 2 figs.
- (2) Manelli, H. Le variazioni del quadro ematico elettroforetico di *Amblystoma* in rapporto all'età, al sesso e alla metamorfosi. Riv. Biol. **50** 1958: 15–26 4 figs. [English summary.]
- Manner, H. W. The metabolic effect of cortisone acetate during salamander limb regeneration. Growth 22 1958: 93-98 1 pl.
- Marco, F. Di. Influenza della Streptomicina e del Cloramelnicolo sulla metamorfosi dei girini di Rana esculenta L. Boll. Accad. gioenia (4) 4 1958 : 281–287. [English etc. summaries.]

Marcot, M., see Jullien, A.

Martin, P. S. A biogeography of reptiles and amphibians in the Gomez Farias Region, Tamaulipas, Mexico. Misc. Publ. Mus. Zool. Univ. Mich. No. 101 1958: 1-102 7 pls.

Martin, S. J., see Fulchiero, E.

Martof, B. S. & Thompson, E. F. Reproductive behaviour of the chorus frog, *Pseudacris nigrita*. Behaviour 13 1958: 243-258 l pl.

Marx, H. Catalogue of type specimens of reptiles and amphibians in Chicago Natural History Museum. Fieldiana, Zool. 36 (4) 1958: 409–496.

Matsuyama, S., see Hashimoto, K.

Matsuzaki, H., see Asayama, S.

Maturana, H. R. Efferent fibres in the optic nerve of the toad (*Bufo bufo*). J. Anat., Lond. 92 1958: 21-27 2 pls.

Mayer, D. T., see England, M. C.

Mazzi, V. Effetti di lesioni alla eminenza mediale sul sistema ipotalamo-ipofisario, adenoipofisi, tiroide e testiciol del tritone crestato. Z. Zellf. 48 1958: 332-348 15 figs.

Medda, A. K., see Bhattacharya, G. C.

Meltzer, R. I., see Money, W. L.

Meng, M. Untersuchungen zum Farben und Formensehen der Erdkrote (*Bufo bufo* L.). Zool. Beitr., Berl. **3** 1958: 313–363, 21 figs.

Menon, K. R., see Barnabas, T.

Merrick, A. W., see Fitch, K. L.

- (1) Mertens, R. Welche Bedeutung haben die "Haare" des Haarfrosches (*Trichobatrachus robustus*). Zool. Anz. **161** 1958 : 82–85,
- (2) Mertens, R. Wie findet der Grasfrosch seine Laichgewässer? Kosmos, Stuttgart 54 1958: 290.
- (3) Mertens, R. Ergänzende Bemerkungen über Batrachomyia-Larven und ihre Wirte. Senek. biol. 39 1958: 197–198.
- (4) Mertens, R. Als Zoologe in Australien, III. Im Herzen des fünften Kontinents. Natur u. Volk 88 1958: 264-276 10 figs.
- Mews, H.-H. Über die Temperaturadaptation der eiweisssppaltenden und synthetisierenden Zellfermente von Froschen. Z. vergl. Physiol. 40 1957: 356–362.
- Meyer, M. Tryptaflavinwirkungen auf Amphibienmitosen. Z. Zellf. 47 1958: 731-759 14 figs.

Michael, M. I., see Sedra, S. N.

Michalowski, J. Geographische Verbreitung der Unken (Bombina Oken) zwischen den Flüssen Weichsel, Skawa und Raba (Woiwodschaft Kraków). Acta zool. Cracov. 3 1958: 247—282 2 pls.

Michel, R., see Volpent, E.

Mighorst, J. C. A., see Burgers, A. C. J.

Miller, R. L., see Olson, E. C.

Mitchison, J. M., see Swann, M. M.

Miyamori, H., see Asayama, S.

- Moisset de Espanés, B. Essais quantitatifs sur la spermiation par gonadotrophine chorionique chez le Crapaud mâle. C.R. Soc. Biol., Paris 151 1958: 1963–1964.
- Money, W. L., Meltzer, R. I., Young, J. & Rawson, R. W. The effect and change in chemical structure of some thyroxine analogues on the metamorphosis of Rana pipiens tadpoles. Endocrinology 63 1958: 20-28 5 figs.
- (1) Moore, J. A. Transplantation of nuclei between Rana pipiens and Rana sylvatica. Exp. Cell. Res. 14 1958: 532-540.
- (2) Moore, J. A. A new genus and species of Leptodactylid frog from Australia. Amer. Mus. Novit. No. 1919 1958: 1-7 2 figs.
- (3) Moore, J. A. The transfer of haploid nuclei between Rana pipiens and Rana sylvatica. Exp. Cell. Res. suppl. 6 1958: 179-191 5 figs.

Morawska, B. & Zaleska, A. The influence of the temperature on the pulse rate of tadpoles and metamorphosed individuals of Xenopus laevis D., Rana temporaria L. and Rana esculenta L. Acta Biol. exper. Warszawa 18 1958: 39-53 7 figs.

Morel, F., see Maetz, J.

- (1) Morgan-Davies, A. M. Some notes on the reproduction, metamorphosis and the ecology of the Ceylonese tree frog, *Rhacophorus cruciger cruciger* (Blyth). J. Bombay nat. Hist. Soc. **55** 1 1958: 11-16 5 figs.
- (2) Morgan-Davies, A. M. Notes on the eggs, tadpoles, metamorphosis, and ecology of the Ceylonese narrow-mouthed frog Ramanella obscura (Günther). J. Bombay nat. Hist. Soc. 55 2 1958: 307-312 2 pls. 4 figs.
- (3) Morgan-Davies, A. M. An introduction to the Amphibia of Ceylon. Loris 8 (1) 1958: 1-5 6 figs.

Morita, Y., see Hashimoto, K.

Moriwaki, K. Influence of temperature in altering the augmenting effect of sodium azide on the energy-rich phosphate content of the toad embryo. J. Fac. Sci. Univ. Tokyo (Zool.) 8 1958: 287–295.

Moses, M. J. The relation between the axial complex of meiotic prophase chromosomes and chromosome pairing in a salamander (*Plethodon cinereus*). J. biophys. & biochem. Cytol. 4 1958: 633-638 3 figs.

Mosimann, J. E. The evolutionary significance of rare matings in animal populations. Evolution 12 1958: 246-261 6 figs.

Moulin, J., see Jullien, A.

n

0

f

n

Moursi, A. A., see Sedra, S. N.

Mühl, N., see Kolatat, T.

Mullally, D. P. Daily period of activity of the western toad. Herpetologica 14 1958: 29-31.

Mullally, D. P. & Powell, D. H. The Yosemite toad: northern range extension and possible hybridization with the western toad. Herpetologica 14 1958: 31-33.

Muller, M., see Simon, S. E.

Müller-Gehring, R. Eine Methode, lebende Amphibien und Reptilien abzugiessen. Präparator 1 1955: 27–28 1 fig.

Mullins, L. J., see Sjodin, R. A.

Munday, K. A. The relation of endogenous melanophore-expanding hormone to hyperglycaemia in Xenopus laevis. J. Endocrin. 15 1957: 190-198.

Murray, R. W. The response of the lateralis organs of Xenopus laevis to electrical stimulation by direct current. J. Physiol. 134 1956: 408-420.

- (1) Myers, C. W. Notes on the eggs and larvae of Eurycea lucifuga Rafinesque. Quart. J. Fla. Acad. Sci. 21 1958: 125-130 1 fig.
- (2) Myers, C. W. Subspecific identity of Rana sylvatica in Missouri. Copeia 1958 (1): 46.
- (3) Myers, C. W. Amphibia in Missouri caves, Herpetologica 14 1958: 35–36.

Nacimiento, A. C. Titration de l'adrénaline et de la noradrénaline par la préparation de Laewen Trendelenburg du Crapaud. C.R. Soc. Bioî. Paris 152 1958: 195-196 2 figs.

Nagasawa, M., see Jimbo, G.

- Nagatani, Y. Studies on the cytoplasmic granules stainable with carcinogenic azo-dye, O-amino-azo-toluene. I. Classification of cytoplasmic granules in amphibian crythrocytes and their cytochemical natures. Jap. J. Zool. 12 1958: 163-177 2 pls.
- Nagler, A. L. & Zweifach, B. W. The effect of bacterial endotoxins and biogenic amines on the phagocytic behaviour of endothelial elements in the frog Rana pipiens. Biol. Bull. Woods Hole 115 1958: 357. [Abst.]
- Natarajan, R. Contributions to the cytology of Indian Anura (Amphibia): A study of chromosomes of the two sexes of *Bufo melanostictus* Schneider (Bufonidae). Proc. Indian Acad. Sci. **48B** 1958: 297-300 1 pl.
- Nath, V., Gupta, B. L. & Kaur, R. Histochemical and morphological studies of the lipids in oogenesis, VI. The frog, *Rana tigrina*. Res. Bull. Panjab. Univ No. 153 1958: 223–234a 5 figs.
- Nayler, W. G. Further observations on the seasonal variation in drug sensitivity of the isolated toad heart (*Bufo marinus*). Aust. J. exp. Biol. med. Sci. 36 1958: 133-144 5 figs.

Nechaeva, N. V., see Brodsky, V. Y.

Neidleman, S., see Bagnara, J. T.

- (1) Neill, W. T. The rapid mineralization of organic remains in Florida and its bearing on supposed Ploistocene records. Quart. J. Fla. Acad. Sci. 20 1 1957: 1-13.
- (2) Neill, W. T. The occurrence of amphibians and reptiles in saltwater areas, and a bibliography. Bull. mar. Sci. Gulf & Caribbean 8 1 1958: 1-97.
- (3) Neill, W. T. The varied calls of the barking treefrog Hyla gratiosa Leconte. Copeia 1958 (1): 44-46.
- Neill, W. T., Gut, H. J. & Brodkorb, P. Animal remains from four preceramic sites in Florida. Amer. Antiq. 21 4 1956: 383–395.
- Noisen, O. E. Increased partial pressure of O<sub>3</sub> and the acceleration of development in early chick and frog embryos; significance of early primitive streak activities. Growth 22 1958: 189-213.
- (1) Newth, D. R. On regeneration after amputation of abnormal structures. I. Defective amphibian tails. J. Embryol. exp. Morph. 6 1958: 297-307, 3 figs.
- (2) Newth, D. R. On regeneration after the amputation of abnormal structures. II. Supernumerary induced limbs. J. Embryol. exp. Morph. 6 1958: 384-392 2 pls.
- Nicholas, J. S. Factors influencing symmetry, posture and reduplication of the developing limb of Amblystoma punctatum. Anat. Rec. 131 1958: 475–485. 11 figs.

Nieder, K., see Skjelkvale, L.

Niedergerke, R., see Lüttgau, H. C.

Nielsen, L., see Jørgensen, C. B.

Nieuwkoop, P. D., see Seno, T.

Nigrelli, R. F., see Jakowska, S.

Nikolsky, N. N. Changes of functional characteristics of frog nerve at the action of various agents. Vest, Leningr. Univ. No. 15 1958: 102-110.

Nishi, S., see Koketsu, K.

Niweliński, J. The effect of prolactin and somatotrophin on the regeneration of the forelimb in the newt *Triturus alpestris* Laur. Folia biol., Kraków 6 1958: 9-36 25 figs.

Noseda, V., see Aguggini, G.

Notarnicola, L. Esperienze di asportazione e successivo trapianto ortotopico di arto negli anfibi anuri. Conseguenze sul sistema nervoso. Rev. Biol. 50 1958: 137–167 6 figs. [English summary.]

Nougarède, F., see Lapparent, A. F.

(Novikov, G. A.) IV Amphibia. Zhivot. Mir. 5 1958: 558. [In Russian.]

Numata, J., see Tamura, J.

Oberheim, K. W. & Luther, W. Versuche über die Extremititenregeneration von Salamanderlarven bei umgekehrter Polaritat des Amputationsstumpfes. Roux Arch. EntwMech. Organ 150 1958: 373-382 2 figs.

Obika, M., see Hama, T.

- Ocser, R. Flectonotus goeldii ein brutpflegender Bromelienfrosch. Aquar. Terrar. Z. 11 1958: 146-149 l fig.
- (1) Oliver, J. A. Vida de batracios y reptiles sudamericanos by M. A. Freiberg. [Review.] Copeia 1958 No. 1: 63.
- (2) Oliver, J. A. Sounds of north American frogs by C. M. Bogert. [Review.] Copeia 1958 No. 3: 241-2
- Olson, E. C. Fauna of the Vale and Choza: 14 Summary, Review and Interpretation of the Geology and the Faunas. Fieldiana Geol. 10 1958: 397-448 figs.
- Olson, E. C. & Miller, R. L. Morphological Integration. 1958: pp. xv, 317, 77 figs. Chicago.
- Oordt, G. J. van, Beenakkers, A. M. Th., Oordt, P. G. W. J. van & Stadhouders, A. M. On the mechanism of the initial stages of spermiation in the grass frog, Rana temporaria. Acta Endocr. 17 1954: 294-301.
- Oordt, P. G. W. J. van. Regulation of the spermatogenetic cycle in the common frog (Rana temporaria), G. W. Van der Wiel & Co., Arnhem 1956: xiii+11653 figs.

Oordt, P. G. W. J. van., see Oordt, G. J. van.

Organ, J. A. Courtship and spermatophore of Plethodon jordani metcalfi. Copeia 1958 No. 4: 251-259 3 figs.

Ormsby, A. I. Partial neoteny in an Australian frog. Brit. J. Herpet. 2 1958: 107-108.

Oros, I., see Pora, E.-A.

Osanai, K. Cortical change at fertilization in the toad eggs. Bull. mar. biol. Sta. Asamushi 9 1958: 47-48.

O'Steen, W. K. Regeneration of the intestine in adult urodeles. J. Morph. 103 1958: 435-477 4 pls.

Page, L. B., see Leaf, A.

Pantelouris, E. M. Protein synthesis in newt occytes. Exp. Cell. Res. 14 1958: 584-595 5 figs.

Pantelouris, E. M. & Jacob, J. Nuclear chimaeras in the newt. Experientia 14 1958: 99.

Paola, D. de., See Schulz, H.

- Parker, H. W. Caecilians of the Seychelles Islands with description of a new subspecies. Copeia 1958 No. 2: 71-76 3 figs.
- (1) Pasteur, G. Sur la systematique des espèces du genre Pleurodeles (Salamandridés). Bull. Scc. Sci. nat. phys. Maroc 38 1958 : 157–165.
- (2) Pasteur, G. Sur les tendances évolutives et la phylogénie des *Pelobates* (Batraciens Anoures) actuels. C. R. Acad. Sci., Paris 247 1958: 1037–1039.

Payne, K. E., see Ferguson, D. E.

- (1) Pearson, P. G. Further notes on the population ecology of the spadefoot toad. Ecology  $38\ 1957:580-586\ 2$  figs.
- (2) Pearson, P. G. Body measurements of Scaphiopus holbrooki. Copeia 1958 No. 3: 215-217.

Pécot-Dechavassine, M. Différenciation histochimique des cholinestérases au niveau des jonctions neuro-musculaires et musculotenineuses du muscle droit abdominal de la Grenouille. C.R. Acad. Sci., Paris 247 1958. 2489-2491.

Pécot-Dechavassine, M., see Jacob, J.

- Penhos, J. C. & Cardeza, A. F. Les caractères sexuels du Crapaud castré traité par les hormones sexuelles et l'acide folique ou l'aminoptérine. C.R. Soc. Biol., Paris 151 1958: 1601–1602.
- (1) Perret, J.-L. Observations sur des rainettes africaines du genre Leptopelis Günther. Rev. suisse Zool. 65 1958: 259-275 11 figs.
- (2) Perret, J.-L. Notes sur des Batraciens du genre Kassina du Cameroun et d'Afrique nord-équatoriale. Bull. Inst. franç. Afr. noire 20A 1958: 1437-1447 7 figs.
- (3) Perret, J.-L. Curieux batraciens du Cameroun. Bull. Mensuel Mus. et Collections Ville Genève 15 5: [no pagination] 2 photos.
- (1) Perri, T. Fattori intrinseci ed estrinseci nello sviluppo del midollo spinale e dei gangli spinali negli anfibi anuri. Ulteriori risultati e considerazioni. Riv. Biol. 50 1958: 1-14 1 pl., 1 fig. [English summary.]
- (2) **Perri, T.** Embryonic implantations of the limb buds of *Bufo viridis* upon *Bufo vulgaris*. Riv. Biol. **50** 1958: 319–326. [English sumary.]

Perseca, T., see Preda, V.

- Peters, J. J. & others. Rhythmic electrical activity of skeletal muscle in the salamanders and its modification by convulsant and anti-convulsant drugs. Physiol. Zool. 31 1958: 288–294.
- Petit, G. & Lomont, H. Nouvelle pullulation de Pelobates dans la région du Canet (Pyr.-Or.). Vie et Milieu 9 1958: 131-133.

nel Buj [En

E

131

fau and 217

the I mic

gra

gra

log

A

cu

eu 19

in an hi

1

d C

1

in

vt.

11

i.

a

B)

n

)-

Petrucci, D. Citocromo c-ossidasi e respirazione nel corso dello sviluppo larvale e della metamorfosi di Bufo bufo. Ric. sci. 28 1958: 541-548 4 figs. [English summary.]

Philipp, G. A. Myobatrachus gouldii in the coastal hills near City Beach. W. Aust. Nat. 6 1958: 131-132.

Philippi, R. A., see Cei, J. M. (4).

Phillipe-Knoepffler, L., see Duguy, R.

Phillips, W. W. A. Some observations on the fauna of the Maldive Island. Part IV. Amphibians and reptiles. J. Bombay nat. Hist. Soc. 55 1958: 217-220.

Piatt, J. & M. Transection of the spinal cord in the adult frog. Anat. Rec. 131 1958: 81-95 13 figs.

Piatt, M., see Piatt, J.

Pick, J. Sympathectomy in amphibians. (Anatomical considerations.] J. comp. Neur. 107 1957: 169-208.

Pickering, D. C., see Lucas, J. W.

Pigon, A., see Levtrup, S.

Pilz, A., Sickel, W. & Birke, R. Elektroretinographische Prüfung ortlicher Empfindlichkeitsunterchiede an der Netzhaut von Rana exculenta. Pflüg. Arch. Physiol. 266 1958: 193–198 3 figs.

(1) Pimentel, R. A. On the validity of Taricha granulosa Skilton. Herpetologica 14 1958: 165-168.

(2) Pimentel, R. A. Southern occurrence of the sharp-tailed snake in a unique community [amphibians from San Luis Obispo Co. Pine Mt. listed.] Herpetologica 14 1958: 169-170.

Pinacci, R., see Baffoni, G.-M.

Pisanó, A. Variaciones somáticas en larvas de Telmatobius hauthali schreiteri de la Cordillera Andina, Acta zool. lillo. 16 1958: 5-12 3 figs.

Plutner, J. M. Western red-bellied newt in Lake County, California. Copeia 1958 No. 4: 339.

Poczopko, P. Further investigations on the cutaneous vasomotor reflexes in the edible frog in connexion with the problem of regulation of the cutaneous respiration in frogs. Zool.polon. 8 (2-3) 1957 [1958]: 161-175 3 figs.

Poglayen, I. & Smith, H. M. Noteworthy herptiles from Mexico. Herpetologica 14 1958: 11-15, 1 pl.

Polyak, S. The vertebrate visual system. Its origin, structure, and function and its manifestations in disease with an analysis of its role in the life of animals and in the origin of man. Preceded by a historical review of investigations of the eye, and of the visual pathways and centers of the brain. Chicago, (University Press), 1957, pp. xviii+1390, 18 pls. frontis (col.), 546 figs.

Pora, E.-A. & Oros, I. Contributions à l'étude de l'action du système nerveux sur le travail mecanique du gastrocnémien de la grenouille. Stud. Cercet. Biol. Cluj 9 1958: 345–350 l fig. [French summary.]

Pora, E. A. & Rejep, A. La modification de la motricité du tractus gastro-intestinal de la grenouille au cours d'une année. Stud. Cercet. Biol., Cluj 8 1958: 349-357 9 figs. [French, etc. summaries.]

Pora, E. A. & Toma, V. L'action des extraits de thymus sur le travail méchanique du gastroenémien de la grenouille. Stud. Cercet. Biol., Cluj 8 1957: 343-347 2 figs. [French, etc. summaries.]

Powell, D. H., see Mulially, D. P.

Powers, T. H., see Peters, J. J.

Preda, V. & Persecă, T. La régénération de la queue du triton et le rôle du cerveau antérieur dans la première étape du processus régénératif. Bull. Univ. Vi Babeş "şi "Bolyai "Cluj. Ser. Sti. nat. 2 1957: 509-514 4 pls. [French, etc. summaries.]

Prum, P., see Lamotte, M.

Pyburn, W. F. Size and movements of a local population of cricket frogs (Acris crepitans). Texas J. Sci. 10 1958: 325-342 5 figs.

Pylka, J. M. & Warren, R. D. A population of *Haideotriton* in Florida. Copeia 1958 No. 4: 334-336 2 figs.

Rabau, E., see Lunenfeld, B.

Rabb, G. R. On certain Mexican salamanders of the plethodontid genus *Chiropterotriton*. Occ. Pap. Mus. Zool. Univ. Mich., No. 587 1958: pp. 37, 3 pls.

Rabor, D. S., Alcala, A. C. & Gonzales, R. B. A brief list of the land Vertebrates of Negros Island. Silliman J. 5 1958: 286-300.

Radovanovic, M. Einige eigentümliche Lurche und Kriechtiere der jugoslawischen Karstgebietes. Verh. dtsch. zool. Ges. 1957 [1958]: 512-516.

(Raitrina, S. S.) [Regeneration in newts and in some mammals.] Bull. Soc. Nat. Moscou biol. 63 (2) 1958: 169–170. [In Russian.]

Ramaswami, L. S. The development of the apodan vertebral column, Zool. Anz. 161 1958: 271-280 9 figs.

Ramaswami, L. S. & Lakshman, A. B. Ovulation induced in frog with mammalian hormones. Nature, Lond. 181 1958: 1210.

Rawson, R. W., see Money, W. L.

Ray, C. Vital limits and rates of desiccation in Salamanders. Ecology 39 1958: 75-83 4 figs.

[Razumovich, M. B.] [Frog haemoglobin content.] Priroda 7 1958: 99-101. [In Russian.]

(1) Reed, C. F. Contributions to the herpetology of Maryland and Delmarva. No. 17: Southeastern herptiles with northern limits on coastal Maryland, Delmarva, and New Jersey. J. Wash. Acad. Sci. 48 1958: 28-32 16 maps.

(2) Reed, C. F. The carpenter frog in Worcester Co., Maryland. Herpetologica 13 1958: 276.

(3) Reed, C. F. Contributions to the herpetology of Maryland and Delmarva, 13: Piedmont herpetofauna in coastal Delmarva. J. Wash. Acad. Sci. 48 1958: 95–99 17 maps.

Reig, O. A. Proposiciones para una nueva macrosistemática de los anuros (Nota preliminar). Physis, B. Aires 21 60 1958: 109–118.

Reig, O. A., see Stipanincie, P. N.

Rejep, A., see Pora, E. A.

Rey, P., see Lamotte, M.

Reyer, R. W. Studies on lens induction in Amblystoma punctatum and Triturus viridescens viridescens. I. Transplants of prospective belly ectoderm. J. exp. Zool. 138 1958: 505-548 4 figs. 3 pls.

Reynolds, A. E. A parallelism of cheek skin colour in Plethodontid salamanders, Proc. Indiana Acad. Sci. 67 1958: 303.

Rhodes, J. R., see Ferguson, D. E.

Richards, C. M. The inhibition of growth in crowded Rana pipiens tadpoles. Physiol. Zool. 31 1958: 138-151 13 figs.

(1) Riemer, W. J. Giant toads of Florida. Quart. J. Fla Acad. Sci. 21 1958: 207-211.

(2) Riemer, W. J. Variation and systematic relationships within the salamander genus *Taricha*. Univ. Calif. Publ. Zool. **56** (3) 301–390 24 figs.

Riley, H. L., see Deck, J. D.

Ripplinger, J., see Jullien, A.

Ris, H., see Lafontaine, J. G.

Robertson, J. D. The ultrastructure of Schmidt-Lauterman clefts and related shearing defects of the myelin sheath. J. biophy. & biochem. Cytol. 4 1958: 39-46 4 pls 3 figs.

Roche, J., see Volpert, E.

Roguski, H., see Skowron, S.

Romanovsky, A. Regeneration of the central nervous system in the early developmental stages of the axolotl (Ambystoma mexicanum Cope). III. Regeneration after extirpation of one half of the brain rudiment in the advanced tail-bud and later stages of embryogenesis. Mém. Soc. zool. tchécosl. 22 1958: 371-380 1 fig. 4 pls.

Romer, A. S. An embolomere jaw from the midcarboniferous of Nova Scotia. Breviora No. 87 1958:

Rondell, P. A. & Wright, P. A. Metabolic studies of ovulation in the frog. Physiol. Zool. 31 1958: 236-243.

Rose, S. M. A feed back mechanism of growth control in tadpoles. Biol. Bull. Woods Hole 115 1958: 320. [Abst.]

Rosenbaum, R. M. Histochemical observations on the cortical region of the occytes of Rana pipiens. Quart. J. micr. Sci. 99 1958: 159-169 4 figs.

Rosenbaum, R. M., see Wittner, M.

Rossi, A. Sull'incompatabilità larvale nei trapianti xenoplastici di Urodeli su Anuri. Processi degenerativi delle code di *Triton taeniatus* trapiantate nel campo dell'arto posteriore di *Discoglossus pictus* e di *Hyla arborea*. R.C. Accad. Lincei. (8) 24 1958: 193–200.

(1) Rossman, D. A. The treefrog Hyla gratiosa in Tennessee. Herpetologica 14 1958: 40.

(2) Rossman, D. A. A rew race of Desmognathus fuscus from the South-Central United States. Herpetologica 14 1958: 158-160.

(1) Rostand, J. Sur un nouveau type de polydactylie chez la Grenouille verte (Rana esculenta L.). C.R. Acad. Sci. Paris. 247 1958: 607-609.

(2) Rostand, J. Effet tératogène des rayons ultraviolets sur les larves d'Amphibiens. C. R. Acad. Sci. Paris 247 1958: 972–974.

(1) Roth, P. C.-J. Senil d'action de la tetraiododesaminothyronine sur la metamorphose des tetards de Rana temporaria L. Ann. Endocr. Paris 17 1956: 725-728.

(2) Roth, P. C.-J. Action de la thyroxamine sur la métamorphose des tetards de Rana temporaria L. C.R. Soc. Biol., Paris 151 1958: 1130-1132.

Rotter, J. Die Reise nach Transkaukasien. Aquar. u. Terr. 5 1958: 286–291 9 figs.

Rounds, D. E. & Flickinger, R. A. Distribution of ribonucleoprotein during neural induction in the frog embryo. J. exp. Zool. 137 1958: 479-499.

Ruben, L. N. & Frothingham, M. L. The importance of innervation and superficial wounding in urodele accessory limb formation. J. Morph. 102 1958: 91-117 11 figs.

Rudrick, D. Embryonic nutrition. (Chicago University Press). 1958: pp. xi+113, text figs.

Ruhland, G. Die Ursachen der Lebensunfähigkeit haploider Amphibien-und Knochenfischlarven (haploidkerniges Syndrom). Z. mikr. anat. Forsch. 64 1958: 30–43 9 figs.

Ruhland, G., see Hertwig, G.

(1) Rühmekorf, E. Beiträge zur Ökologie mitteleuropäischer Salientia. I. Abhängigkeit der Laichabgabe von Aussenfaktoren. Z. Morph. Ökol. Tiere 47 1958: 1-19 6 figs.

(2) Rühmekorf, E. Beiträge zur Ökologie mitteleuropäischer Salientia. II. Temperaturwahl der Larven. Z. Morph. Ökol. Tiere 47 1958: 20–36 3 figs.

(3) Rühmekorf, E. Beiträge zur Ökologie mitteleuropäischer Salientia. III. Wirkungen des Lichtes auf Laich und Larven. Z. Morph. Ökol. Tiere 47 1958: 37-53 9 figs.

Ruibal, R., see Hecht, M. K.

Saadi-Nazim. Action du fluorure de sodium sur la métamorphose de la Grenouille. C.R. Soc. Biol., Paris 151 1958: 1137-1138.

Sabatani, D. D., see Lasansky, A.

Sacarrão, G. F. Cromosomas de cauda do girino do Tritão e da gónada de larva de Salamandra. Naturalia 7 1958 : 226-229 5 figs.

St. Clair, W. A., see Trethewie, E. R.

Sampath Kumar, K. S. V., Sundararajan, T. A. & Sarms, P. S. Biosynthesis of phosphoprotein in the frog. Proc. Soc. exp. Biol. Med. 99 1958: 539-541.

Samtleben, H. Vergleichende Untersuchungen an Hirnen verschiedener Anuren. Z. wiss. Zool. 161 1958: 38–83 20 figs.

Sarma, P. S., see Sampath Kumar, K. S. V.

to Act

[

d'as

8

Cor

eolo

Cy Lir 72

ve

Ru

ver pre 15

57

Bi 39

d Z

ta

.

i

i.

le

n

0

Э

- Sarzana, G. & Castro, V. Diversa elettiortà d'azione dell' adrenalina sui depositi di glicogeno del colombo e della rano. Arch. Fisiol. Firenze 55 1955 52-57.
- Savage, J. M. Zoogeography: the geographical distribution of animals by P. J. Darlington. [Review.] Copeia 1958 No. 2: 153-155.
- Saxén, L. The onset of thyroid activity in relation to the cytodifferentiation of the anterior pituitary. Acta anat. 32 1958: 87-100 3 figs.
- Schenk, R. & Töndury, G. Der Einfluss von Cytostatica auf Wachstum und Differenzierung der Linsen larvaler Tritonaugen. Acta Anat. 30 1957: 725-738.

#### Schjeide, O. A., see Flickinger, R. A.

- (1) (Schmalhausen, I. I.) [Morphology of the vertebral column in Urodela. II. The origin of the centra.] Zool. Zh. 37 2 1958: 229-239 4 figs. [In Russian with English summary.]
- (2) (Schmalhausen, I. I.) [Morphology of the vertebral column of Urodela. III. The transverse processes and ribs.] Zool. Zh. 37 3 1958: 405–429 15 figs. [In Russian with English summary.]
- (3) (Schmalhausen, I. I.) [Nasolachrymal duct and septomaxilla of Urodela.] Zool. Zh. 37 4 1958: 570–583 13 figs. [In Russian with English summary.]
- (4) Schmalhausen, I. I. The origin of the Amphibia. Bull. Acad. Sci. U.R.S.S. Sér. Biol. 1958: 1 1958: 39-58 12 figs. map. [In Russian.]
- Schmidt, A. J. Forelimb regeneration of thyroidectomized adult newts. J. exp. Zool. 137 1958: 197–226 13 figs.

Schmidt, K. P., see Cei, J. M. (3).

Schmidt, S. E., see Butler, E. G.

- (1) Schmidt, W. J. Zur durodentinbildung bei Urodelen-Zähnen. Z. Zellforsch. 46 1957: 281–285.
- (2) Schmidt, W. J. Zur Histologie und Färbung der Zähne des japanischen Riesensalamanders. Z. Zellforsch. 49 1958: 46-57 9 figs.

Schott, J., see Vivien, J. H.

- Schotté, O. E. & Wilber, J. F. Effects of adrenal transplants upon forelimb regeneration in normal and in hypophysectomized adult frogs. J. Embryol. exp. Morph. 6 1958: 247-261 4 pls.
- **Schulerer, F. W.** Factors affecting neoteny in the salamander *Dicamptodon ensatus* (Eschscholtz). Bull, S. Calif. Acad. Sci. **57** 1958: 119-121.
- Schulz, H. & Paola, D. de. Delta-Cytomembranen und lamelläre Cytosomen, Ultrastruktur, Histochemie und ihre Beziehungen zur Schleinsekretion. Z. Zellforsch. 49 1958: 125–141 12 figs.
- (J) Schumacher, G. H. Ein Beitrag zur Kaumuskulatur der Amphibien Untersuchungen an Cryptobranchus japonicus. Anat. Anz. 105 1958: 361–378.
- (2) Schumacher, G. H. Zur Morphologie des Mundbodens der Urodelen. Untersuchungen an Crytobranchus japonicus. Gegenbaurs Jb. 99 1958: 344-371 10 figs.

- (1) Schwartz, A. A new frog of the auriculatus group of the genus Eleutherodactylus from Western Cuba. Herpetologica 14 1958: 69-77 2 figs.
- (2) Schwartz, A. Another new large Eleuthero-dactylus (Amphibia: Leptodactylidae) from Western Cuba. Proc. biol. Soc. Wash. 71 1958: 37-42 1 fig.
- (3) Schwartz, A. Another new species of *Eleutherodactylus* (Amphibia: Leptodactylidae) from Western Cuba. J. Wash. Acad. Sci. 48 1958: 127–131 1 fig.
- (4) Schwartz, A. Four new frogs of the genus Eleutherodactylus (Leptodactylidae) from Cuba. Amer. Mus. Novit. No. 1873 1958: 1-20 4 figs.

Schwartz, A., see Duellman, W. E.

- Sculthorpe, D. Behaviour and reproduction of various news species. Aquarist & Pondkpr. 22 1958: 258-260 3 figs.
- Sedra, S. N. Seven interesting abnormal urostyles of *Bufo regularis* Reuss. Čs. Morfologie 6 1958: 1-6 7 figs.
- Sedra, S. N. & Michael, M. I. The metamorphosis and growth of the hyobranchial apparatus of the Egyptian toad, Bufo regularis Reuss. J. Morph. 103 1958: 1-30 20 figs.
- Sedra, S. N. & Moursi, A. A. The ontogenesis of the vertebral column of *Bufo regularis* Reuss. Čs. Morfologie 6 1958: 7–32 37 figs.
- Selman, G. G. The forces producing neural closure in amphibia. J. Embryol. exp. Morph. 6 1958: 448–465 1 pl. 5 figs.
- (2) Selman, G. G. An ultra-violet light method for producing haploid amphibian embryos. J. Embryol. exp. Morph. 6 1958: 634-637.
- Seno, T. & Nieuwkoop, P. D. The autonomous and dependent differentiations of the neural crest in amphibians. Proc. Akad. Wet. Amst. 610 1958: 489-498 1 fig.
- Sentein, P. Action du phénol sur les mitoses de segmentation des oeufs d'amphibiens, Acta anat. 34 1958 : 201-234.
- Serventy, D. L. Ludwig Glauert-Museum Director and Naturalist. [A Biography.] W. Aust. Nat. 5 7 1957: 148-165 photograph.
- **Sexton, O.** Observations on the life history of a Venezuelan frog, *Atelopus cruciger*. Acta biol. venezuel. **2** 21 1958: 235–242 1 fig.
- Shannon, F. A. & Humphrey, F. L. A discussion of the polytypic species, *Hypopachus oxyrrhinus*, with a description of a new sub-species. Herpetologica 14 1958: 85-95 1 fig.
- Shapiro, B. E. Influences of the salinity and pH of blastocoelic perfusates on the initiation of amphibian gastrulation. J. exp. Zool. 139 1958: 381-3964 figs. 2 pls.
- Sharp, J. E. The vertebrates of the Dernford district. A Survey of Dernford Fen, Sawston, Cambridgeshire 1950-58 1958: 42-44.

Shaver, J. R., see Barch, S. H.

Shaw, C. E. Southern salamanders are slender. Zoonooz 31 1958: 10-11 2 figs.

Shaw, F. H., see Briner, G. P.

Shaw, F. H., see Simon, S. E.

Shimozawa, A. Histological studies of corium in tadpoles of Rana catesbiana. Zool. Mag. Tokyo 67 1958: 380–382 2 pls.

Sibabin, K., see Kleerekoper, H.

Sickel, W., see Pilz, A.

Sidman, R. L. & Wislocki, G. B. Histochemical observations on rods and cones in retinas of vertebrates. J. Histochem. Cytochem. 2 6 1954: 413-433 4 pls.

Silva, P. H. D. H. de. Notes upon the range of body colour variation in the Ceylon Rhacophoridae. Part I Rhacophorus leucomystax maculatus (Gray) and Rhacophorus cruciger (Blyth). Spolia zeylan. 28 1958: 151-156 l pl.

Simon, S. E., Shaw, F. H., Bennett, S. & Muller, M. The relationship between sodium, potassium, and chloride in amphibian muscle. J. gen. Physiol. 40 1957: 753-777 9 figs.

Simon, S. E., see Briner, G. P.

Simons, J. R. The blood pressure and the pressure pulses in the arterial arches of the frog (Rana temporaria) and the toad (Bufo bufo). J. Physiol. 137 1957: 12-21 5 figs.

Singer, M. The regeneration of body parts. Sci. Amer. 199 1958: 79-88 figs.

(1) Sirakami, K. Studies on the "double nucleus" in the early developmental stage of *Bufo vulgaris*. 1. Cyto-embryological considerations on its possible participation in the normal course of cell multiplication. Embryologia 3 1 1956: 1-21 7 figs.

(2) Sirakami, K. Studies on the "double nucleus" in the early developmental stage of Bufo vulgaris. II. On the frequency of double nucleus under various experimental conditions, with special references to the oxygen concentrations of culture media. Embryologia 3 1 1956: 37–56 3 figs.

Sjodin, B. A. & Mullins, L. J. The action potential of frog nerve as affected by temperature, narcosis, and stimulation frequency. J. cell. comp. Physiol. 51 1958: 425-438 6 figs.

Skjelkvale, L., Nieder, K. & Huf, E. G. Metabolic studies on isolated epithelium of frog skin. Va. J. Sci. N.S. 9 1958: 437.

Skowron, S. & Roguski, H. Regeneration from implanted dissociated cells. I. Regenerative potentialities of limb and tail cells. Folia biol., Kraków 6 1958: 163–173 11 figs.

Skowron-Cendrzak, A., see Konieczna-Marczynska, B.

Slevin, J. R., see Anon (6).

Smith, H. M. & Chrapliwy, P. S. New and noteworthy Mexican herptiles from the Lidicker collection. Herpetologica 13 1958: 267–271.

Smith, H. M. & Grant, C. Noteworthy herptiles from Jalisco, Mexico. Herpetologica 14 1958: 18-23 2 figs.

Smith, H. M., see Poglayen, I.

Smith, M. A., see Grandison, A. G. C. (1).

Smith, P. W. & Hensley, M. M. Notes on a small collection of amphibians and reptiles from the vicinity of the Pinacate Larva Cap in Northwestern Sonora, Mexico. Trans. Kansas Acad. Sci. 61 1958: 64-764 figs.

Smith, S. Induction of triploidy in the South African clawed frog, Xenopus laevis (Daudin). Nature, Lond. 181 1958: 296.

Smith, S., see Elsdale, T. R.

Sochurek, E. Zur Verbreitung der Lurche und Kriechtiere in Burgenland. Burgenland. Heimat. Blatt. 20 1958: 149–151.

Solomon, A. K., see Durbin, R. P.

Sparrow, A. H., see Jakowska, S.

Speidel, C. C. Motion pictures of some changes in cells induced by X-ray treatments of tadpoles and tetrahymenae. Biol. Bull. Woods. Hole 115 1958: 322. [Abst.]

Spirito, A. & Bondi, C. Sul trapianto di abbozzi di *Triton* su embrioni di *Rana esculenta* e possibilità di differenziamento e di maggior resistenza alla distruzione a seguito di un nuovo metodo sperimentale. R.C. Acad. Lincei (8) 24 1958: 100-102 1 pl.

Springmuller, V. & Concha, J. Método para estabilizar y hacer reproducibles las medidas de consumo de oxígeno en animales de pequeña talla. Invest. zool. chil. 4 1958: 138-145, 2 figs. [English summary.]

Stadhouders, A. M., see Oordt, G. J. van.

Stagg, M. W., see Goss, R. J.

(1) Stärk, O. J. Entwicklung der Gonaden und Geschlechtszellen bei Triton alpestris, cristatus und taeniatus, mit besonderer berücksichtigung ihrer verschiedenheiten. Z. Zellforsch. 41 1955: 285–334.

(2) Stärk, O. J. Die extragonalen Geschlechtszellen von *Triton alpestris* und degenerative Veränderungen ihrer Kerne. Verh. dtsch. zool. Ges. 1957 [1958]: 454–469 15 figs.

Stefani, R. de. Sviluppo e differinziamento dell' ipofisi in girini di Rana dalmatina BP trattati con tiroxina e metilticuracile. Atti Ist. veneto Sci. mat. nat. 116 1958: 99-111 3 pls.

Sternfeld, R., see Anon (2).

Stevenson, H. M. A record of Hemidactylium scutatum in Florida. Copeia 1958 No. 1: 49.

Stewart, M. McB. Seasonal variation in the teeth of the two-lined salamander. Copeia 1958 No. 3: 190-196. 3 figs.

Stieve, H. Die Abhängigkeit der Schwelle und des Aktionisstroms isolierter markhaltiger Nervenfasern von Temperatur und Stoffwechsel. Z. Naturf. 13B 1958: 96–108 6 figs.

Stille, W. T. The water absorption response of an anuran. Copeia 1958 No. 3: 217-218.

Stine, C. J. & Fowler, J. A. The Hellbender. Welch Biology General Science Digest 6 1956: 7-8 2 figs. Sti Ruib

[19

Sti

resist

nucle ampl Ams

(3) in th 243.

St

newi

(2

St resto retir Zool

St

St

Ran

diere 1958

dev

J. T Shom tran 139

S the Wis

8

of 103 eor

2nd Est 4 I

ou 34 Stinson, B. D. Chemical enhancement of radioresistance in regenerating and developing limbs of urodele larvae. J. Morph. 103 1958: 387-433 18 figs.

Stipanincic, P. N. & Reig, O. A., see Hecht, M. K. & Ruibal, R.

(1) Stolk, A. Special condition of the interphase nucleus in normal and tumorous cells of fishes, amphibians and reptiles IA-IB. Proc. Akad. Wet. Amst. 61C 1958: 395-416 12 figs.

(2) Stolk, A. Tumours of amphibians. IV A-B. Proc. Akad. Wet. Amst. 61C 1958: 610-630 15 figs.

(3) Stolk, A. A transplantable fibroma of the skin in the newt *Triturus taeniatus*. Experientia 14 1958: 243.

Stone, L. S. Inhibition of lens regeneration in newt eyes by isolating the dorsal iris from the neural retina. Anat. Rec. 131 1958: 151-171 13 figs.

Stone, L. S. & Gallagher, S. B. Lens regeneration restored to iris membranes when grafted to neural retina environment after cultivation in vitro. J. exp. Zool. 139 1958: 247-258 l pl.

Storm, R. M., see Ferguson, D. E.

Streett, J. S., Jr. The relation between liver weight and degree of restoration after partial hepatectomy in Rana pipiens. Texas J. Sci. 10 1958: 236-243.

Stettler, P. H. De verzorging van planten en dieren in het terrarium I. Aquarium, Den Haag 28 1958: 148-152.

Su, J. & Tsai, C. Effects of colchicine on the early development of the frog, Rana limnocharis. Quart. J. Taiwan Mus. 11 1958: 87-101 4 pls.

Subtelny, S. The development of haploid and homozygous diploid frog embryos obtained from transplantations of haploid nuclei. J. exp. Zool. 139 1958: 263–298 9 figs. 4 pls.

Sundararajan, T. A., see Sampath Kumar, K. S. V.

Suznki, T., see Csapo, A.

Suzuki, H. K. A study of leg length variations in the wood frog, Rana sylvatica Le Conte. Trans. Wisc. Acad. Sci. 46 1958: 299-303.

Swann, M. M. & Mitchison, J. M. The mechanism of cleavage in animal cells. Biol. Res. 33 1958: 103-135 9 figs.

Sweetman, H. L. The principles of biological control. Dubuque, Iowa 1958: i-xii, 1-560 328 figs.

Swinton, W. E. Fossil Amphibians and Reptiles. 2nd. edn. Lond. 1958: 1-118.

Szabó, I. Contributions à l'herpétofaune de la montagne Dunazug entre Szentendre, Visegrád et Esztergom. Állat. Közlem. 45 3-4 1956 : 123-131 4 pls. 1 fig. [French summary.]

Szejnberg, A., see Lunenfeld, B.

Takagi, Y. The conduction velocity in intracutaneous nerve fibres of the frog. Jap. J. Physiol. 8 1958 340-346 5 figs.

Takayanagi, H., see Yanai, T.

Takeuchi, A. The long lasting depression in neuromuscular transmission of frog. Jap. J. Physiol. 8 1958: 102-113 14 figs.

Takeuchi, N. The effect of temperature on the neuromuscular junction of the frog. Jap. J. Physiol. 8 1958: 391-404 14 figs.

Takeuchi, T. Effects of boric acid on the development of the eggs of the toad, *Bufo vulgaris formosus*. Sci. Rep. Tôhoku Univ. (4) 24 1958: 33-43 20 figs.

Taleisnik, S., see Astrada, J. J.

Tamura, J., Yashiki, K., Kondo, E. & Numata, J. On the commissural systems of the cerebellum in Megalobatrachus japonicus. Okajimas Folia Anat. Jap. 29 1955: 1-5 17 figs.

Tandler, C. J. The localization of inorganic phosphate in the occytes of *Bufo arenarum* heterogeneity of the nucleoli. Exper. Cell. Res. 14 1958: 408-413 2 figs.

Tanner, W. W. Herpetology of Glen Canyon of the Upper Colorado River Basin. Herpetology 14 1958: 193-195.

Tanner, W. W., see Hansen, A. M.

Tansley, K., see Dobrowolski, J. A.

Taurand, G. Sur l'ovulation des amphibians anoures. Bull, mens, Soc. Univ. Lyon 27 1958: 225.

(1) Taylor, E. H. Additions to the known herpetological fauna of Costa Rica with comments on other species. No. III. Univ. Kans. sci. Bull. 39 1958: 3-40 17 figs.

(2) Taylor, E. H. Notes on Costa Rican Centrolenidae, with descriptions of new forms. Univ. Kans. sci. Bull. 39 1958: 41–68 13 figs.

Taylor, E. H. & Elbel, R. E. Contribution to the herpetology of Thailand. Univ. Kans. sci. Bull. 38 1958: 1033-1189 36 figs.

Taylor, E. H. & Weyer, D. Report on a collection of amphibians and reptiles from Harbel, Republic of Liberia. Univ. Kans. sci. Bull. 38 1958: 1191-1230.

Tencer, R. Étude autoradiographique de l'incorporation de <sup>14</sup>CO<sub>2</sub> dans des gastrulae d'Axolotl, J. Embryol. exp. Morph. 6 1958: 117-123, 3 figs.

Ten Kate, I. B., see Kappers, J. A.

Thomas, L. J., Jr. An antagonism in the action of calcium and strontium ions on the frog's heart. J. cell. comp. Physiol. 50 1957: 249-264 6 figs.

Thomas, R., see Jullien, A.

Thompson, E. F., see Martof, B. S.

Thoraton, C. S. The inhibition of limb regeneration in urodele larvae by localized irradiation with ultraviolet light, J. exp. Zool. 137 1958: 153-179 8 figs.

**Tihen, J. A.** Comments on the osteology and phylogeny of ambystomatid salamanders. Bull. Fla Acad. Sci. 3 1958: 1-50 11 figs.

Tisserand, M., see Jullien, A.

Toivonen, S. The dependence of the cellular transformation of the competent ectoderm on temporal relationships in the induction process. J. Embryol. exp. Morph. 6 1958: 479–485, 1 pl. 2 figs.

Tokin, B. P. & Filatova, A. G. On the breaking of morphogenetic processes of animals having transplanted tumours and under the influence of cancer causing agents. Vest. Leningr. Univ. No. 15 1958: 59-63.

Toma, V., see Pora, E. A.

Töndury, G., see Schenk, R.

Trethewie, E. R. & St. Clair, W. A. Effect of extracellular potassium on the intracellular action current in the beating heart in the intact toad. Aust. J. exp. Biol. med. Sci. 36 1958: 441-445 3 figs.

Triplett, E. L. The development of the sympathetic ganglia, sheath cells, and meninges in amphibians. J. exp. Zool. 138 1958: 283-308 11 figs.

Tsai, C., see Su, J.

Tschumi, P. A., see Hughes, A.

Turcotte, M. R., see Fulchiero, E.

Turner, F. B. Life-history of the western spotted frog in Yellowstone National Park. Herpetologica 14 1958: 96–100.

Tyler, M. J. On the diet and feeding habits of the edible frog (Rana esculenta Linnaeus). Proc. zool. Soc. Lond. 131 1958: 583 5 figs.

Ugurgieri, B., see Baldisserotto, A.

Ultangi, J. C. On the morphology of five new species of Nyctotherus found in frogs and toads from Dharwar. J. Univ. Bombay N.S. No. 43 1958: 50-64 13 figs.

Umezawa, S.-I. Role of intracellular phosphate ions on contraction and coagulum formation of an isolated frog muscle fibre. Annot. Zool. jap. 31 1958: 127-135 2 pls.

Uranga, J. Variations saisonnières de l'action antidiurétique de l'encéphale et de l'hypophyse du Crapaud. C.R. Soc. Biol. Paris 152 1958: 186–187.

Vainio, T. An approach to the problem of embryonic induction. Exper. Cell. Res. 15 1958: 184-190 3 figs.

Vandebroek, G. Les moments cruciaux de l'evolution des poissons jusqu'à l'homme. Rev. Quest.sci (5) 16 1955 508-537 13 figs.

Vandevelde, F. Kikvorsen: koelbloedige kannibalen beschermd bovendien. Lacerta 16 1958: 62-63.

Van Dyke, R. H. & Detwiler, S. R. Further studies on ear and capsule development in *Amblystoma*. Anat. Rec. 131 1958: 61-80 12 figs.

Veit, O. Über eine chordaähnliche Umbildung des Darmdaches vor der Praechordalplatte bei Rana fusca. Gegenbaurs Jb. 99 1958: 257-261 1 pl. 4 figs.

Verbaak, K. J. Schuimnestbouwers onder de kikkers. Lacerta 16 1958: 83-85.

(Vereshchagin, N. K.) [Systematic review of sea animals of the Caucasian isthmus.] IV Amphibia. Zhivot. Mir 5 1958: 281-286 figs. [In Russian.]

Viaud, G. & Dreyfus, E. Le galvanotropisme des tétards de Rana temporaria. La force électromotrice d'opposition produite par ces animaux et sa signification. C.R. Soc. Biol., Paris 151 1958: 1227-1231 2 figs.

Vivien, J. H. & Schott, J. Contribution à l'étude des corrélations hypothalamo-pituitaires chez les Batraciens. Le contrôle de l'activité gonadotrope. J. Physiol. Paris 50 1958: 561-563.

(1) Voitkevich, A. A. A peculiar reduction of the shin bone in the doubling of the limb in frog in nature. Dokl. Akad. Nauk SSSR (Transl.) Biol. Sci. 118 1958: 28-31 2 figs. [See also: C.R. Acad. Sci. U.R.S.S. 118 4 1958: 841-844. In Russian.]

(2) Voitkevich, A. A. Spatial relations between true limbs and accessory limbs developing under natural conditions. Dokl. Akad. Nauk SSSR (Transl.) Biol. Sci. 120 1958: 373-376 3 figs. [See also: C.R. Acad. Sci. U.R.S.S. 120 2 1958: 434-437 3 figs. In Russian.]

Volkin, E., see Finamore, F. J.

Volpe, E. P. Interspecific gene exchange between Fowler's and the Southern toad. Anat. Rec. 131 1958; 605–606.

Volpe, E. P. & Harvey, S. M. Hybridization and larval development in *Rana palmipes* Spix. Copeia **1958** No. 3: 196-207 5 figs.

Volpe, E. P., see Anderson, S. C.

Volpert, E., Michel, R. & Roche, J. Sur la concentration d'iodures et d'hormones thyroidiennes par l'ovaire de la Grenouille (Rana temporaria L.). C.R. Soc. Biol. Paris 152 1958: 406-408.

Vonderahe, A. R., see Peters, J. J.

Waddington, C. H. A note on the effects of some cytotoxic substances on amphibian embryos. J. Embryol. exp. Morph. 6 1958: 363-364 1 pl.

(1) Wager, V. A. The Pond reed frog. Afr. wild Life 12 1958: 55-57.

(2) Wager, V. A. The spotted burrowing frog. Afr. Wild Life 12 1958: 201-205 4 figs.

Wahlert, G. v. Biogeographische und ökologische Tatsachen zur Phylogenie amerikanischer Schwanzlurche. Zool. Jb. (Syst.) 85 1957: 253–282 2 figs.

Warren, R. D., see Pylka, J. M.

Warwick, T. Observations on the spawning site of the common frog, Rana temporaria temporaria L. Proc. R. phys. Soc. Edinb. 26 1958: 69-75.

Wasserman, A. O. Relationships of allopatric populations of spadefoots (genus Scaphiopus). Evolution 12 1958: 311–318 2 figs.

Webb, R. G. Frogs of the genus Eleutherodactylus from Coahuila, Mexico. Herpetologica 13 1958: 263–264.

Weinberg, J. A., see Wilson, L. D.

Weisel, G. F. The comparative effects of freshwater and marine teleost pituitary on the water balance of frogs. Copeia 1958 No. 2: 86-91.

Weiss, I., see Hertwig, G.

Wessing, A. & Claes, H. Die Herstellung von orientierten Schnittpräparaten dotterreicher Objekte. Die Verarbeitung von Amphibieneiern. Z. Wiss. Mikr. 63 1958: 477–483 6 figs. W W for e

[19

d'Ala

W

933-W reco

pitro Aca hyp

W

ehro mier 25 f

bioc pole

V

tion Soci ten Stu

[Fr

in dts So

sp

ec 81

Z

les

00.

he

re.

18

en

ler

al.)

R.

08.

58:

nd eia

m-

es

4.).

me J.

ild

g.

he

Z-

ite

ric

us

8:

h-

er

on te. Wettstein, L. Quelques notes concernant l'élevage d'Alytes obstetricans Laurenti. Rev. verviét. Hist. nat. 15 1958: 69-72.

Weyer, D., see Taylor, E. H.

Wilber, J. F., see Schotté, O. E.

Wilkie, D. R., see Jewell, B. R.

Williams, D. D. Rearing newts in the laboratory for experimental purposes. Nature, Lond. 181 1958: 933-934.

Williams, K. L. & Chrapliwy, P. S. Selected records of amphibians and reptiles from Arizona. Trans. Kansas Acad. Sci. 61 1958: 299-301.

Wills, J. P., see Huf, E. G.

Wilson, J. W. & Archdeacon, J. W. The effect of pitressin on lymph drainage in the frog. Trans. Ky Acad. Sci. 19 1958: 67–69.

Wilson, L. D., Weinberg, J. A. & Bern, H. A. The hypothalamic neurosecretory system of the tree frog Hyla regilla. J. comp. Neur. 107 1957: 253–272.

Wischnitzer, S. A study of the lateral loop chromosomes of amphibian occytes by phase contrast microscopy. Amer. J. Anat. 101 1957: 135–167 25 figs.

Wislocki, G. B., see Sidman, R. L.

Wisniewski, W. L. Characterization of the parasitofauna of an eutrophic lake. [Parasitofauna of the biocoenosis of Druzno Lake-part I.] Acta Parasit. polon. 6 1958: 1-64 map.

Witschi, E., see Chang, C.

Witschi, E., Foote, C. L. & Chang, C. Y. Modification of sex differentiation by steroid hormones in a tree frog (*Pseudacris nigrita triseriata* Wied.) Proc. Soc. exp. Biol., N.Y. 97 1958: 196-197.

Wittenberger, C. & Fabian, N. Recherches sur le temps réflexe dans les réflexes spinaux de la grenouille. Stud. Cercet. Biol. Bucuresti 8 1957: 409-417 2 figs. [French, etc. summaries.]

Wittner, M. & Rosenbaum, R. M. Resistance and susceptibility to high oxygen pressures in the early development of the frog, Rana pipiens. Physiol. Zool. 31 1958: 294-303 7 figs.

Woellwarth, C. v. Zur Frage der Induktionsfelder in jungen Embryonalstadien der Amphibien. Verh. dtsch. zool. Ges. 1957 [1958]: 424–430 8 figs.

Woodbury, A. M. Notes on the toads of eastern South Dakota. Copeia 1958 No. 2: 149-150.

Woodford, J. Breeding behaviour of the redspotted newt. Ont. Field Biol. 10 1956: 1-7.

Wright, P. A., see Rondell, P. A.

Wyllie, J. H., see Denton, E. J.

Yamada, T. Induction of specific differentiation by samples of proteins and nucleoproteins in the isolated ectoderm of *Triturus*-gastrulae. Experientia 14 1958: 81-87 11 figs.

Yamamoto, T. Double embryos induced by split organizer of toad eggs treated with potassium oxalate. Zool. Mag., Tokyo 67 1958: 84–87 2 figs.

Yanai, T. Notes on the hatching glands of the frog, Polypedates buergeri. Annot. zool. jap. 31 1958: 222-224 3 figs.

Yanai, T. & Takayanagi, H. Notes on the hatching glands of the frog, *Hyla arborea japonica*. Annot. zool. jap. 31 1958: 39–42 3 figs.

Yashiki, K., see Tamura, J.

(1) Yoshida, M. Acetylcholine hydrolizing enzyme in the isolated inner and outer segments of the receptor cell of frog retina. Jap. J. Physiol. 8 1958: 155-159 2 figs.

(2) Yoshida, M. Histochemical studies on alkaline phosphatase of the salamander retina in the late embryogenesis. Jap. J. Physiol. 8 1958: 31-406 figs.

Young, J., see Money, W. L.

Zaleska, A., see Morawska, B.

Zelený, A. & Kozák, J. Action of acetone on the tonic muscle of the frog. Nature, Lond. 181 1958: 1340-1341 2 figs.

Zieleniewski, J., see Ber, A.

Ziemann, C. Vergleichende volumetrische Untersuchungen an den Zellkernen und dem Zellplasma verschiedener Amphibienarten. Z. mikr. anat. Forsch. 64 1958: 18-29 3 figs.

Ziemann, C., see Hertwig, G.

Zweifach, B. W., see Nagler, A. L.

(1) Zweifel, R. G. Results of the Puritan-American Museum of Natural History Expedition to Western Mexico. 2. Notes on reptiles and amphibians from the Pacific coastal islands of Baja California. Amer. Mus. Novit. No. 1895 1958: 1-17 1 fig.

(2) Zweifel, R. G. Results of the Archbold Expeditions. No. 78. Frogs of the Papuan hylid genus Nyctimystes. Amer. Mus. Novit. No. 1896 1958: 1-51 21 figs.

#### II.-SUBJECT INDEX

References to the "Titles" are by the name(s) of the Author(s) printed in clarendon type.

#### GENERAL LITERATURE AND HISTORY

Text-books.—Fossil Amphibians and Reptiles, Swinton.

General Works,—Importance of amphibian remains in stratigraphy, Lapparent, Claracq & Nougaréde.— Evolution of Amphibia, Schmalhausen, (4).

Bibliography.—Of Mookerjee, see Bhaduri.—Distributional reports on amphibians of E. Canada and zoogeography, Bleakney, J. S.—List of scientific publications by L. Glauert, Serventy.

Biography.-Of Ludwig Glauert, Serventy.

Museums and Collections.—Catalogue of amphibian and reptile types in the Chicago Natural History Museum, Marx.

Nomenclature.—Suppression of Acrodytes Fitzinger, 1843, of tibiatrix Laurenti, 1768 as published in the combination Hyla tibiatrix and interpretation of Rana venulosa Laurenti, 1768, Opinion 520.

Obituary.—An appreciation of Karl P. Schmidt, Anon [E. H. C.]—Of Joseph R. Slevin, Anon (6),—Of H. K. Mookerjee, Bhaduri,—Of K. P. Schmidt, see Cei (3),—Of Malcolm A. Smith, Grandison (1).

Reviews.—Normal table of Xenopus laevis edited by P. D. Nieuwkoop and J. Faber, Anon (1).—Die Reptilien und Amphibien Mitteleuropas by R. Sternfeld (2nd edition), Anon (2).—A Field Guide to Reptiles and Amphibians of Eastern North America by Roger and Isabelle Conant, Adler (1).—Terrarienkunde. 2. Teil Lurche by W. Klingelhöffer, Gans.—Amphibia of Ceylon by P. Kirtisinghe, Grandison (2).—Vertebrates of the United States by Blair, Blair, Cagle, Brodkorb and Moore, Hecht (1).—A short note on the finding of Anuran in the "porphyritic complex of the extraandean Patagonia" with considerations of Geological compositions of same by P. N. Stipanincic and O. A. Reig, Hecht & Ruibal.—Vida de batracios y reptiles sudamericanos by M. A. Freiberg, Oliver (1).—Sounds of N. American frogs by C. M. Bogert, Oliver (2).—Zoogeography: the geographical distribution of animals by P. J. Darlington, Savage.

#### STRUCTURE AND HISTOLOGY

Alimentary System.—Comparison digestive systems of vertebrates, Baer,—Distribution of hepatic glycogen in Pleurodeles walthis, Bellec (1).—Histology of oesophagus of Rana chensinensis, Eksaeva.—Buccopharyngeal mucous membrane of urodeles and caecilian, Elkan, (1).—Development and structure of the pancreas in Amblystoma, Frye.—Masticatory musculature of Cryptobranchus japonicus, Schumacher (1).

Biochemistry.—Hepatic glycogen in Pleurodeles waltivi, Bellee (1).—Histochemical location of acid phosphatase in dendrites of Rana, Cerf & Chacko.—Histochemical studies on the distribution of phosphatases and mucopolysaccharides in the oviduct of the toad Bufo, Chu & Chin.—Nucleic acid content of liver nuclei compared, England & Mayer.—Skin extracts of Leptodactylus and their biochemical use, Erspamer.—Mononucleotide composition of ovarian nucleic acids of Rana catesbeiana, Finamore & Volkin.—Glycogen contents of ovarian, body cavity and uterine eggs of Rana pipiens, Fitch, & Merrick.—Localisation of phosphorus and the site of calcium binding in yolk protein of frog's egg, Flickinger & Schjeide.—Chemistry of venoms produced by Bufo sp., Frèrejacque.—Some constants of toad fat, Fromm.—Mucopolysaccharides in the kidney of Rana, Geyer, G.—Fluorescent pterines and flavines in eye of Rana ingromaculata, Hama & Horiuchi.—Histochemical tests and pigment extractions from Plethodon cinereus, Hertzler.—Histochemical study of peroxidase, Kawasaki.—Amino acid composition of Xenopus laevis skin gelatin, Leach.—Cytochemical nature of cytoplasmic granules in amphibian erythrocytes, Nagatani.—Morphological study of lipids in Rana oogenesis, Nath, Gupta & Kaur.—Protein synthesis in Triturus oocytes, Pantelouris.—Histochemistry of rods and cones in frog and Amphiuma retinas, Sidman & Wislocki.—Inorganic phosphate localised in nucleoli of oocytes of Bufo arenarum, Tandler.—Histochemical studies on alkaline phosphatase of salamander retina, Yoshida (2).

Circulatory System.—General points of interest in the amphibian circulatory system, Baer.—Blood

system in the hypophysis of anurans, Cruz.—Vascularisation of the respiratory surfaces in Ambystoma mexicanum, Czopek.—Spleen modification and development in Taricha torosa, De Lanney.—Methylthiouracil on blood corpuscles of Rana ridibunda, Grabie-Broda.—Histology of the heart of adult Triturus cristatus, Gyévai.—Evolution of a potential in the ventricle wall and Purkinje fibres of the frog heart, Lacquet.—Comparative researches on the catalasic activity of blood in the various classes of Vertebrata, Liotti.—Classification of cytoplasmic granules in amphibian crythrocytes, Nagatani.—Retinal blood supply in Anura, Polyak.

Cytology.—Observations on spinal ganglion cells of Rana pipiens, Anderson & Breemen.—Heteroploid embryos of Pleurodeles walthii, Beetschen (2).—Mitotic anomalies at the site of origin of genetic aneceploids of Pleurodeles, Beetschen (3).—Distribution of hepatic glycogen in Pleurodeles walthi, Bellee (1).—Research into the amount of cellular nephrostomial tissue incorporated in the development of the female genital opening of Rana dalmatina, Cambar & Dechambe.— Electron microscope studies of mitochondria and the Golgi substance in cells of amphibian larvae, Carasso. -Histological structures of the toad oviduct after the laying of eggs, Chu & Chin.—Structure of the oesophagus of Rana chensinensis, Eksaeva.—Buccopharyngeal mucous membrane of Schistometopum and urodeles, Elkan (1).—A mutation reducing nucleolar number in Xenopus laevis, Elsdale, Fischberg & Smith.—Comparative nucleic acid content of liver nuclei, ova nuclei and spermatozoa in Rana pipiens, England & Mayer.—Chromosome number constancy in larval salamanders, Fankhauser & Dluhy.— Cytomorphosis of Rohon-Beard cells during embryonic and larval development of Rana, Filippio.—Amino acids, arginine and sulphydriles proteins in pancreatic cells of salamanders, Gabe.—Heart of adult Triturus cristatus, Gyévai.—Nuclear size in experimental hybrid toads, Hertwig, Ruhland & Weiss.—Histological differentiation in the marginal zone of amphibian eggs, Ikushima.—The Golgi apparatus of the cells of the amphibian pancreas, Kanwar.—Mast cells in the choroid plexus of the axolotl, Kappers, Ten Kate & Bruyn.—Electron micrograph study of the crystalline lattice in yolk platelets of *Triturus pyrrhogaster*, Karasaki & Komoda.—Electron microscope study of lampbrush chromosomes of Amphibia, Lafontaine & Ris.—The cytoplasmic inclusions of the ageing neurones of Rana tigrina, Malhotra.—Chromosomes from the spermatogonial and oogonial metaphase plates of Bufo melanostictus, Natarajan.—Cytoplasmic granules in erythrocytes stainable with carcinogenic azo-dyes, Nagatani.—Chromosomes of *Triturus boscai* and of the gonads of larval salamanders, Sacarrão.— Anterior pituitary and thyroid gland of Xenopus laevis, Saxén.—Teeth of Megalobatrachus, Schmidt, W. J. (2).—Histological studies of corium in tadpoles of Rana catesbiana, Shimozawa.—Histochemistry of rods and cones of Amphiuma and frog, Sidman & Wislocki. —Histology of frog isolated epithelium cells, Skjelk-vale, Nieder & Huf.—Nuclear degeneration in the gonad of Triturus under external influence, Stärk (2).—Special condition of the interphase nucleus in normal and tumorous cells of fishes, amphibians and reptiles, Stolk (1).—Abnormal tumours and normal skin development in Triturus taeniatus, Stolk (2).— Commissural systems of the cerebellum of Megalo-batrachus japonicus, Tamura, Yashiki, Kondo & Numata.—Histological effects of ultra-violet light irradia
—Cyte
of Hyl
of the
Nature
chrom
T. vire
analys
Ziema

japoni in †

> Amph ment sylvate

other

Baer.
Gla
basop
ders,
tive a
morph
Prum
Yanai
Yanai

Interest of Dislarization Czope reflect terization Kobli of am positi Trituthe sl

and s

moni
Deme
stage
an i
casti
tenar
Foot
frog
Varie
ampl
orien
& Cl

vallic vary (1). soble paratures Of th Stru micr irradiation of the limb tip of urodele larvae, Thornton.
—Cytology of the hypothalamic neurosecretory cells of Hyla regilla in normal animals and after transection of the infundibular stalk, Wilson, Weinberg & Bern.—Nature of chiasmata in and size of, the lateral loop chromosomes of occytes of Trisurus pyrrhogaster and T. viridescens, Wischnitzer.—Comparative volumetric analysis of nuclei and cells of several amphibians, Ziemann.

Dentition.—Histology of teeth of Megalobatrachus japonicus, Schmidt, W. J. (2).—Arrangement of teeth in † Paracyclotosaurus, Swinton.

Excretory System.—Urinogenital systems of Amphibia; comparative account, Baer.—Development and degeneration of pronephros in Rana sylvatica tadpoles, Hurley.

Fat Body.—Saponification value, iodine value and other fat constants of Bufo marinus fat body, Fromm.

Glands.—Cutaneous glands present in amphibians, Baer.

Glands: Endocrine.—Histochemical tests on the basophil character of the pancreatic cells of salamanders, and salivary glands of snails, Gabe.—Quantitative analysis of thyroid development during metamorphosis of Nectophrynoides occidentalis, Lamotte & Prum.—Hatching glands of Polypedates buergeri, Yanai.—Hatching glands of Hyla arborea japonica, Yanai & Takayanagi.

Integument.—Analysis of the dorsal pigmentation of Discoglossus pictus, Bedate.—Structure and vascularization of the skin of larval Ambystoma mexicanum, Czopek.—Preserved dermal scutes of Laidleria reflecting the nature of the skin, Kitching.—Characterization and localization of frog skin cholinesterase, Kohlick.—Iodine accumulation in pigmented tissues of amphibian larvae, Koshida (1).—Amino acid composition of Xenopus laevis skin gelatin, Leach.—Triturus taeniatus adepidermal reticular network of the skin in normal and tumourous newts, Stolk (2).

Laboratory Technique.—A procedure for staining and sectioning heads of adult Anura, Baldaut, (2).— Isolation of an m-hydroxyphenyl-ethyl-trimethylammonium from the skin of Leptodactylus, Erspamer.— Demonstration of operation technique at yolk plug stage of determining the potency of the notochord as an inductor, Eyal-Giladi.—A new substance for casting amphibians for exhibition, Fendler.—Maintenance of gonads from Rana larvae in different media, Foote, C. L. & F. M. (1).—Sectioning heavily yoked frog eggs and calcified skin, Griffiths & Carter.—Various methods for anaesthetising and casting amphibians, Müller-Gehring.—The preparation of orientated section preparations of Amphibia, Wessing & Claes.

Myology.—M. depressor mandibulae in Bufo valliceps, Baldauf (1).—Glycogen phosphorylation in varying conditions in frog muscle, Cordier & Duveau (1).—Major features of myology of Centrolene prosoblepon, Eaton.—Musculature of Rana esculenta after parasitism by cystic helminths, Gianotti.—The structures of fast and slow muscle fibres in the frog, Gray.—Of the vocal sac region in Bufo alvarius, Inger, (1).—Structure of myocardium in frog studied by electron microscope, Khristoliubova,—Striated muscles in

frogs, Koloss.—Study on the head musculature of Cryptobranchus japonicus, Schumacher (1).—Morphology of urodele musculature, Schumacher (2).

Nervous System.—Cranial morphology of Bufo valliceps, Baldanf (1).—Organisation in aggregates of anterior neural plate cells of Triturus alpestris, Boternbrood.—Composition of the spinal ganglia of Xenopus, Hughes & Tschumi.—Mast cells in the choroid plexus of the axolotl, Kappers, Ten Kate & Bruyn.—Gomori-positive reactions of sulphydrite substances in the hypothalamus and hypophysis of the toad, Lasansky & Sabatini.—Ageing neurones of Rana tigrina; a cytological study, Malhora.—Effect on them of nerve section, Maturana.—Effect on nervous system of successive orthotopic limb grafts in Bufo vulgaris, Notarnicola.—Extrinsic and intrinsic factors in the development of the spinal cord and ganglia in Anura, Perri (1).—Anatomy of sympathetic rami in Rana pipiens, Pick.—Ultrastructure of Schmidt-Lauterman Clefts and shearing defects of the myelin sheath in soistic nerves of Rana, Robertson.—Comparative analysis of Anuran brains, Samtleben.—Commissural systems of the cerebellum of Megalobatrachus japonicus, Tamura, Yashiki, Kondo & Numata.—Anatomy of the hypothalamic neuroscertory system of Hyla regilla, Wilson, Weinberg, & Bern.

Osteology.—Osteology of amphibians, Baer.—Cranial morphology of Bufo valliceps, Baldauf (1).—Of Centrolene prosoblepon, Eaton.—Of species of Bolitoglossa, Hansen & Tanner.—Interclavicula of the first middle Triassic Labyrinthodont recorded from China (affinities with Heptasaurus and Mastodonsaurus), Huene (1).—The labyrinth and middle ear of Xenopus laevis, Kahmann & Frettlöh.—Tissue staining with particular reference to cartilage, Kawasaki.—Skeleton of a new amphibian from the Lower Triassic of S. Africa, Kitching.—Measurements of disarticulated Rana pipiens skeletons, Olson & Miller.—Morphology of urodele vertebral column; the transverse processes and ribs, Schmalhausen (2).—Nasolachrimal duct and septomaxilla of urodeles, Schmalhausen (3).—Metamorphosis and growth of the hypobrachial apparatus in Bufo regularis, Sedra & Moursi.—Skulls and skeletons of fossil Amphibia, Swinton.—The teeth, premaxillae, prevomers, parasphenoid, hypobranchium and vertebrae of the Ambystomidae, Tihen.

Reproductive System.—Morphological and histochemical changes in the testes of Rana pipiens after injections of gonadotrophins, Burgos & Ladman.—Rana dalmatina female genital opening and amount of pronephric material combined in its development, Cambar & Dechambe.—Experiments on the development of the genital opening after removal of the pronephros, Dechambe.—Cellular changes in gonads of frog larvae on synthetic media for varying periods of time, Foote & Foote (2).—Effect of testosteron-propionate on sex differentiation and structure of reproductive system in Hynobius, Hanaoka (3).

Respiratory System.—Bucco-pharyngeal mucous membrane of Schistometopum and urodeles, Elkan (1).

—Vascularization of respiratory surfaces in ontogeny of Ambystoma mexicanum, Gzopek.—Origin, development and changes of the gills and lungs of Kaloula borealis, Giang & Li.

Sense Organs.—Amphibian sense organs compared with those of other vertebrates, Baer.—Method of measuring the spectral absorption of photopigment in Xenopus rods, Dobrowolski, Johnson & Tansley.—Localisation of fluorescent pterines and flavines in the eye of Rana nigromaculata, Hama & Horiuchi—The labyrinth and middle ear of Xenopus laevis, Kahmann & Frettlöh.—Historical review of investigations of the eye and of the visual pathways and centres of the brain, Polyak.—Histochemistry of rods and cones in retinas of frog and Amphiuma, Sidman & Wislocki.

Teratology.—New type of polydactyly in Rana esculenta, Rostand (1).—Accessory limbs and reduction of shin bone in frogs, Voitkevich, (1).—Spatial relations between true limbs and accessory limbs developing under natural conditions, Voitkevich (2).

Tongue.—Tongue action in Hydromantes, Elkan (2).

#### PHYSIOLOGY

General Works.—Hypothetical interpretation to explain the physiological mechanism operating in the determination of bilateral symmetry in amphibian embryos, Levirup.

Biochemistry.-Distribution of inorganic phosphate in Bufo marinus muscle, Briner, Simon & Shaw.— Tests on frogs to determine whether the melanophore dispersing substance in human pregnancy urine is a 3-4-dihydroxybenzene derivative, Burgers Mighorst.—Effect of an oxygen lack on the phosphorylation of glycogen in frog muscles, Cordier & Duveau (1).—Effect of asphyxiation on the phosphorylation of the hepatic cells glycogen in Rana esculenta, Cordier & Duveau (2).—Phosphorylation of liver glycogen in Rana esculeata, Cordier, Buveau & Brandon.—Diminution of dispetidase activity by X irradiation, Coromaldi (1).—Evidence of a ribonucleoprotein concerned in the adhesion of cells and the binding of calcium to the cell surface, Curtis.—Vitamins A, B1, C and PP in new muscle regeneration, Degan & Asandei.—Peculiarity of carbohydrate metabolism of the bullfrog heart, Hashimoto, Morita & Matsuyama. -Histochemical reactions of the skin glands of Rana temporaria and Discoglossus pictus after injections of reserpine, Liébecq-Hutter & Bacq.—Temperature influence on relation of liver tissue respiration to glycogen content in Rana esculenta, Locker & Doneff. Temperature adaptation of cell enzymes in Rana temporaria, Mews.—Relation of melanophore-expandtemporaria, Mews.—Relation of melanophore-expanding hormone to hyperglycaemia in Xenopus laevis,
Munday.—Histochemical study of lipids in Rana
oogenesis, Nath, Gupta & Kaur.—Cytochrome-coxidase cycle in larval Bufo bufo, Petrucci.—Cytochemical investigation of the cortical cytoplasm of the
developing occyte of Rana pipiens, Rosenbaum.—
Biosynthesis of phosphoprotein in the frog, Sampath
Kumar et al.—Enzyme activity in the isolated inner and outer segments of the receptor cells of frog retina, Yoshida (1).

Body temperature.—Temperature economy in salamanders, Freytag (3).

Circulation.—Temperature effects on the electrocardiogram of Xenopus, Biörck & Johansson.— Reactions of blood circulation to irradiation, Bücker & Hanke.—Circulation of blood in hypophysis of anurans, Cruz. - Spectral absorption curve of haemoglobin of Xenopus laevis, Dobrowolski, Johnson & Tansley.—Histochemistry of the hearts of Triturus cristatus adults, Gyévai.—Carbohydrate metabolism of the isolated bullfrog heart, Hashimoto, Morita & Matsuyama.—Effect on the pulmonary circulation of introducing chemical solutions to the pulmonary eavity of the frog, Julien, Cardot & Moulin.-Frog heart rhythm and a cardiomotor substance of nervous origin, Jullien, Ripplinger & Cardot.-Arterial liga-Ripplinger, Cardot, Applinger & Cardot, —Arterial ligations and subsequent survival in frogs, Jullien, Ripplinger, Cardot & Baylé.—Cessation of the pulmonary circulation and the possibility of survival in the frog, Jullien, Ripplinger, Cardot, Baylé & Tisserand.—Various effects of cultivary of children. Various effects of solutions of sodium, potassium and calcium on the heart of larval Triton alpestris, Jullien, Ripplinger, Cardot & Marcot.-Action of alkaline solutions on the pulmonary circulation of the frog, Jullien, Ripplinger, Cardot & Moulin.—Modifications in lung circulation caused by alkaline solutions of sodium potassium and calcium introduced into the pulmonary cavity, Jullien, Ripplinger, Cardot, Moulin & Thomas.—Nervous activity and dynamics of frog heart, Kolatat, Kramer & Mühl.—Blood oxygen capacity in frogs, Leftwich.—Cellular membrane potentials and Ca and Na antagonism in frog heart, Lüttgau & Niedergerke.—Seasonal variation in drug sensitivity of Bufo marinus heart, Nayler.—Cutaneous vasomotor reflexes in the edible frog, Poczopko.-Blood pressure and pressure pulses in the arterial arches of Rana and Bufo, Simons.—Calcium and strontium antagonism in frog heart, Thomas.—Extra cellular potassium effect on beating heart of intact toad, Trethewie & St.Clair.

Coloration: Pigments.—Melanophores of Xenopus laevis, Bagnara (2).—Hormone control of anuran pigmentation, Bagnara & Neidleman.—Reaction of melanophores of Xenopus laevis to various substances, Burgers.—Light absorption and bleaching of visual purple in frog retina, Denton & Wyllie.—Spectral absorption curve of the photopigment in single rods of Xenopus laevis, Dobrowolski, Johnson & Tansley.—Migration of chromatophores from grafts on the flank of ghost hosts, Finnegan.—Study of absorption spectra of "Rana chrome I", Goto & Hama.—Fluorescent substances of pterin type in the adult skin of Bufo vulgaris formosus, Hama & Obika.—The red and black pigments of Plethodon cinereus, Hertzler.—Figmentation of Bufo bufo, Meng.—Colour variation in Bombina bombina and B. variegata, Michalowski.—Skin colour parallelism in plethodontid salamanders, Reynolds.

Digestion.—Fluid and electrolyte movement across the intestinal wall of Rana catesbiana, Chalfin, Cooperstein & Hogben.—Water flow through frog gastric mucosa, Durbin, Frank & Solomon.—Gastric acid secretion and chloride transport in frog, Durbin & Heinz.—Experiments on chloride transport across the stomach mucosal membrane, Heinz & Durbin.—Influence of low temperatures on the digestive rates of Triturus spp., Joly.—Modifications of the motivity of the gastro-intestinal tract of the frog during the year, Pora & Rejep.

Diseases.—Rana esculenta infected by Flavobacterium hydrophilum causing "red leg", Dubedout.—
Tumours induced in Triturus cristatus by implantation of methyl cholar.trene, Leone.—The origin, structure,

func verte in n and fibro skin spec Stoll cano

E

tive

bian of t

trea

Pola

by 1

Ten

Xen

[1

pad ribo emb Doll reac egge acro tem Ger Gian ager Reg fed and On read Ace neu & I

frog Ca

Eff

of n

Inf

flue gly Do fro the Jar enz che effe Me on Zal aug pho for lar

ten the ner tur ker

no-

rus ism

of

ary

ous

ga-

en,

on-

the

and

ris,

ica-

s of

the

alin

rog

gen

ane

art,

rug

ous.

rial

and

ctra.

tact

pus

iran

of ces.

sual

tral

rods

tion

a.—

The

enia.

lour *ata*, ntid

ross

lfin.

frog

stric rbin

TOBE

98 of

y of

ear,

cter-

tion

n.-

y.—

of the function and its manifestations in disease of the vertebrate visual system, Polyak.—Interphase nuclei in normal and tumorous cells of fishes, amphibians and reptiles, Stolk, (1).—Development of the multiple fibroma of the adepidermal reticular network in the skin of the newt Triturus tenniatus, Stolk (2).—Species specific fibromas in the skin of Triturus taeniatus, Stolk (3).—9, 10, dimethyl-1, and 2-bensantrazen as cancer causing substances, Tokin & Filatova.

Effect of Chemicals and Temperatures.-Comparative effects of phosphorylated nucleosides on amphibian morphogenesis, Ambellan.—The relative stability of the guanophore contracting agent under various treatments, Bagnara (1).—Thermal stadia is the development of Rana temporaria from South West Poland, Bebak,-Production of sterile female triploids by refrigeration of pleurodele eggs, Beetschen (3).— Temperature effect on the electrocardiogram of Xenopus, Biörck & Johansson.—HCG, FSH, LH, and LTH injections and their effect on testes and thumbpads of frogs Burgos & Ladman.—Action of desoxyribosenuclease on lampbrush chromosomes, Callan & Macgregor. -- Action of a thermal gradient on the embryonic development of Triturus alpestris, Cury & Dollander.—Contamination of the diphenylamine reaction for deoxyribonucleic acid in Rana pipiens eggs, Finamore.—Role of the corium in ion transport across isolated frog skins, Fleming.—Influence of temperature on the development of Xenopus laevis, Gerbes .- Effect of insecticide dust on Rana esculenta, Gianotti.—Sulphonylurea compounds as hypoglycemic agents in frogs, Gourley.—Influence of methylthioura-cil on the haematology of Anura, Grabie-Broda.— Regional distribution of catalase activity in tails of fed and starving Xenopus larvae, Hahn, H. P. van.— K tracer in the form KHCO<sub>3</sub> in tracing permeation and diffusion of ions in the sartorius muscle, Harris. On dehydration, critical temperature, activity and reactions to light of Eurycea lucifuga, Hutchison.— Acetyl choline potentiation by noradrenaline in the neuromuscular facilitation of frog leg muscle, **Hutter** & Loewenstein.—Eserine-acetylcholine interaction in frog muscle, Jenden.—Effect of tenfold increase of Ca++ in sartorius muscle fibres of frog, Jenerick.— Effect of temperature on the series elastic compound of muscle, Jewell & Wilkie.—Influence of low temperatures on digestion in Triturus spp., Joly.— Influence of temperature on the respiratory quotient of Rana esculenta skin, Locker.-Temperature influence on relation of liver tissue respiration to glycogen content in Rana esculenta, Locker & Doneff.—Ca and Na effect on heart contractions of frog, Lüttgau & Niedergerke.—Action of steroids on the active transport of sodium in frog skin, Maetz, Jard & Morel.—Temperature adaptation of cell enzymes in Rana temporaria, Mews.—Change in chemical structure of thyroxine analogues and their effect on Rana pipiens metamorphosis, Morey, Meltzer, Young & Rawson.—Influence of temperature on pulse rate of Xenopus and Rana, Morawska & Zaleska.—Influence of temperature in altering the augmenting effect of sodium azide on the energy rich phosphate content of toad embryos (Bufo bufo formosus), Moriwaki.—Temperature selection by larvae of Salientia, Rühmekorf (2).—Influence of temperature on breeding displays in newts, Scul-thorpe.—Effect of narcosis and temperature on frog nerve action potentials, Sjodin & Mullins.—Tempera-ture shock before fertilisation inducing triploidy later in Xenopus laevis, Smith, S.—Influence of temperature

on the threshould of medullated nerve fibres, Stieve.—Chemical enhancement of radio resistance in urodele larvae, Stinson.—Effect of temperature on the neuromuscular junction of frog, Takeuchi, N.—Cancer causing substances of a chemical, physical and biological nature, Tokin & Filatova.—Role of intracellular phosphate ions on contraction and coagulum formation in frog muscle fibre, Umezawa.—Progressive shift of the regional effect by heat treatment of bone marrow of Triturus, Yamada.—Effect of light on retinal enzyme activity, Yoshida (1).

Excretion.—Influence of diuretic hormones on Bufo arenarum, Becerra.—Experiment to show that the different regions of the primary ureter blastema possess at first the same morphogenetic potentialities, Cambar & Gipouloux.—Histochemistry of various parts of the kidney tubules and ureters of Rana, Geyer, G.—Electrical potential measurements on single nephrons of Necturus, Giebisch.—Role of thryoid in kidney development of Rana sylvatica tadpoles, development and degeneration of pronephros, Hurley.—Active sodium transport by the isolated toad bladder, Leaf, Anderson & Page.—Endotoxins in frog renal portal system and release of epinephrine, Nagler & Zweifach.—Seasonal variations in the effects produced by antidiuretics on Bufo arenarum, Uranga.

Experimental Studies .- Action of drugs on the potential in frog skin, Aguggini & Noseda.—Sperm elimination and hormonal injections in Bufo arenarum, Allende & Astrada.—Effects of sodium, potassium and azide on the ionic movements accompanying frog nervous activity, Asano & Hurlbut.-Gonadial responses of Triturus pyrrhogaster to methyltestosterone, Asayama & Matsuzaki.—Effect of estrogens on larval gonads of Hynobius, Asayama & Miyamori.-Effects of inhibitory solutions on sperm production in the toad, Astrada & Taliesnik.—The effect of guanophore contracting hormones on individual guanophores, Bagnara (1).—Precocious accumulation of glycogen in larval Pleurodele livers cultured in glucose, Bellec (2),-Sodium loss through the skin in Bufo after vasopressin injections, Bentley (1).—Effect of pituitrin on water transport in isolated Bufo bladder, Bentley (2).— Movement of sodium across the skin of the dehydrated Bufo marinus, Bentley (3).—Effect of acetylcholine-sterase on the surviving frog heart, Beznák.—Effect of penicillin on the metamorphosis of Bufo melanostictus, Bhattacharya & Medda.—Inactivation of newt larvae temporarily by benzimidazole and its alkyl derivatives, Billett (2).—The effect of acetylcholine in the absence of potassium on the myocardium of Rana temporaria, Borghgraef & Carmeliet.—Action of DCG on the active transport of sodium under the isolated skin of the frog: influence of calcium, Bourguet .-Effect of β-mercaptoethanol on morphogenesis in amphibian eggs, Brachet.-Effect on frog's electroretinogram of varying the amount of retina illuminated, Brindley (2).—Retinal responses to light in excised eye of Rana temporaria, Brindley (3).-Action of LH and FSH on spermiation in Bufo arenarum, Burgos & Houssay.—Effects of gonadotrophins on testes and thumb pads of normal and hypophysectomized Rana pipiens, Burgos & Ladman.—Effect of injections of benzopyrene into Triturus cristatus, Capuani (2).—Effects of benzopyrene injections on Triturus cristatus, Capuani (1).—Effects of asphyxiation and lack of oxygen on frog liver glycogen, Cordier,
Duveau & Brandon.—Influence of drugs on water
inhibition and ion loss by frog gastroenemius,

me in .

Poi

ma Ra

tra

Mu

mi

in

ace of

Sir

Css

im

log

me

po of

Gy

fac

ac

de

RAI

ree

m

fre

m

dr

ga ex

Po

Be Ba Co

in Ed no in tie re of

Eckhardt & Govier.—Effect of heat shock treatment of eggs of Triturus palmatus, T. vulgaris, on chromosome number of resulting embryos, Fischberg.—Curarizing effect of neomycin and its fractions in the frog, Fulchiero, Turcotte & Martin.—Hypophysectomixed efts treated with hormones and study of produced water drive, Grant & Grant,—Action of caffeine on the brain component of motor synaptic potentials, Gualtierotti (1).—Action of caffeine injections on the spinal reflexes of frog, Gualtierotti (2).— Effect of thyroxine on Triturus limb regeneration, Hay (1).-Metabolic energy for chloride transport in the gastric mucosa of Rana pipiens, Heinz & Durbin.

—Effect on oxygen consumption by isolated frog skin of temperature, Huf & Doss.—Effects of metabolic inhibitors and drugs on NaCl transport and O<sub>2</sub> consumption in the skin, Huf, Doss & Wills.—Suppression of growth in limb areas of Xenopus larvae by treatment with Nitrogen Mustard, Hughes & Tschumi.— Effect of azide and chloretone on sodium and potassium contents of frog sciatics, Hurlbut.-Enzymetic hydrolysis of propionylcholine, acetylthiocholine and butyrythiocholine by extracts of frog rectus abdominis, Jacob & Pécot-Dechavassine.—Prepotential, resting potential and latent period relations in frog muscles, Jenerick.—Stress-strain and force-velocity experiments on striated muscle, Jewell & Wilkie.— Effect of synthetic lysine-vasopressin on adenohypophysial activity in Bufo bufo, Jergensen & Nielsen.—Micro-electrode study of cerebellar electrical activity, Joynt.—Effect of alkaline and alkaline-earth solutions on the pulmonary circulation of the frog, Jullien, Cardot & Moulin.—Ligaturing the pulmocutaneous trunks and branches and the survival of the frog and possibility of recommencing the circulation, Julien, Ripplinger, Cardot, Baylé & Tisserand.—Toxicity of various chemical solution on larvae Triton alpestrishearts, Julien, Ripplinger, Cardot & Marcot.—Atropine vasodilation in experiments on frog pulmonary circulation, Jullien, Ripplinger, Cardot & Moulin.— Effect on circulation of introducing alkaline solutions to the pulmonary cavity, Julien, Ripplinger, Cardot, Moulin & Thomas.—Various tissue explants to determine blastopore lip potencies in Triturus gastrulae, Kato (3).—Application of the indirect peroxidase reaction to tissue staining, Kawasaki.-Uptake and turnover of injected I181 in tadpoles of Rana clamitans, Kaye & Le Bourhis.—Combined effect of methyl testosterone and ovarectomy on crest production in female newts, **Kiortsis** (1).—Enzyme inhibitors on resting potential and ion distribution of the sartorius muscle of Rana pipiens, Kloot.—Effect of desoxy-corticosterone acetate on thyroxine induced metamorphosis of Anura, Kobayashi.—Stimulation of spinal ganglion cells in sodium-free media, Koketsu, Cerf & Nishi.—Influence of calcium chloride on muscle of frog, Krolenko (1).—Afferent fatigue and habituation in Bufo bufo due to constant skin stimulations, Kuczka.-Changes in the neurosecretory and neuroglial cells of fresh frog brains when ultracentrifuged, Kumamoto.—Active sodium transport by the fuged, Kumamoto.—Active sodium transport by the isolated toad bladder, Leaf, Anderson & Page.—Disappearance from the skin glands of 5-hydroxy-tryptamine after injections of reserpine, Liébecq-Hutter & Bacq.—Effect of HCG on explusion of spermatozo in 3 Bujo viridis, Lunenfeld, Rabau, H-Steinberg & Szejnberg.—Action of aldosterones on the active transport of sodium in frog skin, Maetz, Marsh. Lunenfeld, exploration of explanation of explanations of explanatio Jard & Morel.-Implantation of embryonic and tadpole tissues into adult frog limbs, Malinin & Deck .- Metabolic effect of cortisone acetate during

limb regeneration in Triturus v. viridescens, Manner .-Effect of lesions of the middle eminence on Triturus cristatus carnifex, Mazzi.—Drug sensitivity of the isolated toad heart during hibernation, Nayler.— Lability and accommodation in frog nerve under various chemical stimuli, Nikolsky.-Prolactin and somatotrophin on forelimb regeneration in Triturus alpestris, Niweliński.—Modifications in skeletal muscle activity of salamanders caused by convulsant and anticonvulsant drugs, **Peters et al.**—Total and partial sympathectomy in *Rana pipiens*, **Pick.**—Thymus extract effect on gastrocnemius of female frogs, **Pora** & Toma. - Induction of ovulation in frogs using mammalian hormones, Ramaswami & Lakshman.-Experiments on ATP inhibition of ovulation in Rana pipiens, Rondell & Wright.—Action of tetraiododesaminothyronine on metamorphosis of Rana temporaria tadpoles, Roth. (1).—Experiments on haploid amphibians, Ruhland.—Variations observed in the action blans, Kuniand,—Variations observed in the action of adrenaline on the deposition of glycogen in frog and pigeon muscles, Sarzana & Castro,—Effect of colcemid and TEM on the differentiation of the lens of embryonic Triton alpestris, Schenk & Töndury. — General observations on thyroideotomized newts, Schmidt A. J. — Comparison of effects caused by phenol and other spindle poisons on segmentation in amphibian eggs, Sentein.—Effect on chemical relationships within muscle of altering the external NaCl and KCl levels, Simon, Shaw, Bennett & Muller.—Action potentials of frog nerve and stimulation frequency affect, Sjodin & Mullins.—Metabolic studies on isolated epithelium of frog skin, Skjelkvale, Nieder & Hut.—Relation be-tween liver weight and degree of restoration after partial hepatectomy in Rana pipiens, Streett.— Colchicine on early frog development, Su & Tsai.— Effect of boric acid on the development of eggs of Bufo vulgaris formosus, Takeuchi, T.—Antagonism in the action of calcium and strontium on the frog heart, Thomas.-Effect of extracellular potassium on intracellular action current in the heart of the intact toad, Trethewie & St. Clair.-Electromotive force setup by Rana temporaria tadpoles in opposition to an electrical stimulus, Viaud & Dreyfus.-Effects of cytotoxic substances on amphibian embryos, Waddington.—Effect of Pitressin on water exchange and lymph drainage in the frog, Wilson & Archdeacon.— Influence of steroids on sex differentiation in Pseudacris, Witschi, Foote & Chang .- The effect of sulfhydryl reagents on oxygen treated embryos of Rana pipiens, Wittner & Rosenbaum,—Effect of various concentra-tions of acetone on the acetylcholine contracture of the rectus abdominis muscle, Zelený & Kozák.

Hormones.—Gonadotrophines and sperm elimination in Bufo arenarum, Allende & Astrada.—Sensitivity of Bufo bufo to chorionic gonadotrophin, Allison.—Responses of Hynobius larval gonads to estrogens, Asayama & Miyamori.—Effect of chromatotrophic hormone on anuran pigmentation, Bagnara & Neidelman.—Diuretic hormones and their action on Bufo arenarum, Becerra.—Supporting role of cortisone and ACTH in hormonal control of ovulation, Chang & Witschi.—Comparative effects of goitrogens on thyroid activity in Triturus and Desmognathus, Dent & Lynn.—Effect of follicular hormone on sex differentiation in Hynobius retardatus, Hanaoka (1).—Effect of testosterone-propionate on sex-differentiation in Hynobius retardatus Hanaoka (2).—Influence of gonadial hormones on the differentiation of the gonad in Rhacophorus schlegelii, Iwasawa (1).—Relation of

]

the

der

and

rus

ind

aus

ing

wia

hi-

ion

rog

of

the

ön-

zed

ects

on

feet

ter-

on

rog

& of be-

fter

t.—

of

ism

rog

on

tact set-

an

of

and

a.—

dryl

ens,

tra-

e of

ina-

siti-

son.

ens.

phic

on

one

g &

on

ent

ffect

in

of

nad

n of

melanophore-expanding hormone to hyperglycaemia in Xenopus laevis, Munday.—Action of sex hormones on the motivity of the gut of Rana during the year, Pora & Rejep.—Role of mammalian hormones in inducing ovulation in frogs, Ramaswami & Lakshman.—Induced ovulation by hormonal treatment in Rana temporaria and R. pipiens, Taurand.—Concentrations of thyroxene and iodine in frog ovaries, Volpent, Michel & Roche.

Laboratory Technique.—Staining technique with thionin blue and Ag. impregnation of Rana dendrites, Cerl & Chacko.—Methods of applying electrical stimulation to lateralis organs of Xenopus laevis, Murray.

Locomotion: Movement.—Coordination of swimming movements in Ambystoma mexicanum after central nervous system regeneration, Romanovsky.—Galvanotropism of Rana temporaria tadpoles, Viaud & Dreyfus.

Muscular activity .- Myocardium of Rana temporaria in the absence of potassium and under the influence of acetylcholine, Borghgraef & Carmeliet.—Distribution of inorganic phosphate in Bufo marinus muscle, Briner, Simon & Shaw.—Longitudinal field, coupled with depolarization in activating frog twitch muscles, Csapo & Suznki.—Chlorpromazine blockage of water imbibition by frog gastrocnemius, Eckhardt & Govier. The effect of trematode infection on the physiological activity of the sartorius, Frank.—Frog muscle metabolism and structure of sulphonylurea compounds, Gourley.—Experiments upon the transmission of stimuli in cardiac musculature of Triturus cristatus, Gyévai.—Rehydration contraction of frozen dried muscle of Rana pipiens, Hopkins.-Neuromuscular facilitation by sympathetic stimulation of frog muscle, Hutter & Loewenstein. - Eserine-acetylcholine interaction in frog rectus abdominis, Jenden,-Latency dependent prepotential of depolarising fibres in sartorius muscle, Jenerick,—Dependence of the heart ectivity of the frog on a substance of nervous origin, Jullien, Ripplinger & Cardot.—Enzyme inhibitors on resting potential and ion distribution of sartorius muscle of frog, Kloot.—Contractions of muscle fibre in frog, Krolenko (2).—Diastole and systole in the ventricle muscles of the frog heart, Lacquet.— Electrical activity of salamander skeletal muscle and modifications by anti-convulsant and convulsant drugs, Peters et al.—Mechanical action of the frog gastrocnemius, Pora & Oros.—Effect of thymus extract (CIF) on the gastrocnemius of female frogs, Pora & Toma.—Relationship between sodium, potassium and chloride in amphibian muscle, Simon, Shaw, Bennett & Muller.—Effect of temperature on frog sartorius neuromuscular junction, Takeuchi, N.—Contraction and coagulum formation of isolated frog muscle fibre, Umezawa.-Effect of acetone on the tonie muscle of Rana temporaria, Zelený & Kozák.

Nervous System.—Observations on spinal ganglion cells of Rana pipiens in normal animals and after injection of malononitrile, Anderson & Breemen.—Effects of chemicals on ionic movements during nerve activity in Rana pipiens, Asano & Hurlbut.—Frog nervous physiology, Burkhardt.—Retrograde reaction in motoneuron dendrites, Cerf & Chacko.—Localisation of neural and lens antigens in frog embryos in relation to induction, Flickinger.—Neomycin and its effects on nerve blocking in frogs, Fulchiero, Turcotte & Martin.—Action potentials in dorsal and ventral

roots of the spinal cord, Gualtierotti (1).-Changes in action potentials in sciatic nerves after injections of caffeine, Gualtierotti (2),-Effect of drugs on sodium and potassium contents and respiration of frog sciatic, Huribut.—Sympathetic innervation of frog leg muscle, Hutter & Loewenstein.—The Schmidt-Lautermann incisure of the peripheral medullated nerve fibre, Ishida.—Cerebellar electrical activity in Rana esculenta, Joynt.—The role played by the spinal cord in the regeneration of the tail of larval salamanders, Kiortsis & Droin.—Electrical responses of Rana pipiens spinal ganglion cells in sodium-free media, Koketsu, Cerf & Nishi.—Effect of the augmented nerve supply on regeneration in postmetamorphic Xenopus laevis, Konieczna, Marczynska & Skowron-Cendrzak.—Physiology of the nervous system in frog and rôle of nucleic acids, Koshtoyants.—Short term afferent fatigue, habituation, reflex actions and instinctive activities and their role in learning behaviour of Bufo bufo, Kuczka.-Distribution of sulphydrile substances in the hypothalamus and neurohypophysis of the toad, Lasansky & Sabatini .-Secondary cerebral modifications of the paraphysis of Rana esculenta, Legait & Legait,—Depolarization in frog's heart during "KCl contractions," Lüttgau & Niedergerke.—Bufo bufo efferent fibres in the optic nerve and their behaviour after section of the nerve. Maturana.—Change of constants a and b in Hoorweg-Weiss formula at the action of substances on frog nerve, Nikolsky .- Histochemical study of cholinesterase at neuromuscular junctions of the abdominal musculature of the frog, Pecot-Dechavassine.—Retinal sensitivity in Rana esculenta, Pilz, Sickel & Birke. Action of the nervous system on the mechanical action of the frog gastrocnemius, Pora & Oros .-Central nervous system regeneration in embryos of Ambystoma mexicanum, Romanovsky.—Quantitative analysis for RNA and protein nitrogen before and after induction of the neural tube, Rounds & Flickinger .-Neural closure in amphibia and the forces concerned, Selman (1).—Action potentials of frog nerves, Sjodin & Mullins.—Threshold dependence of action potentials in medullated nerve fibres, Stieve .--Conduction velocity in intracutaneous nerve fibres of frog, Takagi. -Depression in frog neuro-muscular transmission, Takeuchi, A.—Effect of temperature on frog neuromuscular junction, Takeuchi, N. — Extra cellular potassium effect on intracellular action current of intact toad heart, Trethewie & St. Clair.—Attempts to determine origins of sympathetic ganglia, neurilemma sheath cells, and spinal meninges, Triplett.-The hypothalamic neurosecretory system of Hyla regilla, in normal animals and after transection of the infundibular stalk, Wilson, Weinberg & Bern.—Temporal reflexes associated with spinal reflexes in frogs, Wittenberger & Fabian.—Spina bifida developed at the neural stage in Bufo vulgaris formosus by treatment with potassium oxalate, Yamamoto.

Osmotic relations: Water balance.—Dehydration and hydration and ability to withstand desiccation in frogs of the genera Neobatrachus and Heleioporus, Bentley, P. J. et al.—Effect of vasopressin on water uptake in Bufo in different hypotonic solutions, Bentley (1).—Effect of pituitrin on water transfer across the wall of the isolated urinary bladder of Bufo, Bentley (2).—Water flow through the gastric mucosace of Rana pipiens and R. clamitans, Durbin, Frank & Solomon.—Electromotive chloride transport in the frog, Durbin & Heinz,—Effect of trematode infection on frog's water balance mechanism. Frank.—Action of

 $T_{\mathcal{T}}$ 

nei

Ro

im

Fr

of

po

eff

to

wi

Re

of

mi

tic

T

pla

tic

irı

& de

E

ge in Bi ai & A th

in G se lo et S. hi

st all lo R h lo S A

pitressine and adrenalin on water balance of Bufo paracnemis during and after metamorphosis, Hadler, Berezin & Lison.—Destruction of osmo-regulatory functions in organs of Diemictylus viridescens after X irradiation, Jakowska et al.—Tolerance of water loss by salamanders, Ray.—Poetures of frogs during water absorption, Stille.—Effects of fresh water and marine teleost pituitary on water balance in Rana pipiens, Weisel.—Fluid drainage and the effect of pitressin from the frog dorsal lymph sac, Wilson & Archdeacon.

Pregnancy test use.—Melanophore dispersing substance in human pregnancy urine, Burgers & Mighorst.

Respiration.—Influence of various sound frequencies on respiratory movements, galvanic skin reflex and gastric hunger movements in Rana, Kleerekoper, & Sibabin.—Blood oxygen capacity in frogs, Lettwich.—Respiratory quotient and aerobic glycolysis of Rana esculenta skin under varying temperatures, Locker, A.—Cytochrome-c-oxidase and respiratory rate in larval Bufo bufo, Petrucci.—Regulation of cutaneous respiration in frogs, Poczopko, P.—Increase in respiratory activity during the early development of Siredon mexicanum, Rudnick.—Method for stabilizing Ogonsumption values in frogs, Springmuller & Concha.

Secretion: external.—Investigations on mucilage secretion, Schulz & Paola.—Changes in epidermal glands after removal of the pituitary of clawed toads, Spannhof.

Secretion: internal.—Total catheptic activity of embryos of *Xenopus laevis* and its role in development, Deuchar.—Effect of ultracentrifugation on the hypothalmic neurosecretory system of fresh frog brains, Kumamoto.—Thyroid activity in relation to cytodifferentiation of anterior pituitary of *Xenopus*, Saxén.—The hypothalamic neurosecretory system of *Hyla regilla*, Wilson, Weinberg & Bern.

Senses.—Rana temporaria passive electrical properties of retina, choroid and sclera for radial fields and currents, Brindley (1).—Effect on frog's electroretinogram of varying the amount of retina illuminated, Brindley (2).—Retinal responses to light in sexised eye of Rana temporaria, Brindley (3).—Reaction to light in frog, Brodsky & Nechaeva.—Visual pigments of Amphibia, Crescitelli.—Visual mechanism and sensitivity in Xenopus laevis, Dartnall.—Average light absorption and density photo sensitivity of retinal receptors, Denton & Wyllie.—The representation of the retina on the optic lobe of frog, Gaze.—Reactions caused by stimulation of the horizontal semi-circular canal in Rana viridis, Gribenski.—Yawning behaviour of Bufo calamita in relation to tracking prey by sense of smell, Heusser (3).—Capacity of the lens forming tissue in amphibia, Kankave.—Wiping movements, fatigue and habituation after skin stimulation in Bufo bufo, Kuczka.—Response of lateralis organs of Xenopus laevis to electrical stimulation by direct current, Murray.—Sensitivity of the retina studied by means of an electro retinograph, Pilz, Sickel & Birke.—Acetyl choline hydrolizing enzyme in the receptor cell of frog retina, Yoshida (1).

Transplants, Implants, Grafts: Regeneration Studies.

Grafting experiments on brain size and eyes in Anurs, Balinsky.—Parabiosis experiments with metamorphosing and post metamorphosing newts, Beetschen (1).—Regeneration from supernumary blastemas in adult Triturus viridescens, Bodemer.—Transplanta-

tion experiments with Triton cristatus and Rana esculenta, Bondi.—Excision of the presumptive prosencephalic area of the neural plate of Triturus alpestris, Boternbrood.—Ultraviolet irradiation of upper arm of Ambystoma opacum and its effect on regeneration of the lower part, Bulter, Blum & Schmidt.—Experiment to demonstrate that initially all regions of the primary ureter blastema possess the same morphogenetic potentialities, Cambar & Gipouloux.-Inhibition of regeneration, morphogenesis of regenerates and regenerates on reversed hind limbs, Deck & Riley.—Muscular regeneration in newts under the influence of vitamins, Degan & Asandei.— Bilateral removal of trunk neural folds, transplanting from normal and abnormal hosts, Finnegan.—Parabiosis experiments on Pleurodeles waltlii, Gallien (1).-Skeletal regeneration in amphibians, Goss .- Lower jaw regeneration of newts after excision of the intermandibular regions, Goss & Stagg (1).—Regeneration of lower jaws in adult newts, Goss & Stagg (2).— Sexually mature individuals of Xenopus laevis from the transplantation of single somatic nuclei, Gurdon, Elsdale & Fischberg.—Limb regeneration in *Triturus* viridescens, Hay (1).—Fine structure of blastema cells and differentiating cartilage cells in regenerating limbs of Ambystoma punctatum larvae, Hay (2).—Grafting in parabiosis larvae of Amblystoma, Hughes & Tschumi.—Pronephric regeneration in thiourea treated tadpoles, Hurley.—Explants from the dorsal marginal zone of amphibian eggs, Ikushima.—Wound healing in an intact amphibian and in an isolated organ, Ioff.—Regeneration of the endbrain in postmetamorphic Xenopus laevis, Jordan.-Implantation of the nasal placode in the dorsal area of the forelimb organ district of urodele embryos, Kiortsis (2) .- The role of the spinal cord in the regeneration of the tail of larval Salamandra salamandra, Kiortsis & Droin.-Capacity of heteromorphic limb regeneration in postmetamorphic Xenopus laevis, Konieczna-Marczynska & Skowron-Cendrzak,-Olfactory nerves and endbrain regeneration in Xenopus laevis, Kosciuszko. Orthotopic transplantation of presumptive lateral mesoderm of early Triturus gastrulae, Kotani.— Multiple regeneration of tadpole tails of Xenopus laevis, Kucias.—Limb regeneration experiments in Alytes obstetricans, Lecamp.—Implants of methylcholantrene crystals under the skin of Triturus cristatus, Leone.—Transplantation and trunk muscle development in Anura, Liedke.—Regeneration after amputation Rana pipiens and R. clamitans, Malinin & Deck .- Metabolic effect of cortisone acetate during limb regeneration in Triturus v. viridescens, Manner. Regeneration after amputation of abnormal structures. Newth (1).—Regeneration after the amputation of abnormal structures, Newth (2) .- Prolactin and somatotrophin on forelimb regeneration in *Triturus* alpestris, Niewelinski.—Asportation and successive orthotopic limb grafts in Bufo vulgarie, Notarmicola,-Regeneration of reversed polarity limb extremities in salamander larvae, Oberheim & Luther.—Regeneration of the intestine in adult urodeles, O'Steen .-Transplantation of somatic nuclei to normally fertilised and non-enucleated eggs of Triturus alpestris, Pantelouris & Jacob.—Limb grafts in relation to the development of the spinal cord and ganglia in Anura, Perri (1).—Embryonic limb bud transplantations of Bufo viridis to B. vulgaris, Perri (2).—Regeneration in the spinal cord of the adult frog, Piatt & Piatt.— Regeneration experiments with new tails, Preda & Persecă.—In amphibia, Raitzina.—Homoplastic and heteroplastic grafts between Amblystoma donors and na

ve

us

of

on

&

lly

u-

of

bs,

vts

ng ra-

7er

er-

on

om

on,

248

alls

ng

nes

sal

nd

ed st-

on

nb

he

of

in

ar-

nd

ko. ral

nus

in

yl-

rus

cle

ter

&

ng

es,

of

nd

rus

ive

in

ra-

ti-

·is,

he

ra, of

ion

nd

nd

Triturus hosts, Reyer.—Degree of regeneration of the nervous system of pre neurula embryos of the axolotl, Romanovsky.—Larval incompatability in grafting Anura and Urodela, Rossi.—Nerve dependency in implant-induced supernumary formation, Ruben & Frothingham.—Regeneration in athyroid newts, Schmidt, A. J.—Adrenal transplants and forelimb regeneration in frogs, Schotté & Wilber.-Transplants of neural fold material from three different zones in Triturus alpestris, Seno & Nieuwkoop.—Regeneration of amphibian body parts, Singer.—Regenerative potentialities of limb and tail cells of axolotls, Skowron & Roguski.-Implantation of pituitary tissue and its effect on the epidermal glands of hypophysectomised toads, **Spannhof**.—Preliminary grafting experiments with Triton and Rana esculenta, Spirito & Bondi. Radio resistance in regenerating and developing limbs of urodele larvae, Stinson,-Transplantation experiments with skin fibromas of tumour bearing Triturus taeniatus, Stolk (3).-Retina influence on lens regeneration, Stone.-Lens regeneration and retinal factor in Triturus viridescens, Stone & Gallagher.—Transplantability of haploid nuclei, Subtelny.—Regeneration of inhibited limb stumps after release from irradiation, Thornton.—Weakening of regenerative ability of tissues by cancer causing substances, Tokin & Filatova.—Grafts of embryonic nervous material to determine origins of sympathetic ganglia, Triplett .-Effect of antiserum on inductive capacity of Triturus vulgaris, Vainio.—Ear vesicle grafted in various abnormal regions of Amblystoma, Van Dyke & Detwiler.

Venoms.—Venoms secreted by parotid glands in Bufo spp., Frèrejacque.

X-rays etc. Ionizing irradiation (X-rays and gamma rays) of amphibian tissues, Allen.-Effects of irradiation on parts of Siredon mexicanum heads, Brunst.-Comparison of the effect of ultra violet light and X-rays on the capillaries of the frog's web, Bücker & Hanke.—Ultraviolet irradiation of upper arm of Ambystoma opacum and its effect on regeneration of the lower part, Butler, Blum & Schmidt.—Diminution in dipeptidase activity by X-rays, Coromaldi (1).— Geiger counter assay of radioactivity of K-tracer in sartorii of Rana, Harris.—Hematological and histological effects of whole body X-irradiation, Jakowska et al.—Amputation and despinalisation of the tail of S. salamandra and the reincorporation of irradiated hind limbs, Kiortsis & Droin.—Autoradiographic study on iodine accumulation in pigmented tissues of amphibian larvae, Koshida (1).—Autoradiography of the skin of Rhacophorus larvae, Koshida (2).—Teratological effect of ultra-violet rays on larval amphibia, Rostand (2).—Ultra-violet light method of producing haploid amphibian embryos, Selman (2),—Histological cells changes induced by X-rays, in tadpoles, Speidel .- X-irradiation of larval urodeles, Stinson. Autoradiography to show differences in uptake of a tracer <sup>14</sup>CO<sub>2</sub> in early Axolotl gastrulae, Tencer.— Effects of ultra-violet light irradiation on limb regeneration of larval urodeles, Thornton.

#### REPRODUCTION AND SEX

Experimental. — Modification of Hynobius sex development by methyl-testosterone, Asayama & Amanuma,—Responses of larval gonads to estrogens, Asayama & Myiamori.—Inhibition of sperm production by saline and hypophysial solution in the toad, Astrada & Taleisnik.—Maintenance of frog larvae

gonads on synthetic media, Foote & Foote (2).—Abnormal gonadial differentiation due to hormonal injections in Rhacophorus tadpoles, Iwasawa (1).—Histology of Bidder's organ in Bufo arenarum after castration and after the action of sex hormones, Penhos & Cardeza.—Extra gonadial influence on the gonad of Triturus alpestris, Stärk (2).

Hybridisation.—Discussion of isolating mechanisms between Bufo terrestris-voodhousei group and B. valliceps and within the B. terrestris-woodhousei complex, Gosmer & Black (2).—Distinct hybrids produced, with special regard to cell and nuclear detail, among three European toads, Hertweig, Weiss & Ziemann.—Transference of haploid nuclei between Rana pipiens and R. sylvatica, Moore (3).—Interspecific gene exchange between B. fouleri and B. terrestrie in S.E. Louisiana, Volpe.—Scaphiopus h. hurteri × S. h. holbrooki, skull figured p. 314, Wasserman.

Intersex.—In adult Pelobates punctatus after treating larvae with oestradiol, Gallien & Collenot.

Reproduction.—Response by Q Hyla cinerea to call from G, Blair (2).—Studies on the grassfrog, Bechtle.—Of Bufor. regularie; study of oviposition, Chapman & Chapman.—Reproductive behaviour and breeding pattern of Scaphiopus holbrooki, Hansen.—Of Rhacophorus c. cruciger, Morgan-Davies (1).

Secondary Sex Characters.—The vocal sac of Bufo alvarius, Inger (1).—Thumb-pads in normal and hypophysectomized frogs after gonadotrophin injections, Burgos & Ladman.

Sex Determination.—Modifications in the sexual characters of male Bufo arenarum after castration, Penhos & Cardeza.

Sex Differentiation.—In Rana esculenta and R. dalmatina, Ancona.—Modification of sex differentiation by steroids in Pseudacris, Witschi, Foote & Chang. — Effect of follicular hormone on sex differentiation in Hynobius retardatus, Hanaoka (1).—Effect of testosterone-propionote on sex differentiation in Hynobius retardatus, Hanaoka (2).

Sex Dimorphism.—Dimorphic maxillary dentition in Pseudacris, Goin, C. J. (2).—In Rana albolabris, Guibé & Lamótte (4).

Sex Maturation.—In vitro cultivation of larval anuran gonads, Foote & Foote (3).—Single somatic nuclei of Xenopus laevis producing sexually mature individuals, Gurdon, Elsdale & Fischberg.

Sex Reversal. — Feminization of the gonad differentiation in male salamanders, Asayama & Matsuzaki.—Sex reversal in male Pelobates cultripes with costradiol, Gallien & Collenot.—Sex reversal in thiourea-treated frog tadpoles, Iwasawa (2).

## DEVELOPMENT

Egg: Oogenesis.—Inducted oviposition and incubation of eggs of Batrachoseps, Anderson.—Water absorption by Hynobius eggs, Aoki.—Control of development and differentiation of cells in Rana pipiens gastrulae, Barth & Barth.—Anuran germ cells studied in the blastocoel floor, Blackler.—Eggs of Plethodon g. glutinosus, Brode & Gunter,—Basic requirements for egg laying in salamanders, Carl.—Cortisone effect on frog ovulation, Chang & Witschi.

gain (2) S.-Str

men

st

R

of

of in R K by

a Min o Ba o in

a

e

Streptomycin activity on eggs of Bufo vulgaris, Chiuini.—Triturus alpestris cortical permeability gradient, Dollander.—Chemical analysis of nucleic acid content of nuclei in Rana ova, England & Mayer.— Artificial insemination of newt eggs, Fischberg.— Glycogen content of Rana pipiens eggs at different stages of development, Fitch & Merrick.—Biological observations on Leptodactylus prognathus, Gallardo (2).—Development and description of eggs of Pseudacris brimleyi Gosner & Black (1).—Direct development of Arthroleptis crusculum, Guibé & Lamotte (2).

Fredrich of particular of production of the prognation of the production of the —Evolution of nucleoli in batrachian ovcoytes, Guyénot.—Experimental doubling of triploid nuclei in toad hybrids removed from the ovary, Hertwig, Ruhland & Weiss .- Kinetic and histogenetic pheno-Runand & Weiss.—Americ and instogenetic plenomena in the marginal zone of amphibian eggs, Ikushima.—Crystalline lattice structure of yolk platelets in embryonic Triturus pyrrhogoster, Karasaki & Komoda.—Blastopore potencies of Triturus gastrulae, Kato (3).—Formation of primordial germ cells from Triturus presumptive ectoderm, Kotani.—Electron microscope study of lampbrush chromosomes from amphibian occytes, Lafontains & Ris.—Hypothetical assumption of dorso-ventral polarity giving higher oxidative rates at certain parts of amphibian eggs, LEVITUP.—Effect of oxygen poisoning on gastrulation in Rana pipiens eggs. Malamed (2).—Eggs and larvae of Eurycea lucifuga described, Myers (1).—Lipid bodies in the ocytee of Rana tigrina, Nath, Gupta & Kaur.—Cortical changes at fertilization of toad eggs.

Osanai.—Protein synthesis in cocytes of Triturus palmatus and T. alpestris, Pantelouris.-Influence of pH on ovulatory capacity, inhibitors during incuba-tion and the ovulatory period, and of smooth muscle activity, Rondell & Wright.—Nature and distribution of polysaccharide protein complexes in the cortex of developing Rana pipiens occytes, Rosenbaum.— Influence of factors on haploid or diploid egg production, Ruhland.—Changes in the achromatic apparatus, nucleus and chromosomes of Triturus and Pleurodeles after treatment with phenol, Sentein.—Cleavage in the newt egg, Swann & Mitchison.—Effects of boric acid on egg development in Bufo vulgaris formosus, Takenchi, T.—Heterogeneity of nucleoli in occytes of Bufo arenarum, Tandler.—Ovulation in Rana pipiens and R. temporaria, Taurand.—Concentrations of thyroxene and iodine in the overy of Rana temporaria, Volpert, Michel & Roche.—Orientated section of Rana gastrula, Wessing & Class.—Artificial culture mediums for Alytes obstetricans eggs, Wettstein.— Resistance and susceptibility to high oxygen pressures in the early stages of Rana pipiens development, Wittner & Rosenbaum.

Sperm: Spermatogenesis.—Cytological events in the spermatogenic cycle and the deposition of glycogen in sertoli cells, Burgos & Ladman.—Comparative nucleic acid content of spermatozoa of the frog, England & Mayer.—Discharge of sperm after injections of HCG, Lunenfeld, Rabau, H-Steinberg & Szejnberg.—Action of gonadotrophines on sperm production in Bufo arenarum males, Moisset de Espanés.—Regulation by temperature of the spermatogenetic cycle in Rana temporaria, Oordt.—Spermatophore of Plethodon jordani metcalfi. Organ.

Effects of Environment.—Dependence of spawning on external factors in Salientia, Rühmekorf (1).—Cold temperature and delayed metamorphosis produces accessory limbs in frogs developing in natural conditions, Voitkevich (2).

Electrophoresis.—Changes of the Electrophoretic serum pattern of *Ambystoma* according to age, sex and metamorphosis, Manelli (2).

Embryology. — Amphibian morphogenesis, in particular Rana pipiens, and the effect of phosphorylated nucleosides, Ambellan.—Ammonia -, urea-, and iron-contents in the perivitelline fluid of Rhacophorus schlegelii var arborea, Arima.—Comparative embryology of vertebrates, Baer.—Thermal stadia in the development of Rana temporaria, Bebak.-Correlation between lenticular and total development in embryonic Triturus, Becker.—Cytological study of heteroploid embryos of Pleurodeles walthi, Beetschen (2),-Of Bufo terrestris charlesmithi, B. t. terrestris, (a).—Of Buyo terresirs chartesimin, B. t. terresirs, B. t. americanus, Bragg (2).—Development of Eupsophus taeniatus, Capurro, (2).—Development of Chiromanis petersi kelleri, Cherchi (2).—Activity of streptomycin on the development and differentiation of eggs and larvae of Bufo vulgaris, Chiuini.—Action of the thermal gradient on embryonic development of Triturus alpestris, Cury & Dollander. — Spleen development and experimental modification in Taricha torosa, DeLanney.—Discussion of possible assistance in yolk breakdown derived from catheptic activity in Xenopus laevis, Deuchar.—Development of collagen in Rana pipiens embryos, Edds.—Behaviour of Rohon-Beard cells during embryonic and larval development, Filippio.—Embryonic induction in Rana pipiens and antigen localisation, Flickinger.—Relation of natural and imposed electrical potentials and respiratory gradients to morphogenesis, Flickinger & Blount.—Influence of temperature on the development of Xenopus laevis, Gerbes.—Introduction of a barrier between dorsal and ventral regions of Xenopus embryos and limb field effects, Hughes & Tschumi. Oxygen consumption of Ambystoma maculatum embryos developing in an envelope inhabited by Oophila amblystomatis, numerous protozoa, and bacteria, Hutchison & Hammen,—Differentiation of the cortical cytoplasm and inclusions in occytes of frog, Kemp.—Development of a dorsal crest by female newts under varying conditions, Kiortsis (1).-Development of the ovary in virgin female Nectophynoides, Lamotte & Rey.—Utilization of lipids by embryonic Bufo vulgaris, Lernia, E. D.—Anuran trunk muscle development, Liedke.—Physiological interpretation of the mechanism of determination of bilaterality in Amphibian embryos, Levtrup.— Inversion of the dorso-ventral axis in Siredon mexica-num, Levtrup & Pigon.—Qualitative changes at gastrulation, initiation of a new development dependent chemical system, Malamed (2).—Origin and development of the pronephros and Wolffian duct in Rana, Manelli (1).—Notes on Ramanella obscura, Morgan-Davies (2).—Energy-rich phosphate content of Bufo vulgaris formosus embryos temperature alterations, Moriwaki.—Increased partial pressure of O, accelerates development in Rana embryos, Nelsen. Factors influencing the control of limb development in Amblystoma punctatum, Nicholas.—Developmental capacities of the cell nuclei of various tissues of the newt, Pantelouris & Jacob.—Intrinsic and extrinsic factors in the development of the spinal cord and ganglia in Anura, Perri (1).-Lens induction from prospective belly ectoderm of Ambystoma punctatum and Triturus viridescens, Reyer.—Embryonic nutri-tion, Rudnick.—Forces producing neural closure in amphibia, Selman, (1).—Autonomous and dependent differentiation of the neural crest in Triturus

97

etic

and

in

ory-

and

rus

the

tion em-

of

hen

tris,

of t of

ent y of

tion

n of

een

ible

ptie

t of

our

in

ials

opof a

pus i.—

tum

by

of

ale

cto-

by

ran

ical

of

ica-

at enand in

ıra,

ent

of

en.

ent

tal the

sic

ha

om

tri-

in ent

·us

alpestris, Seno & Nieuwkoop.—Initiation of amphibian gastrulation experiments, Shapiro.—Double nucleus in early development of Bufo bufo, Sirakami (1) & (2).—Induction of triploidy in Xenopus laevis, Smith, S.—Development and variation of gonad and sex cells, Stark (1).—Larval growth and development of experimental haploids and homozygous diploids, Subteiny.—Study of sympathetic ganglia, spinal meninges and neurilemma sheath cells to determine embryological origin, Triplett.—Early developmental studies on Rana fusca, Veit.—Amphibian embryos and cytotoxic substances, Waddington.—Induction fields in early embryonic stages of Triton, Woellwarth.—Induction of double embryos in Bufo vulgaria formosus, Yamamoto.—Histochemical study of salamander retine in late embryogenesis, Yoshida (2).

Embryology: Chemical. — Chemical nature of Rhacophorus perivitelline fluid, Arima. — Bylucuronidase activity during early development of Xenopus laevis embryos, Billett (1). — Study of ribonucleic acid in embryonic Bufo vulgaris, Coromaldi (2). — Lipid utilization by embryonic Bufo vulgaris, Lernia. — Streptomycin and chloromycetin inhibitors of Rana metamorphosis, Marco. — Cytochemical investigation of the cortical cytoplasm of developing Rana pipiens occytes, Rosenbaum. — RNA and protein in neural induction in Rana pipiens, Rounds & Flickinger. — Role of yolk proteins in amphibian embryos, Sampath Kumar et al. — Blocking of mitosis in early metaphase by Colcemid, and lens differentiation in larval Triton alpestris, Schenk & Töndury. — Effect of pH, and salinity on the response to blastocoel perfusion in Ambystoma, Shapiro.— Regional incorporation of 14CO, into Amphibian gastrulae, Tencer.

Embryology: experimental—Inducted oviposition and incubation of eggs of Batrachoseps, Anderson.-Methyl testo-sterone modification of sex development in larval Hynobius, Asayama & Amamuna.—Effect of enzymes on fertilizability of Rana pipiens eggs, Barch & Shaver.—Effect of penicillin, streptomycin and penicillium mycelium on growth and development of Xenopus laevis, Ber & Zieleniewski.—Temporary inactivation of newt larvae by benzimidazole and its alkyl derivatives, Billett (2).—Inhibition of cell division in amphibian eggs by β.-mercaptoethanol, Brachet.-Experiments on the diffusibility of the amphibian evocator, Brahma.—Constriction of newt blastulae and its effect, Brice.—Effect of re-turning eggs on the embryonic development of Pleurodeles walthis, Brouant & Cury.—Effects on development of head structures caused by irradiation of young axolotls, Brunst.—Action of hormones on spermiation in Bufo arenarum, Burgos & Houssay.—Injections of cortisone, estradiol and ACTH to jumbo Q Rana pipiens and their effect on ovulation, Chang & Witschi. Streptomycin on differentiation of Bufo vulgaris, Chiuini .- Diminution of dipeptidase activity in young frogs by X-irradiation, Coromaldi (1).-Early removal of the pronephros and its effect on the development of the female genital opening in Rana dalmatina, Dechambe.—Induced ovulation in Necturus, Fitch, K. L.—Experimental induction of vertebrae in embryonic Pleurodeles waltlii, Gallien (2).—Odd results after hybridizing European toads, Hertwig, Weiss & Ziemann.—Embryonic and larval epiphysectomy in *Taricha torosa* and observations on scoliosis, Kelly.—Effect of desoxycorticosterone acetate on Anuran metamorphosis, Kobayashi.-Formation of primordial germ cells from presumptive

ectoderm of Triturus gastrulae, Kotani,—Gastrular blockage of frog's eggs produced by oxgygen, Malamed (1).—Induction of spermiation in Rana by adrenaline and pregnyl extracts, Oordt.—Young Bufo viridis larvae kept in dilute thyroxin solutions, Perri (2).—Haploid amphibian embryos produced by ultra-violet light, Selman (2).—Preliminary differentiation after grafting Triton to Rana esculenta embryos, Spirito & Bondi.—Differentiation of Rana larvae after treatment with various chemicals, Stefani.—Effect of colchicine on early development of Rana limnocharis, Su & Tsai.—Reaction of gastrula of Triturus vulgaris to implants of guinea pig treated with rabbit immune sera, Vainio.—Regional inductive effect of pentose nucleo-protein (PNP) from the liver and kidney, on Triturus gastrulae, Yamada.—Bufo vulgaris formosus eggs treated with potassium oxalate, Yamamoto.

Fertilisation.—Cortical changes at fertilization of toad eggs, Osanai.

Growth,-Size of brain and eyes in embryos of various Anura, Balinsky.—The mode of growth of the tail in urodele larvae, Bijtel.—Growth of Bufor. regularie studied by marking and recovery; development of fat bodies, ovaries and testes, Chapman & Chapman.—Growth rates of Rana catesbeiana, Cohen & Howard.—Size of species in relation to tooth size and count, Goin, C. J. (2).—Early stages in the normal development of the hind limb of Xenopus laevis, Hughes & Tschumi,-Rapid growth rates and better viability of embryo Ambystoma maculatum symbiotic with Oophila amblystomatis, Hutchison & Hammen.— Tissue growth and wound healing in Amphibia, Ioff.-Interrelation between growth rate, morphological changes of the hypophysis and thyroid and gonadal development, Iwasawa (2).—Growth inhibition in crowded tadpoles, Richards.—Control of tadpole growth by feed back mechanisms, Rose.-Effect of embryonic age on the response to blastocoel perfusion, Shapiro.—Seasonal variation in tooth size and number in Eurycea, Stewart.-The development and growth of haploid and homozygous diploid frog embryos obtained from transplantations of haploid nuclei, Subtelny.-Growth in the amphibian egg, Swann & Mitchison.

Larval Stages.—Development of Hyperolius zonatus and H. lamottei, Arnoult & Lamotte.—Reversed proximodistal polarity in larval and metamorphosing urodeles, Deck & Riley.—Origin, development and changes of the gills and lungs of Kaloula borealis, Giang & Li.—Of Pseudacris brimleyi, Gosner & Black (1).—Criteria for identification of some North American Bufo, Gosner & Black (2).—Larval development of Phrynobatrachus francisci, Lamotte & Dzieduszycka.—Larval development of Ramanella obscura, Morgan-Davies (2).—Larvae of Eurycea lucifuga, Myers (1).—Respiratory rate and enzyme systems in developing Bufo, Petrucci.—Somatic variation and dwarf phenomena in Telmatobius larvae from the Andes, Pisanó.—Inhibition by chemical substances of early embryonic differentiation, particularly of the lens, in Triton alpestris Schenk & Töndury.—Development of hard dentine in Triturus teeth, Schmidt, W. J. (1).—Of Rana palmipes and R. macroglossa, Volpe & Harvey.

Life History.—Biology and development of Eupsophus tacniatus, Cei & Capurro.—Of Pseudacris brimleyi, Gosner & Black (1).—Of Chioglossa lusitanica, Goux.—

ch

cer

fai Pe be de Te

A te C D D b G D A

Of Gastrotheca marsupiata in captivity, Grenard.— Direct development of Arthroleptis crusculum desed, and figd., Guibé & Lamotte (2).—Life cycle of Rana temporaria and the turnover of radiostrontium and radioyttrium by tadpoles, Lucas & Pickering.

Metamorphosis.—The phenomenon of autolysis in Bufo vulgaris, Baldisserotto & Ugurgieri,-Induced Bufo melanostictus metamorphosis, Bhattacharya & Medda.—Life history and metamorphosis of salamanders, Carl.—Rohon-Beard cells and resorption of the tail in Rana esculenta, Filippio.—Of Pseudacris brimleyi, Gosner & Black (1).—Hypophysectomized efts and failure to moult, Grant & Grant.—Influence of the removal of the endbrain on the metamorphosis of Xenopus laevis tadpoles, Jordan.-Effect of desoxycorticosterone acetate on metamorphosis in anuran tadpoles, Kobayashi. — Quantitative analysis of thyroid development during metamorphosis of Nectophrynoides occidentalis, Lamotte & Prum.— Histological changes and the effect on metamorphosis of varying the amounts of strontium-90 taken up by tadpoles, Lucas & Pickering,—Changes in the electrophoretic scrum pattern of Ambystoma, Manelli (2),—Inhibition of metamorphosis of Rana by Streptomycin Marco.-Thyroxine analogues effect on Rana pipiens metamorphosis, Money, Meltzer, Young & Rawson.— Of Rhacophorus c. cruciger desed., Morgan-Davies (1). -Delayed metamorphosis in Mixophyes and variations in size of metamorphosing larva of Limnodynastes dorsalis, Ormsby.—Respiratory rate in metamorphosing Bufo bufo, Petrucci.—Action of tetraiododesaminothyronine on metamorphosis of Rana temporaria tadpoles, Roth (1),—Action of thyroxamine on the metamorphosis of tadpoles of Rana temporaria, Roth (2) .- Action of sodium fluoride on the metamorphosis of Rana temporaria, Saadi-Nazim.-Metamorphosis and growth of Bufo regularis, Sedra & Michael.—Notes on the hatching glands and hatching in Polypedates [Rhacophorus] buergeri, Yanai.

Neoteny.—Factors affecting neoteny in Dicamptodon ensatus, Schuierer.

Ontogeny.—Frequency and repetition rate of frog calls related to body size, Blair (4).—The vertebral column of Bufo regularis, Sedra & Moursi.—Vascularization of respiratory surfaces in ontogeny of Ambystoma mexicanum, Czokep.—Seasonal variation in teeth of Eurycea, Stewart.

Organisers. — Diffusibility experiments on amphibian evocators, Brahma. —Neural induction in Triturus alpestris, Botenbrood. — Implants of competent ectoblast of Pleurodeles walthi to blastocoeles of P. walthi giving nervous tissue, Denis, —The role of neural and non-neural tissues in lens induction, Jacobson. —The dependence of the cellular transformation of the competent ectoderm on temporal relationships in the induction process, Tojyonen.

Organogeny.—Influence of thyroid hormone on mitotic activity of rhomboephalon of Bulo bulo, Baffoni.—Mitotic activity of spinal medulla of Bulo bulo during development, Baffoni & Pinacci.—The notochord as an inductor of the orohypophysis in Pleuvodeles wallti, Eyal-Giladi.—Development of the pancreas in Ambystoma, Frye.—Origin and development of the pronephros and Wolffian duct in Rana esculenta, Manelli (1).—Development of the apodan vertebral column, Ramaswami.—Further studies on

ear and capsule development in Ambystoma, Van Dyke & Detwiler.

Technique.—Method for sectioning refractory tissue (ovaries and eggs of *Eleutherodactylus*, *Sphenophryne* and calcified skin of *Brachycephalus*), **Griffiths & Carter**.

Teratology.—A five legged Rana pipiens from Ohio, Adler (2).—Nerve induced supernumary limbs in adult Triturus viridescens, Bodemer, -- Polydactylous juvenile Rana clamitans from Baltimore and Harford Co. Md., Cooper.-Curvature of spine in Triturus marmoratus, Freytag (2).—Spine curvature and deformed limbs in Triturus alpestris apuanus, Freytag & Hübener.—Parabiosis experiments with teratogenic larvae of Pleurodeles waltlii, Gallien (1).—Supernumerary right arm in Xenopus laevis, Hohson.—Supernumary induced limbs in Rana and Triturus helveticus, Newth (2).—Factors influencing symmetry, posture and reduplication of developing limbs of Amblystoma punctatum, Nicholas. - Effects on the nervous system of asportation and successive orchotopic grafts of limbs asportation and successive orchotopic graits of imps in Anura, Notarnicola.—Nuclear chimaeras in the newt, Pantelouris & Jacob.—Dwarf Telmatobius hauthali schreiteri attributed to growth hormone deficiency, Pisanó.—A new type of polydactyly in Rana esculenta, Rostand (1).—Effect of ultra-violet rays on larval amphibia, Rostand (2).—Importance of wounding and innervation in urodele accessory limb formation, Ruben & Frothingham. - Abnormal urostyles in Bufo regularis, Sedra. - Reduction of shin bone in doubling of frog limb in nature and occurrence of accessory limbs, Voitkevich (1).—Spatial relations between true and accessory limbs developing under natural conditions, Voitkevich (2).

## **EVOLUTION AND GENETICS**

General Works: Theories.—Comments on origin of the herpetofauna of Florida, Goin, C. J. (1).—Evolution of the species of Crinia, Neobatrachus and Heleioporus, Main, Lee & Littlejohn.—Significance of rare matings in Plethodon cinereus, Mosimann.—Amphibians in the Carboniferous Period, the role of the vertebrate visual system in the life of animals and in the origin of man, Polyak.—The origin of the Amphibia, Schmalhausen (4).—Origin and development of Amphibia, Swinton.—Evolution from fish to man via primitive Amphibia, Vandebroek.

Chromosome Problems.—Phase contrast microscopy on a mutation reducing nucleolar number in Xenopus laevis, Elsdale, Fischberg & Smith.—Experimental tetraploidy in newts, Fischberg.—Lampbrush chromosomes from Amphibian oocytes, Lafontaine & Ris.—Relation between axial complex of meiotic prophase chromosomes and chromosome pairing in Plethodon cinereus, Moses.—Chromosomes of the gonads of salamanders, Sacarrão.

Experimental.—On burnsi and kandiyohi pattern variants in Rana pipiens, Anderson & Volpe.—Effects of tryptaflavin on mitosis in Rana temporaria and Bufo bufo tadpoles, Meyer.—Interrelations between nuclei and cytoplasms of R. pipiens and R. sylvatica, cleavage rates in diploid and androgenetic hybrids, Moore (1).

Genetics of Sex.—Experiments supporting the view that sex chromosomes are cytologically unrecognisable in Bufo melanostictus, Natarajan.—The lateral loop

an

aus

me

io.

in

ord

rus

nd

tag

nic er-

OP.

118.

ire

ma

of

he

in

let

of

nb

lan

nin

ns

ler

of

lu-

io-

are

hi-

he

hi-

of via

ro-

in

es, ex

me 1es

rn

cta

nd

en

ca,

ds,

ole

op

chromosomes of Triturus pyrrhogaster and T. viridescens oocytes, Wischnitzer.

Phylogeny.—Origin and relationships of herpetofauna of S. Florida, Duellman & Schwartz.—Of the Pelobates, Pasteur (2).—Salientia phylogeny, Samtleben.—Of the Ambystomidae; family probably derived from the Hynobiidae in late Mesozoic or early Tertiary, Tihen.—Phylogeny of American Plethodontoidea, Wahlert.

Variation.—Experiments on burnsi and kandiyohi pattern variants of Rana pipiens, Anderson & Volpe.
—Population variability in Pleurodema bibroni, Cei (1).

## ECOLOGY AND HABITS

General Works.—Field guide to N. E. American amphibians; keys to the species and notes on ecology, habits, voice and breeding; figs., Conant (2).

Amphibia.—From Negros Island, Philippines, Aleals (2).—The vertebrate fauna of the Sarek mountains and the Padjelanta plain in Swedish Lapland, Curry-Lindahl.—Fauna of the Carpathians, Slovakia, Dudich.—Of some Ceylon amphibians, Morgan-Davies (3).—Some peculiarities of reptiles and amphibia of the alpine tracts of Jugoslavia, Radovanovic.—General ecology of non mammalian vertebrates of Dernford Fen., Sharp.—Biogeography and ecology of American Amphibia, Walhert.

Anura.—General study of Atelopus stelzneri, Adler, J.—Philippine notes on the ecology of Bufo marinus, Alcala (1).—Of some Somalia amphibians, Cherchi (1), (2), (3).—Salientia from Banyuls, Le Canet and the forest of Sorède, Duguy & Phillipe-Knoepffler.—Biology of Leptodactylus prognatius, Gallardo (2).—Rana japonica general ecology and biology, Geyer, H. (1).—Small tree frogs from the United States, Geyer, H. (2).—Of Hyla cinerca and H. squirella, Goin, O. B.—Frogs from Leticia, Colombia, Goin & Layne.—Descriptive list of Anura and Reptilia from the Bamburg area, Heilingbrunner.—Rana dalmatina from Sweden, König.—Life and habits of Bufo debilis, Lang (1).—Survival and disappearance rates of spadefoot toads, Pearson, (1).—Myobatrachus gouldis, its general habits and termite associations, Philipp.—Studies on ecology of middle European Salientia, Rühmekorf (2).—Distribution and general habits of Hyperolius verrucosus, Wager (1).—Life history and ecology of Hemisus guttatus from Hluhluwe, Wager (2).—Alytes obstetricans obstetricans, systematics and general biology, conditions of raising young, Wettstein.

Caudata.—Prehensility of Plethodon g. glutinosus, Brode & Gunter.—Ecology and biology of Chioglossa lusitanica, Goux.—Distribution and ecology of Eurycea lucifuga, Hutchison.—General notes on Ambystoma opacum, Lang (2).—Batrachoseps attenuatus from North America, Lang (3).—Of the genus Taricha, Riemer (2).—General biology and ecology of Batrachoseps attenuatus, Shaw.—The Hellbender, Stine & Flower.—Conditions required for rearing newts in laboratories for experimental purposes, Williams.

Aquaria, Terraria, etc.—Maintenance of Atelopus stelzneri, Adler. J.—Care of amphibians in captivity; setting up an aquarium and feeding captive amphibians;

figs., Conant (2).—Care in captivity of Chiroglessa lusitanica, Goux.—Laboratory rearing of newts for experimental purposes, Williams.

Behaviour.—Notes on habits of large Bufo bufo, Boettger.—Reaction to frogs' calls, Bogert.—Ability of Bufo calamita to track mealworms by sense of smell and yawning, Heusser (3).—Behaviour of various newt species, Sculthorpe.—Aspects of behaviour at spawning sites of the common frog, Warwick.

Biological Control.—Amphibia as predatory vertebrates, Sweetman.

Breeding and Parental Habits.—Daytime breeding in Scaphiopus, Bragg (1).—Habits of Bufor. regularis; breeding dates and reproduction, Chapman & Chapman.—Chiromantis petersi kelleri, Cherchi (2).—Pipu pipa, P. parva in Venezuela, Gines.—Reproductive behaviour of Scaphiopus holbrooki, Hansen, K. L.—Reproduction activities of Bufo bufo and study of spawning migration, Heusser (1).—Of Rhacophorus schlegelii arborea, Kato (2).—Late breeding (September) after heavy rainfall of Pseudaris clarki and P. nigrita, Kennedy.—Q Bufo bufo ×3 B. calamita near Warsaw, Klekowski.—Of Rana nigromaculata, Ling.—Mating behaviour of Hyla versicolor, Little-john (2).—Reproductive behaviour of Pseudaris nigrita, Martof & Thompson.—Spawning in the grassfrog, Mertens (2).—Of Rhacophorus c. cruciger desed. and figd., Morgan-Davies (1).—Of Ramanella obscura desed, and figd., Morgan-Davies (2).—Breeding habits, eggs and larvae of Eurycea lucifuga desed., Myers (1).—Brooding of young by Flectonotus goeldii, Oeser.—Of Acris crepitans, Pyburn.—Activities and egg laying habits of various newt species, Sculthorpe.—Of Atelopus cruciger, Sexton.—Breeding habits of Hemisus guttatus, Wager (2).—Choice of spawn site and number of spawn sites per pond by Rana temporaria, Warwick.—Behaviour in Diemictylus viridescens, Woodford.

Burrowing Habits.—Tunnels of Pseudobranchus and Siren described, Freeman.—Study of Eurycea lucifuga, Hutchison.

Cannibalism.—In Rana sylvatica tadpoles, Bleakney, S.—In Leptodactylus ocellatus, Gallardo (1).—In Rana esculenta, Vandevelde.

Courtship.—Of Pseudacris nigrita, Martof & Thompson.—Reproductive behaviour of Rhacophorus c. cruciger, Morgan-Davies (1.)—Of Ramanella obscura; desed. and figd., Morgan-Davies (2).—Behaviour of Plethodon jordani metcalfi, Organ.—Behaviour pattern in Diemictylus viridescens, Woodford.

Diseases.—Parasitism of Scaphiopus bombifrons tadpoles by Saprolegnia, Bragg, A. N. & W. N.—Skin eruptions in frogs, Jimbo & Nagasawa.

Distribution.—Of Amphibia in the United States in relation to past and present environment, Blair, (1).—Distribution of Amphibia in ricefields in Slovakia, Feriancová.—Records of Anura and Reptilia from the Bamberg area, Heilingbrunner.—Allopatric populations of Scaphiopus, Wasserman.

Ecological Associations.—Studies on the distribution and ecology of amphibians in the ricefields of Slovakia, Bratislavia, Feriancová.—General amphibian ecology of lakes, swamps and marshes in eastern Europe, Germaine.—Winter activities and habitate of Rana clamitans larvae, Getz.—Population study of Typhlotriton in Laclede Co. cave, Missouri, Hendricks & Kezer.—Amphibians of Florida caves, Holman (2).
—Missouri cave dwelling Amphibia, Myers (3).

Ecological Associations.—Amphibians occurring in saltwater areas, a review of literature and comments, Neill (2).

Economics.—Larvae of Lechriodus in Wissel Lakes, New Guinea eaten by Papuans, Brongersma L. D.— Therapeutic uses of toad venoms, Frèrejacque.—Use of herpetofauna of Florida by early Indians, Neill, Gut & Brodkorb.

Effects of Environment.—Seasonal sensitivity of Bufo to chorionic gonadotrophin, Allison.—Effect of environmental factors on anuran mating call, Blair (4).—Eruption after rain of Batrachoseps, Grant.—Seasonal variations in bullfrog carbohydrate metabolism, Hashimoto, Morita & Matsuyama.—Effects of crowding Rana temporaria tadpoles, Hodler.—Late breeding of Pseudacris after heavy rainfall, Kennedy.—Mass migration of Pelobates cultripes in heavy rain at Canet, France, Petit & Lomont.—Dependence on external factors of spawning in Salientia, Rühmekorf (1).—Action of light on the spawning of larval Salientia Rühmekorf (3).

Enemies and Defence.—Defence reactions studied in Bufo arenarum and Leptodactylus occilatus, Gallardo (1).—Glandular secretion of Phyllomedusa bicolor and Phrynohyas zonata, Goin & Layne.—Predators of Eurycea lucifuga, Hutchison.—Defence postures and reactions in Taricha and Cynops, Riemer (2).

Feeding Habits.—Mice eaten by large toads, Boettger.—Salamandra salamandra, Buehholz.—Rana catesbeiana omnivorous; stomach contents listed, Cohen & Howard.—The diets of some eastern North American amphibians, Conant (2).—Stomach contents of some species of S. Florida, Duellman & Schwartz.—Of Leptodactylus ocellatus, Gallardo (1).—Behaviour of Bufo calamita in seeking food by olfactory orientation, Heusser (3).—Ant feeding behaviour of Microhyla carolinensis, Holman & Campbell.—Eurycea lucifuga, Hutchison.—Studies on the diet of Bombina bombina, Lác.—Termites as the diet of Myobatrachus gouldii, Philipp.—Food of N. American Hylidae and Ranidae, Sweetman.—Stomach contents analysis of Rana esculenta, Tyler.

Hibernation.—Sensitivity of *Bufo* to hormones after enforced hibernation, Allison.—Urodeles at low temperatures, a study of their digestive rates during hibernation, **Joly**.

Homing.—In Bufo arenarum and Leptodactylus ocellatus, Gallardo (1).

Larval Stages.—Birth of Gastrotheca marsupiatum tadpoles, Grenard.

Longevity.—Triturus [Cynops] pyrrhogaster 24½ years in captivity, Hensley.

Movement and Migration.—Mass movement of Scaphiopus tadpoles and juveniles, Bragg (1).—
Eruptive mass migration in young Bufo cognatus; discussion of social tendencies, Bragg & Brooks.—
Study of movement and growth in a population of

Bufo r. regularis in S. W. Tanganyika, Chapman & Chapman.—Behaviour of Leptodactylus occilatus and Bufo arenarum, Gallardo (1).—Spawning migration in Bufo bufo; a population study, Heusser (1).—Experimental moving of frogs from their natural territories, Heusser (4).—Observations on Rhacophorus schlegelii, Kato (1).—Seasonal migration of Rana temporaria in Far East river Tadushe, Kurentzov.—Daily cycle of activity of Bufo boreas, Mullally.

Moults.—Skin shedding in Bufo arenarum and Leptodactylus occilatus, Gallardo (1).—Moulting behaviour in salamanders, frogs and toads; figs., Heusser (2),

Nesting Habits.—Of Chiromantis petersi kelleri, Cherchi (2).—Direct development of Arthroleptis crusculum; nest below ground, Guibé & Lamotte (2).—Nest construction by Rhacophorus c. cruciger desed, and figd., Morgan-Davies (1).—Reproduction activities and foam nests of Chiromantis rufescens and C. xerampetina, Verbaak.

Parasitism.—Parasitosis from helminths of the cystic type on batrachia near Lake Trasimeno, Gianotti.—Blood parasites after X-irradiation of newts, Jakowska et al.—Further notes of Batrachomyia larvae and their Anuran hosts, Mertens (3).—Frog and toad hosts of Nyctotherus sp. nov. from Dharwar, Uttangi.—Amphibians and their parasites from Lake Druzno, N. Poland, Wisniewski.

Population Studies.—Bufo r. regularis in S. W. Tanganyika, Chapman & Chapman.—Amphibia in a small area in Kansas, Fitch, H. S.—Comparison of the nonbreeding habits of H. squirella and H. cinerea, Goin, O. B.—Relationship between bat and insect population and population of Typhlotriton in Missouri cave, Hendricks & Kezer.—Behaviour cycle of Bufo bufo; spawning migration, Heusser (1).—Crowding effects on Rana temporaria tadpoles, Hodler.—Eurycea lucifuga, Hutchison.—Amphibians of the Gómez Farias region, Tamaulipas, Mexico; biogeographic analysis, Martin.—Population ecology of the spadefoot toad, Pearson (1).—On a population of Acris crepitans in Texas, Pyburn, W. F.—On Atelopus cruciger in Venezuela, Sexton.

Sex Relations.—Hair of *Trichobatrachus* as a possible means of discrimination by the sexes, Mertens (1).

Social and gregarious habits.—Mass migration and social tendencies of Bufo cognatus, Bragg & Brooks.

Symbiosis.—Ambystoma maculatum embryo-Oophila amblystomatis symbiosis, Hutchison & Hammen.

Technique.—Collecting amphibians; equipment, Conant (2).—Method for collecting Leptodactylus hylaedactylus, Goin & Layne.—Various trapping methods used on frogs, Heusser (4).—Methods described for marking of frogs by scarification by hypodermic needle and banding toes with cylindrical bird bands, Kaplan.—Procaine hydrochloride as a killing agent for amphibians, Livezey.—Procedure for "crowding" experiments with Rana pipiens tadpoles, Richards.

Voice.—Call differences in hylid frogs, Blair, W. F.— Analysis of anuran calls; patterns of geographic variation in call, effect of environment and ontogeny on call, Blair (4).—Absence of call difference between Bogo Cei Calli som Influ and bian Goss repr & Mor Neil

[1

Scar

pur

in to ney, Cold Schr Goir spec Mai the pas, stor

Soc C

I

Sza Die

In

Ne

No

nd

in

ri-

in

of

nd

tis

d,

108

199.-

he

10.

of

nia

ha

ar.

ke

W.

a

he ea.

et

ıri ıfo

ng

ea

ez

nie

ot

ns in

68,

nd 8.

ila

nt.

lus

ng es-

m-

ird

ng

for

08,

hic

ny

Scaphiopus holbrooki and S. hurteri, Blair (5).—On the purpose of frogs' calls with sonograms and an illustration, p. 189, showing calling stations of various species, Bogert.—Warning vibration in Pleurodema bibroni, Cei & Aguilera.—Of Rana obbiana, Cherchi (1).—Calls of North American frogs descd., Conant (2).—Of some S. Florida amphibians, Duellman & Schwartz.—Influence of voice in sex attraction of Bufo arenarum and Leptodactylus ocellatus, Gallardo (1).—Of Colombian frogs, Goin & Layne.—Of Pseudacris brimleyi, Gosner & Black (1).—Importance of the call in reproductive behaviour of Pseudacris nigrita, Martof & Thompson.—Call of Ramanella obscura descd., Morgan-Davies (2).—The varied calls of Hyla gratiosa, Neill (3).

#### GEOGRAPHICAL DISTRIBUTION

General: Zoogeography.—Distribution of Amphibia in the U.S.A., Blair (1).—Of Eastern Canada, Bleakney, J. S.—Biogeography of the Balearic Islands, Colom.—Herpetofauna of S. Florida, Duellman & Schwartz.—Origin of the herpetofauna of Florida, Goin, C. J. (1).—Distribution and evolution of the species of Crinia, Neobatrachus and Heleioporus, Main, Lee & Littlejohn.—Biogeographic analysis of the amphibians of the Gómez Farías region, Tamaulipas, Mexico, Martin.—Zoogeography of the Ambystomidae, Tihen.

## PALEARCTIC REGION

Europe.-Eiselt.

England.—(Dorset), Gooch.—(Dernford Fen, Cambridgeshire), Sharp.

Austria. — (Tirol), Hellmich (1). — (Burgenland), Sochurek.

China.-Huene (1).

France.—(Alpes Maritimes), Aellen.—(Marseilles), Berner.

Germany .- (Gössweinstein), Brunner, G.

Hungary.—Dely.—(Pilis Mt., N. W. Budapest), Szabo.

Japan.—(Honshu and Shikoku), Alexander & Diener.—Loo Choo Is. (Amami group), Koba.

Luxembourg.—Hoffmann.

Mediterranean.—Balearic Islands, Colom.

Mongolia.-Bannikov.

Poland.—Fudakowski.—(S. W. Poland), Bebak.— (near Warsaw), Klekowski.—(Cracow), Mickalowski.— (Lake Druzno, Northern Poland), Wisniewski.

Portugal.-Goux.

Russia.—Crimea, Chernov, S. A.—Kola Peninsula, Novikov. — (Transcaucasia), Rotter. — (Caucasian Isthmus), Vereshchagin.

Spain.-Goux.

Turkey.—(Anatolia), Hellmich (2).

#### ORIENTAL REGION

Ceylon.—Kirtisinghe. — Morgan-Davies (1), (2) & (3).—Silva.

India.-Huene, (2).

Maldive Islands.—Phillips.

Philippine Islands.—Alcala (1).—(Negros Island), Alcala (2).—Rabor, Alcala & Gonzales.

Sarawak.-Inger (2).

Thailand.—Taylor & Elbel.

## AUSTRALIAN REGION AND POLYNESIA

Australia.—Littlejohn (1).—Main, Lee & Littlejohn. —Moore (2).—(C. Australia), Mertens (4).

New Guinea.—Brongersma, L. D.—Zweifel (2).

#### ETHIOPIAN REGION

Central Africa.—(S. Rhodesia: Inyanga district), Fitzsimons.—(Belgian Congo), Laurent (2).—(French Cameroons), Guibé & Lamotte (1).—Perret (1) (2) & (3).

East Africa. -- Somalia, Cherchi (1), (2), (3).

South Africa. - (Eastern Cape Province), Kitching.

West Africa,—French Guinea (Doromou), Guibé & Lamotte (3).—French Guinea (Mount Nimba), Guibé & Lamotte (4).—French Guinea (Mt. Nimba), Laurent (1).—(Liberia), Taylor & Weyer.

São Tomé, Principe and Rolas Islet.-Manacas.

Seychelles.-Parker.

## NEARCTIC REGION

Canada,—Eastern Canada, Bleakney, J. S.—(East of 100th meridian), Conant (2).—British Colombia, Carl & Guiguet.—Nova Scotia, Romer.—Ontario (Long Point), Adams & Clark.

United States .- (East of 100th meridian), Conant (2).

North Eastern States. — Wasserman. — Delaware (Delmarva Peninsula), Conant (1). — Maryland (E. shore), Conant (1). — Maryland & Delmarva, Reed (1). — Maryland (Worcester Co.), Reed (2). — Maryland (Delmarva Penin.), Reed (3). — Michigan (Jackson Co.), Hahn, D. E. — Minnesola (Cass Co.), Brattstrom (1). — New Jersey, Black & Gosner. — Pennsylvania (Delmarva Penin.), Conant (1).

South Eastern States. — Blair (1). — Arkansas, Louisiana, Missouri, Wasserman.—Arkansas (Ozark Plateau and Ouachita Mts.), Dowling.—Florida, Blair, (3).—S. Florida, Duellman & Schwartz.—Florida, Goin, C. J. (1).—Florida, Holman (1).—Florida, Citrus Co. and Alachua Co.), Holman (2).—Florida, Neill, Gut & Brodkorb.—Florida (Jackson Co.), Pylka & Warren.—Florida, Riemer (1).—Florida, Stevenson.—W. & C. Georgia (Coastal Plain), Goodman.—Kentucky, Barbour.—Kentucky (Livingstone Co.), Rossman (2).—S.E. Louisiana, Volpe.—Mississippi (Wayne & Copiah Co.), Brode.—Mississippi (Winston Co.), Ferguson & Rhodes.—Missouri (Laclede Co.), Hendricks & Kezer.—Missouri, Myers

1

Hy1

fro Hy

ang

W Gu

r08

Cu

spe Sill

anl

me

wa Ta use

Rec

(in

mu

tio

(2)

Die

sul

COL

tio

Tse

pa: La

Oz

da

(2) .- E. Missouri, Myers (3) .- Tennessee (White Co.), Rossman (1) .- Virginia (Delmarva Peninsula), Conant (1).

Middle and Western States.—Arizona (Glen Canyon), Tanner. — Arizona, Williams & Chrapliwy. — S. Dakota, Woodbury.—Kansas (Lyon Co.), Clarke, S. Dakota, Woodbury.—Kansas (Lyon Co.), Clarke, Breukelman & Andrews.—Kansas, Fitch, H. S.—Nevada, Brattstorm (2).—Oklahoma, Bragg (3).—Oklahoma, Wasserman.—N.E. Oregon (Baker Co.), Ferguson, Payne & Storm.—Texas (Montague Co.), Goin & Auffenberg.—Texas, Olson.—Utah (Glen Canyon), Tanner.—Wyoming (Niobrara Co.), Goin & Auffenberg.—Wyoming (Yellowstone National Park), Turner.

Pacific Coast States,—California (O'Neals), Cohen & Howard. — California, Karlstrom. — California (Eldorado Co.), Mullally & Powell.—California (San Luis Obispo Co., 3954 ft.), Pimentel (2),-California (Lake Co.), Plutner.

#### CENTRAL AMERICA

Canal Zone.—Fouquette.

Cuba.—(Western), Schwartz (1).—(Pinar del Rio Province), Schwartz (2) & (3).—(Pinar del Rio and Oriente Provs.), Schwartz (4).

Costa Rica.-Taylor, E. H. (1).

Haiti .- Lynn.

Mexico.—Smith & Chrapliwy.—Shannon & Hum-phrey.—Duellman (2).—W. Mexico (Colima), Duellman (1).—(Gómez Farias, Tamaulipas), Martin.—(Chiapas)
Poglayen & Smith.—(N.E.), Rabb.—(Jalisco), Smith
& Grant.—(N.W. Sonora), Smith & Hensley.—
(Coahuila), Webb.—Baja California (S. Todos Santos I.), Zweifel (1).

## SOUTH AMERICA

Argentine.—Gallardo (1).

Brazil.—(Fed. Terr. of Amapa), Bokermann (1) & (2),—(Southeast), Lutz & Carvalho.

Chile. — Capurro. (1). — Cei (1) & (4). — Cei & Capurro.

Colombia .- (Leticia, Amazonas Commissary), Goin & Layne.

Venezuela.—Gines.—Sexton.

## PALEONTOLOGY AND GEOLOGICAL DISTRIBUTION

Carboniferous.-Romer.

Cretaceous.-Goin & Auffenberg.

Permian .- (Vale & Choza), Olson.

Pleistocene.—Brattstrom (2).— Brunner, G. — Holman (1).—Neill, (1).

Pliocene.-Auffenberg.-Heller.

Triassic.—Huene (1) & (2).—Kitching.

## III.—SYSTEMATIC INDEX

THE ORDERS, AND FAMILIES AND GENERA WITHIN THEM, ARE ALPHABETICALLY ARRANGED.

References to the "Titles" is by the name(s) of the Author(s) printed in clarendon type.

#### **AMPHIBIA**

General: Fossil.-+Pre-tertiary saurians of Asia, Huene (2).—† Amphibia, one gen. nov. Eryopid undescribed, from the Permian of Texas, Olson.— Classification of Amphibia with Geological occurrence,

General: Recent.—Classification of the Amphibia, Baer.—Zoogeography of the amphibians of Eastern Canada, Bleakney, J. S.—Notes on amphibians of New Guinea, Brongersma, L. D.—Amphibians of Chile, Capurro (1).—Previously unpublished plates from Philippi's Suplemento los Batraquios Chilenos; discussion on status and synonymy of species, Cei (4).—Systematic review of the amphibia of Crimea, Chernoy, S. A.—Field guide to amphibians of Eastern N. America; descriptions, figs. and colour plates, distribution maps, Conant (2).—Systematic account of S. Florida amphibians. Notes on life history, habits, variation, ecology and distribution, Duellman & Schwartz.—Salientia from the Banyuls region, Duguy Schwartz.—Salientia from the Banyus tegon, Deagn, & Phillipe-Knoepffler.—Frogs from Amazonas region, S. Colombia, Goin & Layne.—Amphibians of Mt. Nimba, French Guinea. (Arthroleptis, Phrynobatrachus and Hyperolius excluded.) Guibé & Lamotte (4).—First descriptions of tadpoles of some species of anura from Ceylon, Kirtisinghe.—Amphibians of the Gómez Farías region, Tamaulipas, Mexico; biogeo-Gomez Farias region, Tamaulipas, Mexico; blogeo-graphic analysis, Martin.—Type specimens in the Chicago Natural History Museum, Marx.—Amphibia of Kola peninsula, Novikov.—Brief list of Negros Island amphibians, Rabor, Alcala & Gonzales.— Amphibians of Glen Canyon, Colorado R., Utah, Tanner.—The Amphibia of Thailand; a list of species known from the country with some descriptions and photos., Taylor & Elbel.—Amphibia of the Caucasian isthmus, Vereshchagin.

Proposed classification of the Anura, as follows.—

Superorder Salientia Order Anura Suborder Amphicoela Family Notobatracidae Letopelmatidae Family Suborder Aglossa Family Pipidae Family Palaeobatrachidae Family Eoxenopoides Archaeobatrachia nov. Suborder Superfamily A Family Discoglossidae Family Rhinophrynidae Family Montsechobatrachidae? Superfamily B Family Pelobatidae Family Pelodytidae Suborder Neobatrachia nov. Superfamily A Family Leptodactylidae Dendrobatidae Family Family Atelopodidae Family Bufonidae Family Hylidae

IIN

the

pid

ce,

oia,

ern

ew ile,

om

dis-

OV,

N.

tri-

f S.

guy

ion,

Mt.

traotte

s of the

geo-

ibia

gros

tah,

cies

and

sian

g.

Family Pseudidae
Superfamily B
Family Ranidae
Family Hyperolidae
Family Phrynomeridae
Family Phrynomeridae
Family Heleophrynidae

Reig.

## SUPER ORDER SALIENTIA ORDER GYMNOPHIONA

## CAECILIIDAE

Caecilians of the Seychelles Is; a new subspecies of Hypogeophis descd., Parker.

Oaccilia rostrata, considered generically distinct from other Seychelles caecilians and assigned to Hypogeophis, Parker.

Dermophis larvatus Ahl, includes Hypogeophis angusticeps Parker, Parker.

Geotrypetes angeli, from Harbel, Liberia, Taylor & Weyer.—G. scraphini occidentalis, from Mt. Nimba, Guibé & Lamotte (4).

Hypogeophis, a monotypic genus. Hypogeophis rostratus considered correct name for Caecilia rostrata, Cuvier. Lectotype designated. Three insular subspecies recognised:—H. r. rostratus from Mahé and Silhouette, H. r. guentheri from Frigate, H. r. pruslini subsp. nov. p. 76 from Praslin, Parker,—H. development of vertebral column, Ramaswami.

Ichthyophis, vertebral column development, Ramaswami.—I. glutinosus, of Thailand; note on variation, Taylor & Elbel.

Prasiinia Boulenger, P. cooperi type species, to be used for following Seychelles caecilians:—Dermophis sechellenis, Hypogeophis alternans, D. larvata (includes H. angusticeps), H. brevis, Parker.

Schistometopum thomense, on bucco-pharyngeal mucous membrane, Elkan (1).

## ORDER CAUDATA

Field guide for E. North American forms; descriptions, figs., colour plates, distribution maps, Conant (2).—General systematics and distribution, Wahlert.

#### AMBYSTOMIDAE

Phylogeny of family discussed. Osteological evidence suggests division into three subfamilies: Dicamptodontinae subfamily nov., Rhyacotritoninae subfamily nov. and Ambystominae, last subfamily containing Rhyacosiredon and Ambystoma. Definitions, ranges and geographical considerations, Tihen.

Ambystoma, parabiosis experiments, Hughes & Tschumi.—Changes in the electrophoretic serum pattern, Manelli (2).—Defined and discussed, Tihen.—Later ear differentiation, Van Dyke & Detwiler.

Ambystoma annulatum, or geographic relations of Ozarkian amphibians, **Dowling**.

Ambystoma gracile, habits, life-history and general data, Carl.-A.g. decorticatum, on salt water association, Neill (2).

Ambystoma jeffersonianum, of E. Canada; zoogeography, Bleakney, J. S.—Of Cass Co., Minnesota, Brattstrom (1).

Ambystoma laterale, of Long Point, Ontario, Adams

Ambystoma macrodactylum, from Baker Co., Oregon, Ferguson, Payne & Storm.

Ambystoma maculatum, of E. Canada; zoogeography, Bleakney, J. S.—On geographic relations of Ozarkian amphibians, Dowling. — Embryo-alga symbiosis, Hutchison & Hammen.—From Piedmont of coastal Delmarva, Reed (3).—Radioresistance in larvae, Stinson.—A. maculatum group, defined and discussed. Referred forms: — A. g. gracile, A. g. decorticatum, A. jeffersonianum, A. laterale, A. m. macrodactylum, A. m. croceum, A. maculatum, Tihen.

Ambystoma mexicanum, ontogeny of vascularisation of respiratory surfaces, Czopek.—Mast cells in the choroid plexus, Kappers, Ten Kate & Bruyn.—Regeneration of the central nervous system in the early developmental stages, Romanovsky.—A. mexicanum group, defined and discussed. Forms referred:—A. kansense, A. lermaensie, A. mexicanum, Then.

Ambystoma opacum, retardation of fat body, prevention of Müllerian duct formation, Asayama & Matsuzaki.—Ultraviolet irradiation of upper arm of larva and its effect on regeneration of the lower part, Butler, Blum & Schmidt.—On geographic relations of Ozarkian amphibians, Dowling.—Development of the pancreas, Frye.—From U.S.A. general, Lang (2).—Hosts for accessory limb formation experiments, Ruben & Frothingham.—Radioresistance in larvae, Stinson.—A. opacum group, defined and discussed. Forms referred:— A. opacum, A. talpoideum, Tihen.

Ambystoma punctatum, tail growth, part played by notochord, Bijtel.—Regenerates on reversed hind limbs, Deck & Riley.—As donor of neural crest material, Finnegan.—Larvae in skeletal regeneration experiments, Goss.—Fine structure of blastema cells and differentiating cartilage cells in regenerating limbs, Hay (2).—Influence of the retina on lens development, Jacobson.—Control of limb development, Nicholas.—Lens induction, Reyer.—Perfusion of blastocoel, Shapiro.—Inhibition of limb regeneration, Thornton.

Ambystoma subsalsum, on ability to withstand salt water, Neill (2).

Ambystoma talpoideum, inhibition of limb regeneration, Thornton.

Ambystoma texanum, habitat in Lyon Co., Kansas, Clarke, Breukelman & Andrews.—On geographic relations of Ozarkian amphibians, Dowling.

Ambystoma tigrinum, habitat in Lyon Co., Kansas, Clarke, Breukelman & Andrews.—On geographic relations of Ozarkian amphibians, Dowling.—On bucco-pharyngeal mucous membrane, Elkan (1).—Radioresistance in larvae, Stinson.—Inhibition of limb regeneration, Thornton.—A. tigrinum group, defined and discussed. Forms referred:—A. amblycephalum, A. bombypellum, A. fluvinatum, A. granulosum, A. hibbardi, A. lacustris, A. ordinarium, A. r. rosaceum, A. r. nigrum, A. r. sonoraensis, A. subsalsum, A. t. tigrinum, A. t. californiense,

A. t. diabloi, A. t. mavortium, A. t. melanostictum, A. t. nebidosum, A. t. stebbinsi, A. t. velasci, Tihen.—A. t. tigrinum, from Cass Co., Minnesota, Bratistrom (1).—A. t. utahensis, of Glen Canyon, Tanner.

† Ambystomichnus, known only from trackways, Tihen.

Bathysiredon, a subgenus of Ambystoma differing from A. mexicanum group in cranial elements. Definition and discussion. Includes A. (B.) d. dumerili, A. (B.) d. queretarensis, Tihen.

Dicamptodon, defined and discussed; range, Tihen.

—D. eneatus, factors affecting neoteny, Schuierer.

Linguaelapeus, a subgenus to include A. annulatum. A. c. cingulatum, A. c. bishopi, A. mabesi, A. schmidti, A. texanum. Definition, discussion and range. Radiograph of type of A. schmidti, Then.

Rhyacosiredon, defined and discussed. Referred forms:— R. altamirani, R. leorae, R. rivularis, R. zempoalensis, Tihen.

Rhyacotriton, defined and discussed; range. Referred forms, R. o. olympicus, R. o. variegatus, Tihen.—Rhyacotriton olympicus variegatus, vital limits and desiccation, Ray.

Siredon mexicanum, irradiation of parts of the head, Brunst.—Grafting of the nasal placode to the fore limb area, Kiortsis (2),—Inversion of the dorso-ventral axis, Levtrup & Pigon.—Tail regeneration after abnormal amputations, Newth (1).—Respiration during development, Rudnick.—Neural closure and the forces involved, Selman. (1).

#### AMPHIUMIDAE

Amphiuma means, Of S. Florida; habitat, stomach contents, nests, range, Duellman & Schwartz.—On bucco-pharyngeal mucous membrane, Elkan (1).—Late Pleistocene recerd from Florida; doubts on accuracy of age of remains due to rapid mineralisation of bones, Neill (1).

## CRYPTOBRANCHIDAE

Ecology and world distribution, Wahlert.

Cryptobranchus a. alleganiensis, C. a. bishopi, on geographic relation of Ozarkian amphibians, Dowling.—C. japonicus, head musculature, Schumacher (1).—Musculature, Schumacher (2).

Megalobatrachus japonicus, histology of teeth, Schmidt, W. J. (2).—Commissural systems of the cerebellum, Tamura, Yashiki, Kondo & Numata.

## HYNOBIIDAE

Hynobius dunni, alkaline phosphatase of the retina in late embryogenesis, Yoshida (2).—H. keiserlingii, cessation of regeneration by synthetic cancer causing substances, Tokin & Filatova.—H. nebulosus, sex development modified by methyl-testosterone, Asayama & Amanuma.—Estrogen administration to larval gonads, Asayama & Miyamori.—Kinetic properties of the marginal zone of eggs, Irushima.—Iodine accumulation in pigmented tissues, Koshida (1).—H. retardatus, water absorbtion by the eggs, Aoki.—Effect of follicular hormone on sex differentiation, Hanaoka (1).—Effect of testosterone-propionate on sex differentiation, Hanaoka (2).

#### PLETHODONTIDAE

Phylogeny and map of distribution in North America, Wahlert.

Aneides ferreus, A. flavipunctatus niger and A. lugubris, vital limits and desiccation, Ray.

Batrachoseps attenuatus, inducted oviposition and incubation of eggs, Anderson.—On bucco-pharyngeal mucous membrane, Elkan (1),—Eruption of individuals from a crack after rain, Grant.—From North America, Lang (3).—From Pine Mt., 3954 ft., San Luis Obispo Co., California, Pimentel (2).—Vital limits and desicoation, Ray.—From San Diego, general notes on its study, Shaw.—B. pacificus leucopus, from S. Todos Santos Island, Baja California, Zweifel (1).

Bolitoglossa, osteological study of following forms:—platydactyla, rufescens, occidentalis. Comparison with Salamandra and Plethodon; figs., Hansen & Tanner.—B. occidentalis, B. rufescens, of Chiapas, Poglayen & Smith.

Chiropterotriton, N. E. Mexico species divided into two groups on basis of size and dentition. Species and subspecies recognised:— C. multidentatus (photo.), C. arboreus; C. mosaueri; (photo.); C.c.chondrostega; C.c.terrestris (photo.); C.c. cracens subsp. nov., Agua Linda, 5950 ft., \(\dip 7\) mis. W.N.W. Gómez Farias, Tamaulipas, p. 28; C. dimidiatus, Rabb. —C. chondrostega, C. multidentata, habitat at Gómez Farías, Tamaulipas, Martin.

Desmognathus fuscus, thyroid activity and the effect of goitrogens, Dent & Lynn.—On geographic relations of Ozarkian amphibians, Dowling.—Structure of meiotic chromosomes from primary spermatocytes, Moses.—Radioresistance in larvae, Stinson.—D. fuscus conanti subsp. nov. p. 158, 400 ft., 2·1 mls. S. Smithland, Livingstone Co., Kentucky, Rossman (2).—D. f. fuscus, of E. Canada; zoogeography, Bleakney, J. S.—Low temperatures, hibernation and its effect on digestion, Joly.—Of Piedmont, coastal Delmarva, Reed (3).—D. ochrophaeus, on buccal and pharyngeal mucous membranes, Elkan (1).—D. ochrophaeus carolinensis, check skin color parallelisin, Reynolds.

Ensatina eschscholtzi, on bucco-pharyngeal mucous membrane, Elkan (1).—E. e. xanthoptica vital limits and desiccation, Ray.

Eurycea bislineata, chromosome numbers in larval somatic cells, Fankhausfr & Dluhy.—E. b. bislineata, of E. Canada; zoogeography, Bleakney, J. S.—Effect of low temperatures and hibernation on digestive rates, Joly.—Of Piedmont of coastal Delmarva, Rec (3).—Seasonal variation in teeth, Stewart.—E. longicaula guttolineata, from Mississippi caves, Brode.—From cave in Mississippi, Brode & Gunter.—E. l. longicaula guttolineata, from caves, Myers (3).—E. longicaula melanopleura, E. lucifuga, E. m. multiplicata, E. m. griseogaster, E. lucifuga, E. m. multiplicata, Compression of Ozarkian amphibians, Dowling.—E. lucifuga distribution, ecology and physiology, Hutchison.—Notes on eggs and larvae; hatchling figd., Myers (1).—On occurrence in E. Missouri caves, Myers (3).

Gyrinophilus porphyriticus inagnoscus, type locality restricted to Salt Creek, 4 mls. S.W. of S. Blooming-ville, Salt Creek Township, Hocking Co., Ohio, Condit.—G. p. porphyriticus, of E. Canada; zoogeography. Bleakney, J. S.

hab geo of coa Flo

[]

E

ton figs Elk pho Dis

rela P. Rel chrof den E. Win Rho des dist

On

Flo sipp pre In cav twe Dec P. des

of P. and coa Mis

Oz

hal

for rec

Ka gra Ind stu Cla

me

rth

gu-

ınd

eal

vi-

rth

ital

om i).

i:—

ner.

nto

tus ();

W.

mez

fect

ons

of tes,

ith-

ney,

fect

rva,

geal

ieus

ous

rval

ata,

fect

tes,

uda

rom

uda, uda

eata,

stri-

s on On

lity

ing-

phy,

).-

8.

Haideotriton of. wallacei, from Jackson Co., Florida; habitat; photo., Pylka & Warren.

Hemidactylium scutatum, of E. Canada; zoogeography, Bleakney, J. S.—On geographic relations of Ozarkian amphibians, Dowling.—Of Piedmont, coastal Delmarva, Reed (3).—From near Tallahassee, Florida, Stevenson.

Hydromantes, On characters and distribution, tongue action; note on reproductive behaviour; figs., Elkan (2).—List of species and distribution, Elkan (3).—H. italicus strinatii subsp. nov., p. 156, photo., Aspremont cave, Alpes Maritimes, France. Distribution of species discussed; maps, Aellen.

Plethodon caddoensis, P. cinereus angusticlavius, P. c. serratus, P. glutinosus P. ouachitae, geographic relations of Ozarkian amphibians, Dowling. P. cinereus, red and black pigments, Hertzler.—Relation between axial complex of meiotic prophase chromosomes and chromosome pairing, Moses.—Study of evolutionary significance of rare matings in low density populations, Mosimann.—P. c. cinereus, of E. Canada; zoogeographic, Bleakney, J. S.—From E. Missouri cave, Myers (3).—P. dorsalis, from Winston Co., Miss., range extension, Ferguson & Rhodes.—P. dunni and P. elongatus, vital limits and desiccation, Ray.—P. glutinosus, map. p 451, of distribution in the Southern United States, Blair (1).—On bucco-pharyngeal mucous membrane, Elkan (1).—P Plethodon glutinosus, Pleistocene of Saber-tooth cave, Florida, Holman (1).—P. g. glutinosus, from Missisaippi caves, Brode.—Egg masses desecd, note on prehensile tail of cave specimens, Brode & Gunter.—In association with Eurycea lucifuga in E. Missouri cave, Myers (3).—P. jordani jordani, parallelism between check skin colour and the "imitator" variety of Desmognathus ochrophaeus carolinensis, Reynolds.—P. jordani metcalfi, courtship and spermataphore deeed, and figd., Organ.

Pseudoeurycea belli, P. cephalica, P. scandens; habitat in Gómez Farías, Tamaulipas, Martin.

Pseudotriton montanus, electron microscope study of lampbrush chromosomes, Lafontaine, & Ris.—P. r. ruber, on bucco-pharyngeal mucous membrane and capillary network, Elkan (1).—Of Piedmont, coastal Delmarva, Reed (3).—P. ruber vioscai, from Mississippi caves, Brode.

Typhlotriton spelaeus, on geographic relations of Ozarkian amphibians, Dowling. — Population and ecological study in Missouri cave, Hendricks & Kezer. —From E. Missouri cave, Myers (3).

#### PROTEIDAE

Necturus diagnoses and keys to eastern N. American forms, survey of distribution. Four distinct species recognised:— punctatus, beyeri, lewisi, maculosus. Discussion on relationships and taxonomic characters; maps, Hecht (2).—N. maculosus, habitat in Lyon Co., Kansas, Clarke Breukelman & Andrews.—On geographic relations of Ozarkian amphibians, Dowling.—Induced ovulation, Fitch, K. L.—Electron microscope study of lampbrush chromosomes, Lafontaine & Ris.—N. m. maculosus, of Long Point, Ontario, Adams & Clark.—Of E. Canada; zoogeography, Bleakney, J. S.

Proteus anguinus, on buccal and pharyngeal mucous membranes, Elkan (1).

#### SALAMANDRIDAE

Systematics, general biology and American distribution, Wahler.

Chioglossa lusitanica, from the Balearic Islands, Colom.—Ecology, biology and distribution; map and note on care in captivity, Goux.

Cynops ensicauda, of Amami group, Loo Choo Is., Koba.—[C.] Triturus pyrrhogaster from Shikoku, Japan, Alexander & Diener.—Gonadial response to methyltestosterone, Asayama & Matsuzaki.—Experimental tetraploidy attempts, Fischberg.—24 years 6 months in captivity, Hensley.—Kinetic properties of the marginal zone of eggs, Ikushima.—Lons induction, Jacobson.—Yolk platelets structure, Karasaki & Komoda.—Differentiating potencies of the dorsal part of the blastopore lip, Kato (3).—Formation of primordial germ cells, Kotani.—Erythrocytes stained with o-amino-azo-toluene: Nagatani.—Study of the lateral loop chromosomes of oocytes, Wischnitzer.

Diemictylus viridescens, supernumary limb development, Bodemer.—Regenerates on reversed hind limbs., Deck & Riley.—Goitrogens studied from their effect on thyroid activity, Dent & Lynn.—Geographic relations of Ozarkian amphibians, Dowling.—Somatic chromosome constancy in larvae, Fankhauser & Dluhy.—In regeneration experiments, Goss.—Lower jaw regeneration, Goss & Stagg (2).—Jaw regeneration, Goss & Stagg (2).—Have refive studies on hypophysectomised efts, Grant & Grant.—Effect of thyroxine on limb regeneration, Hay (1).—Radiobiology, Jakowska et al.—Electron microscope study of lampbrush chromosomes, Lafontaine & Ris.—Intestinal regeneration, O'Steen.—Rhythmic electrical activity of skeletal muscle, Peters et al.—Accessory limb formation, Ruben & Frothingham.—Regeneration in athyroid newts, Schmidt, A. J.—Study of the lateral loop chromosomes of occytes, Wischnitzer.—Breeding behaviour, Woodford.—D. v. louisianensis. From E. Missouri cave, Myers (3).—Notophthalmus v. piaropicola, of S. Florida; habitat, variation, breeding dates, stomach contents, Duellman & Schwartz.—D. v. viridescens, of Long Point, Ontario, Adams & Clark.—Of E. Canada; distribution, Bleakney, J. S.—Metabolic effect of cortisone acetate during limb regeneration, Manner.—From Piedmont of coastal Delmarva, Reed (3).—Studies on lens induction, Stone.—Lens regeneration and retinal factor, Stone.—Lens regenerati

Euproctus montanus, on bucco-pharyngeal mucous membrane, Elkan (1).

Mertensiella caucasica, of Transcaucasia; photo., Rotter.—M. luschani, descrip. of Hamburg Inst. specimen and study of skull (photo. and X-ray). Considered intermediate between M. caucasica and Salamandra, Bašogiu & Freytag.

Molge, see under Triturus.

Notopthalmus, see under Diemictylus.

Pleurodeles, Algerian Tunisian population belong to P. poireti (includes P. hagenmülleri); diatinguished from P. waltli. Discussion on Iberian and Morocoan populations of P. waltli, Pasteur (1).—P. waltli, parabiosis experiments, Beetschen (1).—Cytological study of lethal heteroploid embryos, Beetschen (2).—Hepatic glycogen distribution, Bellee (1).—Effect on

Fer Toi gui Vai tag

1

8-8

Sch

8

Lou

and

Nei

cap

for

F

E.

(col

Cor

vall

hyb

voc

Her

Ble

(1). And

1

I

embryonic development of returning eggs, Brouant & Cury.—Competent ectoblast and nervous differentiation, Denis.—On bucco-pharyngeal mucous membrane, Elkan (1.)—The notochord as an inductor, Eyal-Giladi.—Chromosome number constancy in larval somatic cells, Fankhauser & Dluhy.—Larval parabiosis experiments, Gallien (1).—Induction of vertebrae and surrounding tissues, Gallien (2).—Action of phenol on egg mitoses and segmentation, Sentain.

Salamandra atra, temperature preference, Freytag (3).—Believed not to occur in Poland, Fudakowski.—S. salamandra, from Marseilles, Berner.—Feeding habits, Buchholz.—Taxonomic study based on colour and pattern. Races considered valid and figd.:—salamandra, algira, almanzoris, bejarae, corsica, fastuosa, gallaica, gigliolisi, infraimmaculata, semenovi, terrestris. Colour plates of typical form and of gallaica. Diagnoses, distributions (maps), synonymies and biological notes given, Eiselt.—On buccopharyngeal mucous membrane, Elkan (1).—Distribution in Poland, Fudakowski.—Regeneration of the tail region in larval forms, Kiortsis & Droin.—Regeneration of reversed polarity grafts, Oberheim & Luther.—From N.W. Budapest; habitat, Szabo.—Cytological study, Ziemann.—S. s. quadrivirgata, temperature economy, Freytag (3).—S. maculosa [S. salamandra], S. s. salamandra, S. s. taeniata, digestion at low temperatures, Joly.

Taricha, species and subspecies recognised:—
t. torosa, t. sierrae, t. granulosa, t. mazamae, t. twittyi,
t. simulans, rivularis, Pimentel (1).—Study of variation, systematic relationships habits, ecology and
distribution. Species recognised:— granulosa, torosa
affir rivularis. Recognition of T. g. similans and
T. g. twittyi and T. torosa klauberi not justified but
T. g. mazamae and T. t. sierrae upheld; figs., Riemer
(2).—T. granulosa, considered a subspecies of torosa;
discussion of validity, Pimentel (1).—T. rivularis, from
S. of Kelseyville, California, Plutner.—T. torosa,
spleen development and experimental modification,
DeLanney.—The hypomere and chromatoblast migration, Finnegan.—Gradient potentials across ectodermal flank of larva, Flickinger & Blount,—Lens
induction experiments, Jacobson.—Epiphysectomy
and scoliosis, Kelly.

Triton, see under Triturus.

Triturus alpestris, inactivation of newt larvae by benzimidazole, Billett (2).—Organisation in aggregates of neural plate cells, Boternbrood.—Early gastrulae and evocators, Brahma.—Frontal and saggital constrictions of blastulae, Brice.—Embryonic development and thermal gradients, Cury & Dollander.—Habitats and distribution in Hungary, Dely.—Effect of egg turning on cortical permeability gradient, Dollander.—Distribution in Poland, Fudakowski.—Effects of chemical solutions on the larval heart, Jullien, Ripplinger, Cardot & Marcot.—Grafting of the nasal placode to the fore limb area, Kiortsis (2).—Forelimb regeneration under the effects of drugs and hormones, Niwelinski.—Protein synthesis in occytes, Pantelouris.—Nuclear chimaeras, Pantelouris & Jacob.—Chemical inhibitors of embryonic differentiation, Schenk & Töndury.—Development of hard dentine in teeth, Schmidt, W. J. (1).—Neural closure and the forces involved, Selman (2).—Donors of

neural crest material, Seno & Nieuwkoop.—Development of gonad and sex cells, Stärk (1).—Studies on the gonad development, Stärk (2).—Effects of cytotoxic substances on embryos, Waddington.—Induction fields in young embryos, Woellwarth.—T. a. alpestris, on specimen with flecked venter, Freytag (1).—Temperature preference, Freytag (3).—Digestion under low temperature conditions, Joly.—T. alpestris apuanus, deformed limbs and spine curvature, Freytag & Hübener.

Triturus alpinus, behaviour and reproduction, Sculthorpe.

Triturus boscai, chromosomes, Sacarrão.

Triturus cristatus, transplant experiments, Bondi. Lampbrush chromosomes from oocytes, Callan & Effects of benzopyrene injections, Capuani (1).—Benzopyrene treatment, Capuani (2).—Of Crimea, Chernov.—Regeneration of striated muscle under the action of vitamins, Degan & Asandei.—On bucco-pharyngeal mucous membrane, Elkan (1),-Histology and biochemistry of the heart, examination of surviving parts, Gyévai.—Effect of combined ovarectomy, male sex hormones and nervous supply change on the production of a crest in females, Kiortsis (1).-Grafting of the nasal placede to the forelimb area, Kiortsis (2).-Cytology of induced neoplasms by grafting methylcholantrene, Leone.— Development and variation of gonad and sex cells, Stärk (1),-Effect of cancer causing substances on regeneration, **Tokin & Filatova.**—T. cristatus carnifex effect of lesions of the median eminence on the hypothalamo-hypophysis, adenohypophysis, thyroid and testicle, Mazzi.—T. c. cristatus, temperature preference, Freytag (3).—T. cristatus karelinii, with adenocarcinoma of the skin behind the head, Stolk (1).

Triturus helveticus, effect of frontal and sagittal blastulae constrictions, Brice.—Grafting of the nasal placode to the fore limb area, Kiortsis (2).—Supernumary induced limbs, Newth (2).—T. palmatus [helveticus], Protein synthesis in occytes, Pantelouris.—In experiments to produce haploid embryos, Selman (2).—Action of phenol on egg mitoses and segmentation, Sentein.—Activities and egg laying habits, Sculthorpe.—T. h. helveticus, temperature preference, Freytag (3).—Low temperatures and digestion, Joly.

Triturus marmoratus, on bucco-pharyngeal muccus membrane, Elkan (1).—On spine curvature, Freytag (2).

Triturus taeniatus, mode of tail growth, Bijtel.—Moulting behaviour; fig., Heusser (2).—Larval incompatibility on grafting, Rossi.—Development and variation of gonad and sex cells, Stärk (1).—Multiple fibroma of the adepidermal reticular network in the skin, Stolk (2).—Species-specific tumours of the skin, Stolk (3).—Effect on regeneration of cancer causing substances, Tokin & Filatova. — Comparative cytology, Ziemann.

Triturus viridescens, see under Diemictylus.

Triturus vulgaris, correlation between developmental aspects, Becker.—Artificially inseminated eggs, Fischberg.—On breeding in brackish pools, Reill (2).—Development of hard dentine in teeth, Schmidt, W. J. (1).—Behaviour and reproduction, Sculthorpe. — Experiments to produce haploid embryos, Selman (2).—In backwaters of Dernford

Fen, Sharp.—From N.W. Budapest; habitat, Szabo.—Cellular transformation and induction processes, Toivonen.—Reaction of gastrula to implants of guinea pig liver etc. treated with rabbit immune sera, Vainio.—T. v. vulgaris, temperature preference, Freytag (3).

#### SIRENIDAE

Ecology and world distribution, Wahlert.

† Prosiren gen. nov. p. 449, type species P. elinorae sp. nov. pp. 449, 450, fig., from Trinity beds of Lower Cretaceous, Montague Co., Texas, Goin & Auffenberg.

Pseudobranchus striatus, on burrowing ability; S-shaped tubes desed., Freeman.—P. s. belli, of S. Florida; habitat, food and variation, Duellman & Schwartz.

† Adelphesiren gen. nov., type species A. olivae sp. nov. pp. 453, 454, figs., from Lance Formation, Upper Cretaceous, Niobrara Co., Wyoming, Goin & Auffenberg.

Siren intermedia nettingi, from brackish water Louisana, Neill (2).—S. lacertina, Variation, habitat and stomach contents of S. Florida individuals, Duellman & Schwartz.—On burrowing habit; tunnels desed., Freeman.—From brackish water in Florida, Neill (2).

#### ORDER ANURA

#### ATELOPODIDAE

Atelopus cruciger ecology and breeding behaviour, Sexton.—Atelopus stelzneri, methods of keeping in captivity and general habits, Adler. J.

## BRACHYCEPHALIDAE

 $Brachycephalus\ epihippium\ sectioning\ technique$  for calcified skin, Griffiths & Carter.

Dendrobates, see under Dendrobatidae.

Phyllobates, see under Dendrobatidae.

## BREVICIPITIDAE SOO MICROHYLIDAE

#### BUFONIDAE

Plates from Philippi's Batraquios Chilenos; discussion on status and synonymy of species, Cei (4).—E. North American species and subspecies desed. figd. (colour plates) and distinguished; distribution maps, Conant (2).

Bufo. Criteria for identification of larval Bufo valliceps. B. terrestris americanus, B. t. terrestris. B. woodhousei fowleri and discussion on natural hybridization, Gosner & Black (2).

Bufo alvarius, on the degenerating condition of the vocal sac, Inger (1).—From N.W. Sonora, Smith & Hensley.

Bufo americanus, of E. Canada; zoogeography, Bakmey, J. S.—Of Cass Co., Minnesota, Brattstrom (I).—Of Lyon Co., Kansas, Clarke, Breukelman & Andrews.—Bufo arenarum, sperm elimination, Allende & Astrada.—Influence of diuretic hormones, Becerra.—Action of LH and FSH on spermiation, Burgos &

Houssay.—Behaviour, Gallardo (1).—Studies on sulphydrile groups in the brain, Lasansky & Sabatini.—Sperm inhibition by hormones, Moisset de Espanés.—Effect of adrenaline and noradrenaline and the Laewen Trendelenburg preparation, Nacimiento.—Castrated and treated with sex hormones, Penhos & Cardeza.—In Florida with notes on its introduction, Riemer (1).—Inorganic phosphate in nucleoli of oocytes, Tandler.—Seasonal variations in the action of antidiuretics, Uranga.

Bufo asper, from Thailand; photo., Taylor & Elbel.

Bufo boreas, irradiation of tissues and the subsequent effects, Allen.—Range in California; sympatric with B. canorus, Karlstrom.—B. b. boreas, trom Baker Co., Oregon; note on habitat and breeding date, Ferguson, Payne & Storm.—B. b. halophilus, study of visual pigments during metamorphosis, Crescitelli.—B. b. halophilus, B. b. boreas, on daily cycle of activity, Mullally.

Bufo bufo, sensitivity to chorionic gonadotrophin, Allison.—Influence of thyroid hormone on mitotic activity of rhombocephalon, Baffoni.—Mitotic activity of spinal medulla during development, Baffoni & Pinacci.—Metamorphosis, Baldisserotto & Ugurgieri. From Marseille, Berner.—Studies on germ cells in eggs, embryos and larvae, Blackler.—Notes on behaviour of large specimens; feeding on mice, Boettger. Of Crimes, Chernov.—Activity of streptomycin on the development and differentiation of eggs and larvae, Chiuini.—Embryonie ribonucleie acids study, Coromaldi (2).—Population study of behaviour cycle; factors affecting spawning migration, Heusser (1).— Moulting behaviour, Heusser (2).—Trapping and experimental changes of habitat, Heusser (4).—Effect of synthetic lysine-vasopressin on adenohypophysial activity, Jørgensen & Nielsen.-Cartilage tissue staining using indirect peroxidase reaction, Kawasaki.—Afferent fatigue and habituation after skin stimulation, Kuczka.—Active sodium transport by the isolated bladder, Leaf, Anderson & Page.— Utilization of lipids during embryonic development, Lernia. - Efferent fibres in the optic nerve and recovery of nerve after section, Maturana.—Pigmentation study, Meng.—Successive orthotopic limb grafts, Notarnicola.—Cortical change at fertilization in eggs, Osanai. - As host to limb bud implants from B. viridis, Perri (2).—Cytochrome oxidase and respiratory rate during larval development and metamorphosis, Petrucci.—Larvae and the effect of ultra violet light, Rostand (2).—Spawning and the environment, Rühmekorf (1).—Temperature selection by larvae, Rühmekorf (2).—Effect of light on spawning, Rühmekorf (3).—From Dernford Fen, Sharp.—Blood pressure and pressure pulses in arterial arches, Simons.—Studies on "double, pueleus" in developments. "double nucleus" in development; its role and frequency, Sirakami (1) & (2).—From N.W. Budapest; habitat, Szabo.—Cytological study, Ziemann.— B. b. bufo, effects of tryptaflavin on mitosis in tadpoles, Myer.—B. bufo formosus, fluorescent skin colours of the pterin type, Hama & Obika.—Metamorphosis and effect of hormones, Kobayashi,-Influence of temperature on energy-rich phosphate content of embryos. Moriwaki.—Effect of boric acid on egg development, Takeuchi, T.—Split organiser induction of double Takeuchi, T.—Split organiser induction of double embryos, Yamamoto.—B. vulgaris [bufo] japonicus, from, Honshu, Japan, Alexander & Diener.— Enzymes in the retinal receptor cells, Yoshida (1).

& ons,

m-

on

to-

ion

ris,

der

ua-

on,

On ion ned ply les, the

on ifex the roid ture with (1).

peratus uris. yos, and ying ture iges-

asal

el.—
l inand
tiple
the
skin,
using

ative

cous

elopnated sools, eeth, etion, ploid nford Bufo calamita, of Marseilles, Berner,—On tracking down prey by smell; and yawning behaviour, Heusser (3).—In the neighbourhood of Warsaw, Klekowski.—Spawning and the environment, Rühmekorf (1).—Effect of light on spawning, Rühmekorf (3).

Bufo c. camerunensis, from Mt. Nimba; photo. Distinguished from latifrons, tuberosus and regularis. Range and habitat, Guibé & Lamotte (4).

Bufo canorus, range in California; sympatric with B. boreas, Karlstrom,—From Eldorado Co., California at 7,500 ft. B. boreas-like individual from 8,000 ft.; species may hybridize, Mullally & Powell.

Bufo cognatus, social behaviour in young; mass migration, Bragg & Brooks.—From N.W. Navajoa; range extension, Smith & Chrapliwy.—Of N.W. Sonora, Smith & Hensley.—Of S. Dakota, Woodbury.

Bufo c. compactilis, from Aguascalientes, Smith & Chrapliwy.

Bufo debilis, general description, Lang (1).—
B. d. kelloggi, from Sinaloa; note on intergradation with B. d. retiformis. Hybrids from 33 mls. S.S.W. Navajoa, Sonora; range extension, Smith & Chrapling.—B. d. retiformis, range extension in Pima Co., Arizona, Williams & Chrapliny.

Bufo gardoensis, ecology and variation, Cherchi (3).

Bufo gargarizans, changes in the histochemistry of the oviduct after egg laying, Chu & Chin.

Bufo güntheri, from Limbé, Haiti; breeding dates, call, sex dimorphism, size, Lynn.

Bufo hemiophrys, of S. Dakota, Woodbury.

Bufo holdridgei, photograph of type, Taylor, E. H. (1).

Bufo horribilis, of Colima, Mexico; distribution, Duellman (1).—Habitat in Gómez Farías region, Tamaulipas, Martin.

Bufo kelaartii, first description from Ceylon, Kirtisinghe.—Of Ceylon, Morgan-Davies (3).

Bufo luetkenii, of Costa Rica; photo., Taylor, E. H. (1).

Bufo marinus, ecology in the Philippine Islands, Alcala (1).—Effect of vasopressin on water uptake whilst in different hypotonic solutions, Bentley (1).—Effect of pituitrin on water transfer across the wall of the isolated urinary bladder, Bentley (2).—Loss of sodium from the skin when dehydrated, Bentley (3).—Distribution of inorganic phosphate in muscle, Briner, Simon & Shaw.—On introduction into New Guinea, Brongersma. — Blood system of the hypophysis, Grus.—On occurrence near Miami and introduction into S. Florida, Duellman & Schwartz.—Investigation of fat, Fromm.—Habitat in Letticia, Colombia, Goin & Layne.—Moulting behaviour; fig., Heusser (2).—Active sodium transport by the isolated bladder, Leaf, Anderson & Page.—Seasonal variations in drug sensitivity of the isolated heart, Nayler.—Note on ability to withstand saline conditions, Neill (2).—In Florida, with notes on its introduction, Riemer (1).—Chemical relationships in the sartorius muscle, Simon, Shaw, Bennett & Muller.—Extracellular potassium on intracellular heart action currents, Trethewie & St. Clair.

Bufo marmoreus, distribution in W. Mexico, Duell-man (1).—From Jalisco, Smith & Grant.

Bufo mazaltanensis, of Jalisco; B. nayaritensis considered a synonym, Smith & Grant.

Bufo melanochloris, of Costa Rica; note on variation in specimens; photos., Taylor, E. H. (1).

Bufo melanostictus, induced metamorphosis of tadpoles, Bhattacharya & Medda.—Life history notes; photo., Morgan-Davies (3).—Study of male and female chromosomes, Natarajan.—From Malé, Maldives, Phillips.—From Thailand, Taylor & Elbel.

Bufo microtympanum, of Coylon, Morgan-Davies (3).

Bufo nayaritensis, considered a synonym of B. mazatlanensis, Smith & Grant.

Bufo occidentalis, of Colima; distribution in W. Mexico, Duellman (1).

Bufo paracnemis, action of pitressine and adrenalin on water balance during and after metamorphosis, Hadler, Berezin & Lison.—On its introduction into Florida, Riemer (1).

Bufo punctatus, from late Pleistocene cave in Nevada, Brattstrom (2).—Habitat in Gómez Farías, Tamaulipas, Martin.—From N.W. Sonora, Smith & Hensley.—Of Glen Canyon, Tanner.

Bufo quercicus, of S. Florida; habitat, call, stomach contents, range. Preyed on by Heterodon and Thamnophis, Duellman & Schwartz.

Bufo regularis, from Mt. Nimba; photo.; habitat; larva desed. and figd., Guibé & Lamotte (4).—Origin of nucleoli in ovocytes, Guyénot.—On abnormal urostyles, Sedra.—Metamorphosis and growth of the hypobranchial apparatus, Sedra & Michael.—Ontogenesis of vertebral column, Sedra & Moursi.—Response to pituitrin injection on fluid distribution, Wilson & Archdeacon.—B. r. regularis, Population study of movement and growth; reproduction, breeding dates, food. Chapman & Chapman.—From Inyanga district, S. Rhodesia, Fitzsimons.

Bufo rubropunctatus, distribution and colour, Capurro (1).

Bufo spinulosus, distribution, Capurro (1).

Bufo stomaticus, of Ceylon, Morgan-Davies (3).

Bufo superciliaris from Mt. Nimba; photo. Habitat, primary forest, Guibé & Lamotte (4).

Bufo terrestris, on its introduction into Florida, Riemer (1).—B. t. americanus, on geographic relations of Ozarkian amphibians, Dowling.—Of Piedmont, coastal Delmarva, Reed (3).—Of S. Dakota, Woodbury.—B. t. charlesmithi, eggs desed. and compared with those of other terrestris subsp., Bragg (2).—B. t. terrestris, habitat in S. Florida; variation; food, Duellman & Schwartz.—B. t. terrestris, B. t. copei, B. t. americanus, larval characters, distribution; discussion on isolating mechanisms, Gosner & Black (2).—B. t. terrestris, from Grants cave, Florida, Rolman (2).

† Bufo tiheni sp. nov. p. 14, Middle Pliocene, Alachua formation, Haile VI, A (8W ‡ Sec. 24, T95, R18E), near village of Haile Alachua Co., Florida, Sacral vertebra and ilium figd., Aufenberg. Lar med —I Tar

[

& 1

dipe (1). Lin occ the pes effe disc Sze

cha ver & I Fer am gra Of

pro

to

Cla

Lo

& I du & I her fro

Ta

exe

Sp pr Co (pl p. Tr

ch

Pr Ai C. (p 1

ell-

nsis

ria-

tad-

108

nale

ves,

vies

naz-

W.

alin osis.

into

in

rías,

i Az

aach

and

tat:

rigin

the

nto-

Ri.—

tion.

tion eed-

rom

lour,

oto.

rida,

tions

ood-

2).-

ood,

opei,

lack

rida,

cene.

T95.

rida,

Bufo typhonius, habitat at Leticia, Colombia, Goin & Layne.

Bufo valliceps, cranial morphology, Baldauf (1).— Larval characters; hybrid inviability partial isolation mechanism in fowleri × valliceps, Gosner & Black (2).— Habitat and breeding dates in Gómez Farías, Tamaulipas Martin.

Bufo variegatus, distribution, Capurro (1).

Bufo viridis, of Crimea, Chernov.—Diminution in dipeptidase activity by X-ray irradiation, Coromaldi (1).—On occurrence in N. Kizil Kum, Krivosbeev.—Limb buds implanted in B. bufo, Perri (2).—On occurrence in Transcaucasia, Rotter.—Spawning and the environment, Rühmekorf (1).—From N.W. Budapest; habitat, Szabo.—Comparative study of the effect of HCG dissolved in various solvents on the discharge of sperm, Lunenfeld, Rabau, H-Steinberg & Szejnberg.

Bufo woodhousei, of Lyon Co., Kansas; habitat, Clarke, Breukelman & Andrews.—B. w. fowleri, of Long Point, Ontario, Adams & Clark.—Laval characters discussion of isolating mechanisms preventing hybridization with B. t. americanus, Gosner & Black (2).—Bufo woodhousei, of Union Co., Oregon, Perguson, Payne & Storm.—Of S. Dakota, Woodbury.

—B. w. fowleri, on geographic relations of Ozarkian amphibians, Dowling.—B. w. woodhousei, on geographic relations of Ozarkian amphibians, Dowling.—Of Glen Canyon, Tanner.

Nectophrynoides occidentalis from Mt. Nimba. Reproduction and life history descd.; photo., figs. Key to distinguish from N. vivipara and N. tornieri, Guibé & Lamotte (4).—N. occidentalis, thyroid development during metamorphosis studied quantitatively, Lamotte & Prum.—Development of the ovary, Lamotte & Rey.

Pedostibes ruyosus sp. nov., p. 476, from Menuang, headwaters of Baleh R., Sarawak; fig. Distinguished from P. hosi, Inger (2).

Rhinophrynus dorsalis, habitat in Gómez Farías, Tamaulipas, Martin.

#### HYBRIDS

Bufo fowleri  $\times$  B. terrestris, on interspecific gene exchange in S.E. Louisiana, Volpe.—Bufo bufo  $\circ$   $\times$  Rana temporaria  $\circ$ , Hertwig, Weiss & Ziemann.

#### CENTROLENIDAE

Discussion of systematic position and diagnostic characters of the three genera (Centrolene, Teratohyla and Cochranella) recognised; figs., Eaton.—Costa Rican species discussed; distribution notes; Key. Species recognised and described:— Centrolene prosoblepon (photo.): Teratohyla spinosa (photo): Cochranella pulverata, (photo.) C. albomaculata, (photo.), C. granulosa (photo.), C. decorata sp. nov. p. 50, photo., from Hda. La Florencia, 3 mls. W. of Turrialba, Cartago Prov., C. millepunctata sp. nov. p. 53, photo., La Palma, San José Prov., C. fleischmanni, (photo.), C. chrysops (photo.), C. chrripoi sp. nov. p. 59, photo, Cocales Creek, Suretka, Limón Prov., C. reticulata sp. nov. p. 61, photo., near Inter American Institute of Agric., Turrialba, Cartago Prov., C. valerioi, C. talamancae (photo.), C. colymbiphyllum (photo.), Taylor, E. H. (2).

Vor. 96

Centrolene prosoblepon, anatomical study; figs., and photos., Eaton.

#### DENDROBATIDAE

Dendrobates auratus, D. typographicus (photo.), of Costa Rica, Taylor, E. H. (1).—D. granuliferus sp. nov. p. 10, photo., from low mountains N. of Rio Diquis, approx 3 mls. N. of Palmar, Puntarenas Prov., Costa Rica; distinguished from D. typographicus, Taylor, E. H. (1).—D. pictus, from Amazonas, Colombia; colour in life; photo., Goin & Layne.—D. machadoi, sp. nov. p. 73, Serra do Navio, Fed. Terr. Amapa, Brazil, figs.; related to D. trivittatus, Bokermann (2),

Phyllobates femoralis, from Leticia, Colombia; colour in life, Goin & Layne.

## DISCOGLOSSIDAE

Alytes obstetricans boscai and A. cisternasi from Majorca, Baleario Islands, Colom.—A. obstetricans, limb regeneration experiments, Lecamp.—Rearing conditions, Wettstein.—Comparative cytology, Ziemann.

Bombina bombina, diet studies, Lác.—Distribution south of Cracow, variation in coloration, Michalowski.—From N.W. Budapest; habitat, Szabo.—B. variegata, origin of nucleoli in ovocytes, Guyenot.—Trapping and experimental removal from normal territory, Heusser (4).—Distribution south of Cracow, variation in coloration, Michalowski.—Spawning and environmental factors, Rühmekorf (1).—Bombinator pachypus [B. v. pachypus], lens induction, Jacobson.

Discoglossus sp., appearance of glycogen in larval liver tissue, Bellec (2).—D. pictus, pigmentation experimental analysis, Bedate.—From the Balcarie Islands, Colom.—Intraperitoneal injections of reserpine, Liébecq-Hutter & Bacq.—Larval incompatibility on grafting, Rossi.

#### HYLIDAE

Call difference as an isolation mechanism in Florida species of hylid frogs, Blair (3).—E. North American species and subspecies desed., figd. (colour plates) and distinguished, Conant (2).—Maxillary dentition of some hylids, Goin, C. J. (2).

Acris crepitans, population size, movement and behaviour of a population in Texas, Pyburn. — A. gryllus on the call with sonagram p. 42, Blair (3),— Habitat in Lyon Co., Kansas, Clarke, Breukelman & Andrews.—On geographic relations of Ozarkian amphibians, Dowling.—Ecology, Geyer, H. (2).—From Jackson Co., Michigan, Hahn, D. E.—From E. Missouri cave, Myers (3).—A. g. blanchardi, A. g. crepitans, A. g. dorsalis, tooth count, Goin, C. J. (2).—A. g. dorsalis, of S. Florida. Note on food, predators range, pattern variation, Duellman & Schwartz.—Acrodytes Fitzinger, 1843 generic name suppressed, Opinion 520.

Agalychnis dacnicolor, of Colima and W. Mexico, Duellman (1).—A. d. A. helenae, maxillary dentition, Goin, C. J. (2).

Flectonotus goeldii, brooding young, Oeser.

Gastrotheca marsupiata, life history in captivity, Grenard.—G. oviferum, maxillary dentition, Goin, C. J. (2).

[

1

hal

on

fro

Bla

600

Sel

Ke

(1)

lisl

Off

Du

Fa

or Or

Bl

ste

H.

(8) in

Do

ne

th

Ne

sp

Sp

N.

ku

5,

no

re

pe M

h

1

Hyla. Discussion and tabulation of maxillary dentition in various species, Goin, C. J. (2).

Hyla alleni, from Leticia, Colombia; colour and habitat, Goin & Layne.

Hyla alvaradoi, of Costa Rica; note on habitat and colour; photo., Taylor, E. H. (1).

Hyla arborea, from Marseilles, Berner.—Of Crimea, Chernov.—Larval grafts, Rossi.—Spawning habits, Rühmekorf (1).—From N.W. Budapest; habitat, Szabo.—H. a. japonica, from Honshu and Shikoku, Japan, Alexander & Diener.—Hatching glands, Yanai & Takayanagi.

Hyla arenicolor, distribution in the U.S.A., map p. 456, Blair (1).—Of Glen Canyon, Tanner.

Hyla baudini, of Colima and W. Mexico, **Duellman** (1).—Smilisca baudini, habitat and breeding times in Gómez Farías region, Tamaulipas, **Martin**.

Hyla boulengeri, of Costa Rica; photo., Taylor, E. H. (1).

Hyla cinerea, photograph, Anon (3).—Note on response of ♀ to call of ♂, Blair (2).—Call discussed, sonagram p. 42, Blair (3).—Figured with sonagram p. 187, Bogert.—On geographic relations of Ozarkian amphibians, Dowling.—Pattern variation of S. Florida individuals. Note on food, breeding, habitat. Includes H. evittata Miller, Duellman & Schwartz.—Nonbreeding habits compared with H. squirella, Goin, O. B.—On occurrence in saline water, Neill (2).—H. c. cinerea × evittata, from Kent Co., Delaware, Conant (1).

Hyla crucifer, discussion of the call, sonagram p. 36, Blair (3).—Of Cass Co., Minnesota, Brattstrom (1).—On geographic relations of Ozarkian amphibians.—Ecology, Geyer, H. (2).—H. c. crucifer, of E. Canada; zoogeography, Bleakney, J. S.

Hyla dominicensis, from Limbé, Haiti; size; sex dimorphism; colour variation; glandular secretion, Lynn.

Hyla duicensis sp. nov. p. 37, photo., Golfito, Puntarenas Prov., Costa Rica; related to H. eleachroa, Taylor, E. H. (1).

Hyla eleachroa, of Costa Rica; photo. Related to H. dulcensis, Taylor, E. H. (1).

Hyla evittata, considered a synonym of H. cinerea, Duellman & Schwartz.

Hyla eximia, habitat in Gómez Farías region, Tamaulipas, Martin.—H. wrightorum, from Coconino Co., 8,000 ft., Arizona, Williams & Chrapliwy.

Hyla faber, colour photograph p. 149, Stettler.

Hyla femoralis, distribution in the U.S.A., map p. 456, Blair (1).—Call discussed, sonagram p. 34, Blair (3).—Ecology and breeding in S. Florida; food; pattern variation, Duellman & Schwartz.

Hyla foliamorta sp. nov. p. 125, 7 mls. N.W. Miraflores Locks, Canal Zone, photo. Related to H. boulengeri, H. acuminata and H. proboscidia, Fouquette.

Hyla gabbi, of Costa Rica; photo., Taylor, E. H. (1).

Hyla gilleni, of C. Australia; photo., Mertens (4).

Hyla gratiosa, on New Jersey population, Black & Gosner.—Call discussed sonagram p. 36, Blair (3).—Figured with sonagram p. 187, Bogert.—Of S. Florida. Note on pattern variation, ecology and range, Duellman & Schwartz.—Note on arboreal and fossorial habits and variety of call, Neill (3).—From White Co., Tennessee, Rossman (1).

Hyla infrafrenata, of New Guinea; fig., Brongersma,

Hyla japonica, first record from Mongolia, Bannikov.

Hyla lanciformis, from Leticia, Colombia; distinguished from H. albopunctatum and H. spegazzini. Call and habitat descd.; photo., Goin & Layne.

Hyla laynei, from Leticia, Colombia; call, habitat and colour descd.; Goin & Layne.

Hyla legleri sp. nov. p. 33, photos., approx. 15 km. W.S.W. San Isidro del General, San José Prov., Costa Rica; related to H. nigripes, Taylor, E. H. (1).

Hyla leucophyllata, from Leticia, Colombia; note on call, habitat, sex difference, colour, Goin & Layne.

Hyla maxima, probably largest tree frog (5" body length); photograph, Anon (4),—Habitat in Leticia Colombia; call and colour desed., Note on validity of name, Goin & Layne.

Hyla membranacea, from Leticia, Colombia. Call, habitat and colour desed., Goin & Layne.

Hyla m. microcephala, first record from Costa Rica; photo. Distinguished from H. underwoodi and H. phlebodes, Taylor, E. H. (1).

Hyla miotympanum, habitat and range in Gómez Farías region, Tamaulipas, Martin.

Hyla misera, from Leticia, Colombia; note on call, habits, habitat and colour, Goin & Layne.

Hyla monticola, from Costa Rica; redescd. Type lost. Photograph, Taylor, E. H. (1).

Hyla ocularis, call discussed, sonagram p. 40, Blair

Hyla phaeocrypta, call discussed, sonagram p. 34, Blair (3).

Hyla phaeota cyanosticta, of Chiapas; colour in life note, Poglayen & Smith.

Hyla picadoi, of Costa Rica, Taylor, E. H. (1).

Hyla pickeringi, ecology, Geyer, H. (2).

Hyla pseudopuma, of Costa Rica; note on colour variation; photos., Taylor, E. H. (1).

Hyla pulchrilineata, of Haiti; call desed.; colour in life, Lynn.

Hyla punctata, from Leticia, Colombia; colour, call and habitat descd.; photo., Goin & Layne.

Hyla regilla, study of visual pigments during metamorphosis, Crescitelli.—Habitat in Baker Co., Oregon, Ferguson, Payne & Storm.—From Pine Mt. 3954 ft, San Luis Obispo Co., California, Pimentel (2).—Development of sympathetic ganglia, Triplett.—Anatomy and cytology of the hypothalamic neurosecretory system, Wilson, Weinberg & Bern.

Hyla rubella, of C. Australia; photo., Mertens (4).

Hyla rubra, from Leticia, Colombia; colour, call, habits descd., Goin & Layne.

Hyla sarayacuensis, from Lecticia, Colombia. Note on colour, sex dimorphism and habitat; distinguished from leucophyllata, Goin & Layne.

Hyla septentrionalis, call discussed, sonagram p. 42, Blair (3).—Of S. Florida. Note on pattern variation, ecology, food. Larva desed. and figd., Duellman & Schwartz.—Breeding in brackish pools in Florida Keys, Meill (2).

Hyla smithi, of Colima and W. Mexico, Duellman (1).

Hyla spilomma Cope, 1877, specific name as published in the above combination to be placed on Official List of specific names, **Opinion 520**.

Hyla squirella, Kentucky record incorrect, Barbour.—Call discussed, sonagram p. 36, Blair (3).—Of S. Florida. Notes on variation, food, ecology, call, Duellman & Schwartz.—Nonbreeding habits compared with H. cinerea, Goin, O. B..—From Houston Co., Georgia, Goodman.

Hyla staufferi, colour, call and habitat of Gómez Farías, Tamaulipas individuals, **Martin**.

Hyla tibiatrix Laurenti, 1768, specific name suppressed as published in the above combination, **Opinion 520.** 

Hyla versicolor, call discussed, sonagram p. 40, Blair (3).—Discussion on mating call in speciation; geographic variation in call, Blair (4).—Taxonomic status in Oklahoma. Distribution and characters of H. v. versicolor and H. v. chrysoscelis discussed, Bragg, (3).—Of Cass Co., Minnesota, Brattstrom (1).—Habitat in Lyon Co., Kansas, Clarke, Breukelman & Andrews.—On geographic relations of Ozarkian amphibians, Dowling.—Mating behaviour, Littlejohn (2).—H. v. versicolor, of E. Canada; zoogeography, Bleakney, J. S.

Hyla zeteki, of Costa Rica, Taylor, E. H. (1).

Hyla zonata Spix, 1824, specific name published in the above combination is rejected, Opinion 520.

Limnaoedus ocularis, habitat in S. Florida, Duellman & Schwartz.—From Chatahoochee Co., Georgia, Goodman.—On occurrence near brackish ditches, Neill (2).

Nyctimystes generic diagnosis, key to component species, maps, figs. Component species:— avocalis sp. nov. p. 6, E. side Goodenough I., D'Entrecasteaux Group, Papua, fig.: daymani sp. nov. p. 8; fig., N. Slope Mt. Dayman, Papua: fluviatilis sp. nov. p. 11, fig., Bernhard Camp, İdenburg R., Netherlands New Guinea: granti: gularis (humerus 3 figd.): kubori sp. nov. p. 18, fig., Kubor Mts., near Kup, 5,000-8,000 ft., N.E. New Guinea: montana: narinosa sp. nov. p. 26, fig., Mt. Hagen, Wahgi Valley region of N.E. New Guinea, over 8,000 ft.: papua: perimetri sp. nov. p. 31, fig., 830-1170 ft., W. slope of Mt. Riu, Tagula I., Louisiade Archipelago, Papua: persimilis sp. nov. p. 34, fig., N. slope of Mt. Dayman, Papua: rueppelli (transferred from Hyla): semipalmata (foot and leg figd.): humeralis (includes Hyla humeralis Blgr.). The following species are trans-

ferred to Hyla:— amboinensis, darlingtoni, loveridgei, thesaurensis. Pattern of palpebral venation as one of diagnostic characters; figs., Zweifel (2).

Phrynohyas Fitzinger, 1843 (type sp: Rana venulosa Laurenti, 1768) generic name accepted, Opinion 520.—
P. inflata, of Colima and W. Mexico, Duellman (1).—
P. spilomma, of Gómez Farias region, Tamaulipas,
Martin.—P. zonata, maxillary dentition, Goin, C. J.
(2).—From Leticia, Colombia; note on variation from
Trinidad form, call and glandular secretion, Goin &
Layne.

Phyllomedusa bicolor, P. burmeisteri, maxillary dentition, Goin, C. J. (2),—P. bicolor, from Leticia, Colombia; note on colour, habits, song and glandular secretion, Goin & Layne.

Pseudacris brachyphona, P. brimleyi, P. clarki, P. nigrita (and subspecies), P. ornata, P. streckeri, maxillary dentition, Goin, C. J. (2).—P. brimleyi, life history, figs., Gomer & Black (1).—P. clarki, P. nigrita. Habitat in Lyon Co., Kansas, Clarke, Breukelman & Andrews.—P. clarki, P. nigrita, late breeding activity (September) after heavy rainfall in Harris Co., Texas, Kennedy.—P. nigrita, call discussed, sonagram p. 40. Blair (3).—On geographic relations of Ozarkian amphibians, Dowling.—Of Union Co., Oregon, Ferguson, Payne & Storm.—Reproductive behaviour and the importance of vocalization, Martof & Thompson.—P. n. feriarum, ecology, Geyer, H. (2).—P. n. triseriata, of E. Canada; zoogeography, Bleakney, J. S.—From 8,000 ft., Coconino Co., Arizona, Williams & Chrapliwy.—Hormonal influence on sex differentiation, Witschi, Foote & Chang.—P. n. verrucosa, ecology in S. Florida; food-ants, beetles, Duellman & Schwartz.—On occurrence in halophytic grasses, Neill (2).—P. ornata, map of distribution in the U.S.A., p. 455, Blair (1).—Call discussed, sonagram p. 34, Blair (3).—From Fort Benning, Georgia. Locality records for Upper Coastal Plain, Goodman.—P. streckeri, distribution in the U.S.A., map p. 455, Blair (1).—P. streckeri, on geographic relations of Ozarkian amphibians, Dowling.

Pternohyla fodiens, of Colima and W. Mexico, Duellman (1).

Smilisca, see under Hyla,

Sphoenohyla aurantiaca, habitat and colour of Leticia Colombia specimens; distinguished from S. dorisae, Goin & Layne.—S. aurantiaca, S. habra, S. nana, maxillary dentition, Goin, C. J. (2).—S. dorisae, habitat in Leticia, Colombia; colour desed., Goin & Layne.—S. habra, from Leticia, Colombia; habitat, call and colour desed., Goin & Layne,—S. seabrai sp. nov. Serra do Navio, Federal Territory of Amapá, Brazil, p. 43, figs. Related to S. orophila and S. habra, Bokermann (1).

#### Hyprids

 $Hyla\ gratiosa imes cinerea$  figure and sonagram of hybrid, p. 187, Bogert.

#### HYPEROLIIDAE

References at present retained under RHACOPHORI-DAE.

dis-

itat

1

4).

\$ 8

ida.

iell-

rial

Co.,

ma.

km. ov., (1).

ody ticia y of

Call, ica ; and

mez call,

l'ype Blair

34, life

lour ur in

call

gon, i4 ft,

## LEPTODACTYLIDAE

Plates from Philippi's Batraquios Chilenos; discussion of status and synonymy of species, Cei (4).

Batrachyla leptopus, distribution and ecology, Capurro (1).

Calyptocephalella gayi, distribution, Capurro (1).

Orinia. Theory of evolution of the species, Main, Lee & Littlejohn.—C. sloanes, sp. nov., p. 225 from S. bank of Murray R., by main traffic and railway bridge adjacent to Tocumwal on Murray, N.S.W. Grouped with C. insignifera superspecies and sympatric with closely related C. signifera and C. parinsignifera, Littlejohn (1).

Eleutherodactylus. Technique for sectioning ovaries and eggs, Griffiths & Carter.—Eleutherodactylus of Cuba believed to be divisible into five groups:—auriculatus, varleyi, ricordi, dimidiatus, symingioni. E. zeus, sp. nov. p. 38, photo., 0.5 mls. S. of San Vicente Pinar del Rio Province, Cuba; affinities discussed, Schwartz (3).—E. abbotti, of Haiti; colour pattern, Lynn.—E. augusti, E. hidalgoensis, of Gómez Farias region Tamaulipas; habitats and range. Note on stomach contents of hidalgoensis, Martin.—E. a. augusti, from S.E. Coahuila, Webb.—E. augusti fluscofemora, E. a. latrans, from Coahuila; discussion on absence of clinal characters, Webb.—E. augusti, Cuban members of this species group discussed. Recognised:—E. a. auriculatus, E. a. varians, E. sonans, E. eileenae and E. a. olibrus subsp. nov. p. 72, Cueva del Cable, San Vincente, Pinar del Rio Province, Cuba, photo. Distinguished from E. a. varans (photo.) and E. a. auriculatus. E. gehrmanni considered a synonym of E. eileenae, Schwartz (1).—E. dimidiatus amelasma subsp. nov. p. 12, San Vicente, Pinar del Rio Prov. Cuba, (of the ricordi group); E. gehrmanni sp. nov. p. 4, photo., near Guantanamo Bay, Oriente Prov. Cuba, (of the ricordi group); E. phyzelus sp. nov. p. 7, photo., 44 mls. N.W. Vicente, Pinar del Rio Prov. Cuba (of the dimidiatus group); E. phyzelus sp. nov. p. 7, photo., Avanta Schwartz (4).—E. inoptatus, from near Limbé, Haiti; colour, Lynn.—E. monnichorum, first record from Costa Rica; note on variation from typical form; photo., Taylor, E. H. (1).—E. occidentalis, of Colima; distribution in W. Mexico, Duellman (1).—E. pictissimus from near Milot, Haiti; colour, Lynn.—E. schmidti limbensis subsp. nov., p. 155, approx. 3 mls. S.W. of Limbé, Haiti, Lynn.—E. taurus sp. nov., p. 17, photo., Golfito, Puntarenas Prov., Costa Rica; related to E. fleischmanni, Taylor, E. H. (1).—E. vocalie, of Colima; note on variation and distribution in W. Mexico, Duellman (1).—E. protissimus from near Milot, Prov., Cuba. Related to E. gundlachi and E. atkin

Eupsophus bibroni, E. calcaratus, E. coppingeri, E. grayi, E. hidalgoi, E. kriegi, E. maculatus, E. nodosus, E. taeniatus, E. verrucosus, distribution in Chile; ecology, Capurro (1).—E. taeniatus, development, Capurro (2).—Biology and development, figs., Cei & Capurro.

Heleioporus, comparison with Neobatrachus of ability to take up water in response to dehydration,

Bentley, P. J. et al.—Heleioporus, five species recognised:—australiacus, albopunctatus, psammophilus, eyrei and inornatus. Discussion on ecology, distribution and evolution, Main, Lee & Littlejohn.—H. sudelli, considered a synonym of H. pictus. Specimens from St. George, S.W. Queensland. Relationship in pictus-centralis-pelobatoides complex discussed, Hosmer.

Hylella azteca, of Colima; note on habits, a terrestrial hylid breeding in rocky streams and not a bromeliad dweller, **Duellman** (1).

Hylodes hylaeformis Melin, name preoccupied by Cope's Phyllobates hylaeformis New name melini proposed as substitute, **Bokermann** (3).

Hylorina sylvatica, distribution and ecology, Capurro (1).

Kyarranus gen. nov. p. 3; a cycloranine related to Limnodynastes but differs from it and related genera in embryology and in pre-sacral vertebrae. Type species designated, K. sphagnicolus, sp. nov. p. 4, from 5,000 ft. Point Lookout, near Ebor, New South Wales; photos. Genus also includes K. loveridgei (Philoria loveridgei Parker, P. frosti Loveridge), Moore (2).

Lechriodus, larvae made into stew and eaten by natives, Brongersma.

Lepidobatrachus asper, geographic isolation and phenotypic constitution of La Rioja and Grand Chaco populations, Cei (2).

Leptodactylus, isolation of leptodactylina from the skin, Erspamer.—L. hylaedactylus, habitat at Leticia, Colombia; note on call and method for collecting specimens, Goin & Layne.—L. i. intermedius, from Leticia, Colombia; distinguished from podocipinus; Call, habits and colour, Goin & Layne.—L. labialis, L. melanonotus, habitats in Gómez Farias region, Tamaulipas, Martin.—L. melanonotus, distribution in W. Mexico, Duellman (1).—L. ocellatus blood system in the hypophysis, Cruz.—Behaviour; cannibalism. feeding habits, defense, Gallardo (1).—L. p. pentactylus, habitat and habits in Leticia, Colombia, Goin & Layne.—L. podicipinus, from Leticia, Colombia; note on colour, Goin & Layne.—L. prognathus, biology, Gallardo (2).—L. rubido, from Leticia, Colombia, Goin & Layne.

Limnodynastes dorsalis, variations in larval development, Ormsby.

Mixophyes fasciolatus on delayed metamorphosis,

Myobatrachus gouldii, and its associations with termite colonies, Philipp.

Neobatrachus, comparison with Heleioporus of water uptake after dehydration, Bentley, P. J. et al. — N. pictus, N. pelobatoides, N. centralis, N. wilsmorei, distribution, ecology and evolution, Main, Lee & Littlejohn.

Paratelmatobius gen. nov. p. 241, type species P. lutzii sp. nov. p. 241, (photos; figs., osteological characters) from Brejo da Lapa, Alto Itatiaia, state of Rio de Janeiro, Brazil (22°24'S: 44°55'W; 2100 m.). Related to Cyclorhamphus, Telmatobius and Batrachophrynus. P. pictiventris sp. nov. p. 239 from Serra da Bocaina, Parahyba valley, Lutz & Carvalho.

nus;
Pl
note
vibre
and
(1).

[19

PI

sy and phot Nayo notes S. p. S. p. note phot Rela S. cy in

To larve T. if Cape desce (1).

S. m

A pred

Con Sab feed phr. Flo. M. amy Flo. Due

And Ari Gór of G

Tay

mus cun hin Nay Dis

> reg Col

og-

u-

ci-

ed,

<del>08</del>-

by

ini

gy,

to

ra pe

m

th

qei

0),

by

nd

co

he ia.

ng

18;

is, on,

in em

m.

taia,

mus,

ia,

p.

sis,

ith

ter

rei,

ies

cal

of

1.).

ra-

Philoria frosti, P. loveridgei transferred to Kyarranus; figs., Moore (2).

Pleurodema bibroni, P. bufonina, distribution; notes, Capurro (1). — P. bibroni, sexual warning vibration, Cei & Aguilera.—Population variability and phenotypic distribution; statistical analysis, Cei (1).

Syrrhopus, key to forms in W. Mexico, descriptions and figures. S. m. modestus (colour, call, range, photo.): S. m. pallidus subsp. nov. p. 5, San Blas, Nayarit, photo.: S. rubrimaculatus, range: S. leprus, notes on relationships, colour, call and range; photo.: S. p. pipilans, note on colour, call and range; photo.: S. p. nebulosus, characters and range: S. interovitalis, note on colour and range: S. teretistes sp. nov. p. 10, photo., 3 mls. N.W. Tepic, Nayarit (±2950 ft.), Relationships of species discussed, Duellman (2).—S. cystignathodoides, S. latodactylus, S. smithi, habitats in Gómez Farias region, Tamaulipas, Martin.—S. modestus, of Colima; call desed.; note on habitat and distribution, Duellman (1).

Telmatobius hauthali schreiteri, somatic variation in larvae desed.; dwarf form desed., figs., Pisanó.—
T. laevis, T. marmoratus, T. halli, T. montanus,
T. peruvianus, distribution and ecology in Chile,
Capurro (1).—T. bullocki, of Chile, Capurro (1).

Tomodactylus petersi, of Colima and Tecolapa; call desed.; note on distribution in W. Mexico, Duellman (1).

#### MICROHYLIDAE

Asterophrys turpicula, of New Guinea; note on predation on large lizards, Brongersma, L. D.

Breviceps adopersus, from Inyanga district, S. Rhodesia, Fitzsimons.

Gastrophryne carolinensis, From E. Chesapeake Bay, Conant (1).—†Microhyla carolinensis, from Pleistocene Saber-tooth cave, Florida, Holman (1).—On ant feeding behaviour, Holman & Campbell.—Gastrophryne carolinensis, on occurrence on Salicornia flat in Florida and in other brackish habitats, Neill (2).—M.c. carolinensis, on geographic relations of Ozarkian amphibians, Dowling.—G. c. carolinensis, of S. Florida. Notes on variation, call, ecology and food, Duellman & Schwartz.—M. carolinensis olivacea, habitat in Lyon Co., Kansas, Clarke, Breukelman & Andrews.—M.c. mazutlanensis, from Santa Cruz Co., Arizona, Williams & Chrapliwy.—G. olivacea, of Gómez Farías region, Tamaulipas, Martin.—G. u. usta, of Colima, Duellman (1).

Glyphoglossus molossus, from Thailand, photo., Taylor & Elbel.

Hypopachus. Key to Mexican species:— caprimimus, maculatus, alboventer alboventer, cuneus cuneus, cuneus nigroreticulatus, alboventer reticulatus, oxyrrhinus taylori subsp. nov. p. 89, 3 mls. E. San Blas, Nayarit, fig., oxyrrhinus ovis, oxyrrhinus oxyrrhinus, Discussion on relationships and taxonomic status. Shannon & Humphrey.—H. cuneus, of Gómez Farías region, Tamaulipas, Martin.—H. oxyrrhinus, of Colima; distribution in W. Mexico, Duellman (1).

Kaloula borealis, origin, development and changes of the gills and lungs, Giang & Li.—K. picta, confirmed on Negros Island, Alcala (2).—K. p. pulchra, from Thailand, Taylor & Elbel.—K. p. taprobanica, tadpoles first description from Ceylon, Kirtisinghe.

Microhyla berdmorei, from Thailand, photo.; description, Taylor & Elbel.—M. carolinensis, see under Gastrophryne carolinensis.

Ramanella obscura, eggs, tadpoles, metamorphosis and ecology descd., photos., Note on dermal secretion, Morgan-Davies (2), -R. palmata, tadpole first description from Coylon, Kirtisinghe.

Sphenophryne, technique for sectioning ovaries and developing eggs, Griffiths & Carter.

Uperodon systoma, note on demal secretion, Morgan-Davies (2).—Note on burrowing habits; photo., Morgan-Davies (3).

#### PELOBATIDAE

Megophrys aceras, M. hasselti hasselti (and photo.), from Thailand, Taylor & Elbel.

Pelobates discussion of evolutionary tendencies and phylogeny on basis of skull study. Pseudopelobates gen. nov. p. 1037, type species P. transcaucasious (Delwig 1928). Skeletal characters of Pelobates fuscus, P. syriacus (and subspecies), P. cultripes and P. varaldi compared, Pasteur (2).—P. cultripes, from Marseilles; littoral, Berner.—Total sex reversal using oestradiol on larvae, Gallien & Collenot.—Mass numbers during wet weather at Canet (Pyr-Or), Petit & Lomont.—P. fuscus, of Crimea, Chernov.—Spawning habits, Rühmekorf (1).—From N.W. Budapest; habitat, Szaho.

Pelodytes caucasica, of Transcaucasia, Rotter.—
P. punctatus, from Marseilles; habitat, Berner.—
Production of intersex by cestradiol treatments,
Gallien & Collenot.—Present in Luxembourg, Hoffmann.

#### PIPIDAE

Pipa, from Venezuela. Species desed.:— P. pipa, P. parva, key to distinguish, distribution map, notes on habitats, reproduction; photos., Gines.

Xenopus, effects of cytotoxic substances on embryos, Waddington.—X. laevis, melanophores, Bagnara (2).
—Early gastrula transplants, Balinsky.—Effect of penicillin, streptomycin and penicillium mycelium on growth and development, Ber & Zieleniewski.—
β-glucuronidase activity during early development of embryos, Billett (1).—Temperature effects on the electrocardiogram, Biörck & Johansson.—Studies on germ cells in eggs, embryos and larvae, Blackler.—Evocators and their diffusibility, Brahma.—Reaction of melanophores to hormones etc., Burgers.—Melanophore dispersing reaction as a pregnancy test, Burgers & Mighorst.—Disaggregation of gastrulae; and suggestion of a ribonucleoprotein on the cell surface, Curtis.—Visual mechanism and sensitivity, Dartnall.—Embryos, regional differences in catheptic activity, Deuchar.—Spectral absorption of the photopigment, Dobrowolski, Johnson & Tansley.—On a mutation reducing nucleolar number, Elsdale, Fischberg & Smith.—Nuclear transplantation, Fischberg, Gurdon

& Elsdale,—Influence of temperature on development, Gerbes.—Transplantation of somatic nuclei producing sexually mature individuals, Gurdon, Elsdale & Fischberg.—Supernumerary right arm, photo., Hobson.—Dorsal root ganglia and ventral horn, Hughes & Tschumi. — Forebrain regeneration in post metamorphic forms, Jordan.—Morphological study of labyrinth and middle ear, Kahmann & Frettiöh.—Regeneration power in postmetamorphic animals, Konieczna-Marczynska & Skowron-Cendrzak.—Tadpoles, influence of the olfactory nerves on end brain regeneration, Kosciuszko.—Tadpoles, multiple regeneration of tails, Kucias.—Amino acid composition of skin-gelatin, Leach,—Influence of temperature on pulse rate, Morawska & Zaleska.—Blood sugar level relative to melanophore-expansion, Munday.—Response of lateralis organs to electrical stimulation by direct current, Murray.—X. laevis, thyroid activity and cylo-differentiation of anterior pituitary, Saxén.—Induction of triploidy, Smith, S.—X. tropicalis, from Mt. Nimba; photo. Distinguished from X. fraseri; ranges. Larva desed. and figd., Guibé & Lamotte (4).—From Harbel, Liberia, Taylor & Weyer.

## RANIDAE

E. North American species and subspecies descd., figd. (colour plates) and distinguished; distribution maps, Conant (2).

Arthroleptis crusculum, on morphology and direct development; nest below ground; figs., Guibé & Lamotte (2).

Astylosternus occidentalis, from Mt. Nimba; phoso. Note on habitat; larvae desed., Guibé & Lamotte (4).

Cacosternum boettgeri, early gastrula transplants,

Cornufer guentheri, first record from Negros Island, Alcala (2).

Hemisus guttatum, from Natal and Hluhluwe, Wager (2).—H. marmoratum, breeding habits and general ecology, Wager (2).—H. marmoratum guineensis, from Mt. Nimba; habitat; photo.; larva desed. and figd., Guibé & Lamotte (4).

Micrixalus magnapustulosus sp. nov. p. 1066, photo., Ban Na Phua, Kan Luang, Na Kae, Nakhon Phanom, Thailand, 200 m. approx., **Taylor & Elbel**.

Nannophrys ceylonensis ceylonensis, tadpole first description, Kirtisinghe.

Ooeidozyga lima, from Thailand; note on characters and habits, Taylor & Elbel.

Petropedetes natator, from Mt. Nimba; photo. Distinguished from P. palmipes. Note on habitat and range. Larva descd. and figd., Guibé & Lamotte (4).

Phrynobatrachus francisci, larval development; figs., Lamotte & Dzieduszycka.—P. maculiventris sp. nov., pond in forest, near Doromou, French Guinea, p. 257, fig., Guibé & Lamotte (3).—P. natalensis, early gastrula transplants of presumptive ectoderm, Balinsky.—P. ukingensis mababiensis, from Nyamaziwa, S. Rhodesia, Fitzsimons.

Phrynoglossus laevis martensi, from Thailand; characters, Taylor & Elbel.

Ptychadena of the Cameroons. Key to and description and figures of the species:— longirostris, aequiplicata, oxyrhynchus, superciliaris, mascareniensis, taenioseslis, bibroni, perreti sp. nov. p. 1456, Foulassi, Sangmelina, Fr. Cameroun (related to P. grandisonae), Guibé & Lamotte (1).—P. longirostris, from Uele, Belgian Congo, Laurent (2). For further references see under Rana.

Pyxicephalus delalandii and P. adspersus, presumptive ectoderm transplants, Balinsky.

Rana (Ptychadena) aequiplicata, from Mt. Nimba; habitat forest; photo., Guibé & Lamotte (4).—See also under Ptychadena.

Rana (Conraua) alleni, habitat in Mt. Nimba region. Larva desed. and figd. Photo., Guibé & Lamotte (4),

Rana angolensis [R. fuscigula angolensis] from Inyanga district, S. Rhodesia, Fitzsimons.

Rana areolata, habitat in Lyon Co., Kansas, Clarke, Breukelman & Andrews.—On geographic relations of Ozarkian amphibians, Dowling.—R. areolata aesophus, ecology in S. Florida, Duellman & Schwartz.

Rana arvalis, general notes; photo., Duellman.—Spawning and ecology, Rühmekorf (1).—Action of light, Rühmekorf (3).—Adenoma of the skin with cytoplasmic or nucular inclusions, Stolk (1).

Rana aurora, development of sympathetic ganglia, Triplett.

Rana (Ptychadena) bibroni from Mt. Nimba; photo, Note on characters, affinities and range, Guibé & Lamotte (4).—See also under Ptychadena.

Rana breviceps, note on burrowing and nocturnal habits, Morgan-Davies (3).

 $Rana\ capito$ , from Saber tooth cave, Florida, **Holman** (2).

Rana catesbeiana, of Long Point, Ontario, Adams & Clark.—Irradiation of blood forming cells in the tadpole kidney, Allen.—Experimental control of guanophores, Bagnara (1).—of E. Canada; zoo-geography, Bleakney, J. S.—In British Columbia, figured p. 62, Carl & Guignet.—Retrograde reaction in motor neuron dendrites, Cerf & Chacko.—Fluid and electrolyte movement across the intestinal wall, Chalfin, Cooperstein & Hogben.—Habitat in Lyon Co., Kansas, Clarke, Breukelman & Andrews.—Stomach contents; growth rates, Cohen & Howard.—On geographic relations of Ozarkian amphibians, Dowling.—From Baker Co., Oregon, Ferguson, Payne & Storm.—Ovarian egg nucleic acids, Finamore & Volkin.—Larval gonads maintained on synthetic media, Foote & Foote (2).—In vitro cultivation of gonads, Foote & Foote (3).—From Jackson Co., Michigan, Hahn, D. E.—From Grants cave, Florida, Holman (2).—Blood oxygen capacity, Leftwich.—Regeneration experiments, Schmidt, A. J.—Tadpoles histological studies of corium, Shimozawa.

Rana chalconota, from Thailand, Taylor & Elbel.

Rana chensinensis, structure of the oesophagus; figs., Eksaeva.

Role & C. J. S. Britis Polytions through through through through through the conference of the confere

[19

R dime

R. c

Dav

Mali

\_E

rege

R

Frodege fem Exp gen data

Fro Not Gui

Elb

Nos nan egg Tri as : lack Cor pho (2).

An res abe Car and Hi Mc

he Se Vi Rana clamitans, from Long Point, Ontario, Adams & Clark.—Of E. Canada; 200geography, Bleakney, J. S.—Of Cass Co., Minnesota, Bratistrom (1).—In British Columbia, figured p. 61, Carl & Guiguet.—Polydactylous juvenile, Cooper.—On geographic relations of Ozarkian amphibians, Dowling.—Water flow through the mucosal layer of the stomach, Durbin, Frank & Solomon.—Maintenance of gonads from second-year larvae, in different media, Foote, C. L. & F. M. (1).—On winter activities and habitats of larvae, Getz.—Injected with 1<sup>131</sup>, Kaye & Le Bourhis.—Influence of various sound frequencies on respiratory movements, Kleerekoper & Sibabin.—Blood oxygen capacity, Leftwich.—Regeneration after amputation, Malinin & Deck.—From E. Missouri cave, Myers (3).—Effects of adrenal transplants upon fore limbregeneration, Schotté & Wilber.

Rana corrugata, note on habits and habitat, Morgan-Davies (3).

Rana cubitalis, from Thailand; photo.; note on sex dimorphism and sex characters, Taylor & Elbel.

Rana cyanophlyctis, induced ovulation with mammalian hormones, Ramaswami & Lakshman. — R. c. cyanophlyctis, habits and call; photo., Morgan-Davies (3).

Rana dalmatina, sex differentiation, Ancona.—From Marseilles, Berner.—Combination of cells of degenerating pronephros in the formation of the female genital structures, Cambar & Dechambe.—Experiments on the development of the female genital opening, Dechambe.—Occurrence and general data from Sweden, König.—Spawning and ecology, Rühmekorf (1).—Tadpoles differentiation after treatment with thyroxine and methyl uracil, Stefani.—From N.W. Budspest; habitat, Szabo.

Rana (Dicroglossus) occipitalis, from Mt. Nimba. Note on habitat, coloration, larva (figured); photo., Guibé & Lamotte (4).

Rana erythraea, from Thailand; photo., Taylor & Elbel.

Rana esculenta, dorsal skin potentials, Aguggini & Noseda.—Sex differentiation, Ancona.—From stagnant waters, Marseilles, Berner.—Study of germ cells; eggs, embryos and larvae, Blackler.—Transplants with Triturus, Bondi.—Melanophore dispersing reaction as a pregnancy test, Burgers & Mighorst.—Oxygen lack and phosphorylation of glycogen in muscles, Cordier & Duveau (1).—Effect of oxygen lack on phosphorylation of liver glycogen, Cordier & Duveau & Brandon.—Diminution of dipeptidase activity by X-ray irradiation, Coromaldi (1).—Comparison with Ambystoma mexicanum of the ontogeny of respiratory vascularization, Czopek.—Visual purple absorption, Denton & Wyllie.—Ecology from the Carpathians of Slovakia, Dudich.—Rohon-Beard cells and their behaviour during development, Filippio.—Histotopographical study of the kidney, Geyer, G.—Mortality after insecticide dust on the banks of Lake Trasimeno, Gianotti.—Lens production in the absence of the retina, Jacobson.—Micro electrode studies of cerebellar electrical activity, Joynt.—Rhythmical heart potentials in the ventricle wall, Lacquet.—Secondary cerebral modifications, Legait & Legait.—Variations in respiratory quotient of the skin, Locker.

—Temperature influence on relation of liver tissue respiration to glycogen content, Locker & Doneff.—Origin and development of the pronephres and Wolffian duct, Manelli (1).—Inhibition of tadpole metamorphosis using streptomycin, Marco.—Influence of temperature on pulse rate, Morawska & Zaleska.—Electro retinograph study of different fields of perception, Pilz, Sickel & Birke.—Cutaneous vasomotor reflexes and respiration, Poczopko.—Variations in gut motivity throughout the year, Pora & Rejep.—Polydactyly, Rostand (1).—Spawning and ecology, Rühmekorf (1).—Action of light, Rühmekorf (3).—Biosynthesis of phosphoprotein, Sampath Kumar et al.—Grafting experiments with Triturus, Spirito & Bondi.—Action potential dependence on temperature and metabolism in medullated nerves, Stieve.—Habitat in N.W. Budapest, Szabo.—Diet and feeding habits, Tyler.—Cannibalism of adults on young, Vandevelde.—Orientated gastrula section, Wessing & Claes.—Spinal reflexes, Wittenberger & Fabian.—Comparative cytological studies, Ziemann.

Rana f. fasciata, from Nyamaziwa, S. Rhodesia, Fitzsimons.

Rana fusca, determinations of bilateral embryonic symmetry, Levtrup.—Adenoma of the skin, Stolk (1). —Early development, Veit.—Comparative cytological studies, Ziemann.

Rana goliath, general note on habits and habitat; photo., Perret (3).

Rana g. grayii, from Inyanga district, S. Rhodesia, Fitzsimons.

Rana grylio, of S. Florida; call; food; ecology, Duellman & Schwartz.

Rana hexadactyla, habitat in Ceylon; colour, Morgan-Davies (3).

Rana (Hylarana) albolabris, from Mt. Nimba; photo.; range; sex dimorphism; habitat; larva desed. and figd. Distinguished from R. zenkeri, Guibé & Lamotte (4).

Rana (Hylarana) zenkeri, from Mt. Nimba primary forest; photo. Distinguished from albolabris, Guibé & Lamotte (4).

Rana ishikawae, of Amami group, Loo Choo Is.; photos., Koba.

Rana japonica, from Honshu, Japan, Alexander & Diener.—Biology, Geyer, H. (1).—Sex reversal in thiourea-treated tadpoles, Iwasawa (2).—Cartilage tissue staining using indirect peroxidase reaction, Kawasaki.—Effect of hormones on metamorphosis, Kobayashi.

Rana kuhlii, from Thailand; photo., Taylor & Elbel.

Rana limnocharis, of Amami group, Loo Choo Is., Koba.—Effect of colchicine on early development, Su & Tsai.—R. l. limnocharis, R. l. greenii, of Ceylon; thinguished. Photo. of suspected hybrid, Morgan-Davies (3).—R. l. limnocharis, from Thailand; photo.; note on chest fold and sex characters, Taylor & Elbel.

Rana (Ptychadena) longirostris, from humid forest, Mt. Nimba; related to R. christyi; photo., Guibé & Lamotte (4).—See also under Ptychadena.

ae), ele, ces

1

id:

ip-

sis, 881.

oa;

(4).

ke, s of

of with

oto.

mal nan

the of soobia, tion and vall,

Co., ach -On ing. orm.

lood beridies

us ;

Rana (Ptychadena) maccarthyensis, from Mt. Nimba, photo. Larva desed. and figd., Guibé & Lamotte (4).

—R. maccarthyensis, from Harbel, Liberia, Taylor & Wever.

Rana macrodon, eggs lack pigment; considered distinct from R. magna, Inger (3).—From Thailand, Taylor & Eibel.

Rana macroglossa, larvae descd. and figd.; note on status and relationship with R. palmipes, Volpe & Harvey.

Rana m. macrognatha, from Thailand; photo.; diagnosis and description, Taylor & Elbel.

Rana magna, considered a polytypic species and distinct from R. macrodon; subspecies:— acanthi, macrocephala, visayanus. Eggs pigmented, Inger (3)

Rana m. mascareniensis, from Inyanga district, S. Rhodesia, Fitzsimons.—R. (Psychadena) mascareniensis, from Mt. Nimba; photo. Note on larval characters, Guibé & Lamotte (4).—See also under Psychadena,

Rana modesta, considered distinct from R. magna; notes on affinity and pattern, Inger (3).

Rana narina, of Amami group, Loo Choo. Is., Koba.

Rana nigromaculata, study of fluorescent "Ranachrome I", Goto & Hama,—Flavines and pterines in the eye, Hama & Horiuchi,—Cartilage tissue staining using indirect peroxidase reaction, Kawasaki.—Experiments on the neurosecretory system, Kumamoto. — Study of breeding habits in Nanchang suburbs, Ling, K.—Cortical granules in eggs, Osanai.—Conduction velocity in intracutaneous nerve fibres, Takagi.—Neuro-muscular depression, Takeuchi, A.—Effect of temperature on the neuromuscular junction, Takeuchi, N.—Muscle fibres used in determining the role of intracellular phosphate ions, Umezawa.—R. n. nigromaculata, from Honshu, Japan, Alexander & Diener.—Donors of pituitary extract to Hyla arbora japonica, Yanai & Takayanagi.

Rana nigrovittata, from Thailand; photo., Taylor & Elbel.

Rana obbiana, desed.; colour variation, call, habits and habitat; photos., Cherchi (1).

Rana (Ptychadena) ozyrhynchus habitat in Mt. Nimba region; photo. Larva desed. and figd., Guibé & Lamotte (4).—See also under Ptychadena.

Rana palmipes, from Leticia, Colombia; colour and habits; photo., Goin & Layne.—Hybridization and larval development; relationship with R. macroglossa discussed, Volpe & Harvey.

Rana palustris, of E. Canada; zoogeography, Bleakney, J. S.—From Mississippi caves, Brode.—From cave in Mississippi, Brode & Gunter.—On geographic relations of Ozarkian amphibians, Dowling.—From E. Missouri cave, Myers (3).

Rana pipiens, of Long Point, Ontario, Adams & Clark.—Five legged Ohio specimen, Adler (2).—
Embryos treated with solutions of ATP, AMP-3, ADP, IDP or mixtures of AMP-3+ATP, Ambellan.—
Observations on spinal ganglion cells, Anderson & Breemen.—Experiments on pattern variants, Anderson & Volpe.—Ionic movements and activity in frog

nerves, Asano & Hurlbut.-Control of guanophores. Bagnara (1).—Hormonal control of pigmentation, Bagnara & Neidleman.—Enzymes and their effect on eggs, Barch & Shaver.—Gastrulae differentiation, Barth & Barth.—Of E. Canada; zoogeography, Bleakney, J. S.—Of Cass Co., Minnesota, Brattstrom (1).—Effect of gonadotrophins on testes morphology and thumb pads, Burgos & Ladman.—Cortisone effect on ovulation, Chang & Witschi.—Habitat in Lyon Co., Kansas, Clarke, Breukelman & Andrews.—Depolarization in activating frog twitch muscles, Csapo & Suznki.—On geographic relations of Ozarkian amphibians, Dowling.—Of Colima and W. Mexico, Duellman (1).-Variation in S. Florida individuals. ecology, food and adaptation to brackish water, Duellman & Schwarts.—Water flow through the gastric mucosa, Durbin, Frank & Solomon.—Development of collagen in the embryo, Edds.—Comparative nucleic acid contents in liver, ova and spermatozoa nuclei, England & Mayer.—Contaminent in egg extracts, Finamore.—Glycogen content of eggs, Fitch Merrick.-Antigen localisation and induction, Flickinger.—Gradients across vitelline membrane in early stage eggs, Flickinger & Blount.—Tadpoles in hind limb regeneration experiments, Goss.—Sulphony-lurea structure-activity relationships, Gourley.—Chloride transport in the gastric mucosa, Heinz & Durbin,—† Rana cf. pipiens from Saber-tooth cave, Florida, Holman (1).—R. pipiens, rehydration conreaction of frozen dried muscle, Hopkins,—Na Cl transport and O<sub>2</sub> consumption in the skin, Huf, Doss & Wills.—Effect of azide and chloretone on Na and K contents and respiration of sciatic nerves, Hurlbut.-Eserine-acetylcholine interaction in the rectus abdominis muscles, Jenden.—Potential relations in isolated sartorius muscle, Jenerick.—Cartilage tissue staining using the indirect peroxidase reaction, Kawasaki.-Influence of various sound frequencies on respiratory movements, galvanic skin reflex and gastric hunger movements, Kleerekoper & Sibabin. Enzyme inhibitors on resting potential and ion distribution in the sartorius, Kloot.—Spinal ganglion cells in sodium-free media, Koketsu, Cerf & Nishi.—Electron microscope study of lampbrush chromosomes, La fontaine & Ris.—Blood oxygen capacity, Leftwich.—Development of trunk muscles, Liedke.—Oxygen poisoning of eggs, Malamed (2).—Regeneration after amputation, Malinin & Deck.—Altudinal distribution, range and habitat in Gómez Farías region, Tamaulipas, Martin. Tadpole metamorphosis, Money, Meltzer, Young & Rawson.-Interrelations between nuclei and cytoplasms of R. sylvation and pipiens, Moore (1). Transfer of haploid nuclei with R. sylvatica, Moore (3). —From E. Missouri cave, Myers (3).—Increased partial pressure of O<sub>2</sub> accelerates development in embryos, Nelsen.-Measurements from disarticulated skeletons, Olson & Miller.—Transection of the spinal cord, Piatt & Piatt .-- Anatomy of sympathetic rami, effects of total and partial sympathectomy, Pick .-Tadpoles and growth under crowded conditions, Richards.—Studies on ovulation, Rondell & Wright. Histochemistry of oocyte cortical region, Rosenbaum .- RNA and protein in neural induction, Rounds & Flickinger.—Regeneration experiments, Schmidt, A. J.—Effects of adrenal transplants upon forelimb regeneration, Schotté & Wilber.-Adenocarcinoma of the kidney with cytoplasmic or nuclear inclusions, Stolk (1).—Relation between liver weight and degree of restoration after partial hepatectomy, Streett. -Transplantations of haploid nuclei, Subtelny. -Ovulation experiments, Taurand.—Effect of pituitary extracts from marine and freshwater teleosts on water balance, Weisel.—Water exchange and dorsal lymph drainage as affected by Pitressin, Wilson & Archéeacon.—Resistance and susceptibility to 0<sub>2</sub>. Wittner & Rosenbaum.—R. pipiens brachycephala, from Baker Co., Oregon, Ferguson, Payne & Storm.—Of Glen Canyon, Tanner.—R. pipiens sphenocephala, Saber tooth cave, Florids, Holman (2).—On saltwater association in Florids. Neill (2).

Rana pretiosa, life history, Turner.—R. pretiosa luteiventris, from Baker Co., Oregon, Ferguson, Payne & Storm.

Rana pustulosa, of Colima and W. Plateau, Duellman (1).

Rana (Ptychadena) retropunctata, from Mt. Nimba. Distinguished from other Ptychadena; photo., Guibé & Lamotte (4).

Rana ridibunda, of Crimea, Chernov.—Ecology from the Carpathians of Slovakia, Dudich.—Influence of methylthiouracil on the blood, Grabie-Broda.—From Transcaucasia, Rotter.—Habitat in N.W. Budapest, Szabo.

Rana rugosa, distribution of hatching glands, Yanai & Tokayanagi.

Rana septentrionalis. of E. Canada; zoogeography, Bleakney, J. S.—Of Cass Co., Minnesota, Brattstrom (1).

Rana sinaloae, from 1,500 ft., Colima; occurs also on W. Plateau, **Duellman** (1).

Rana subaspera, of Amami group, Loo Choo Is.; photo., Koba.

Rana (Ptychadena) submascareniensis, from Mt. Nimba; photo. Larvae desed. and figd., Guibé & Lamotte (4).

Rana (Ptychadena) supercitiaris, from Mt. Nimba; range; photo., Guibé & Lamotte (4).—See also under Ptychadena.

Rana sylvatica, hormonal control of pigmentation, Bagnara & Neidleman.—Experiments on tadpoles on control of guanophores, Bagnara (1).—Of E. Canada; zoogeography, Bleakney, J. S.—Cannibalism in tadpoles, Bleakney, S.—Habitat in Lyon Co., Kansan, Clarke, Breukelman & Andrews.—On geographic relations of Ozarkian amphibians, Dowling.—Role of thyroid in kidney development; development and degeneration of pronephros, Hurley.—Interrelations between nuclei and cytoplasms of R. pipiens, Moore (1).—Transfer of haploid nuclei with R. pipiens, Moore (1).—Transfer of haploid nuclei with R. pipiens, Moore (1).—Transfer of haploid nuclei with R. pipiens, Rogeneration experiments, Schmidt, A. J.—Northsouth gradient of body/tibia ratio, R. s. cantabrigensis and R. s. sylvatica distinguishable at their extreme ranges but not in Wisconsin, Suzuki.

Rana temporaria, thermal stadia in development, Bebak.—Of Marseilles, Berner.—Studies of eggs, embryos and larvae giving germ cell information Blackler.—Myocardium in the absence of potassium and influenced by acetyl choline, Borghgraef & Carmeliet.—Passive electrical properties of retins, choroid and sclera for radial fields and currents, Brindley (1).—Retinal responses to light, Brindley (3).

-Dark adapted frogs study of photo-sensitive pigments in the pink and green rods, Denton & Wyllie. Fast and slow muscle fibres, Gray.—Origin of nucleoli in ovocytes, Guyénot.—K tracer in muscle for the study of permeation of ions, Harris.—Tadpoles, effects of crowding, Hodler .- Diastolic and systolic potentials in heart muscle, Lacquet.-Intra peritoneal injections of reserpine, Liébecq-Hutter & Bacq.—Direct absorp-tion of dissolved Strontium-90 and Yttrium-90, Lucas & Pickering.—Spawning, Mertens (2).—On temperature adaptation of cell enzymes, Mews.—Influence of temperature on pulse rate, Morawska & Zaleska.— Regeneration experiments, Newth (1).—Mechanism of the initial stages of spermiation, Oordt.—Spermatogenetic cycle; regulation by temperature, Oordt.-Ultrastructure of Schmidt-Lauterman clefts an related shearing defects of the myelin sheath, Robert-son.—Action of tetraiododesaminothyronine on tadpole metamorphosis, Roth (1), -Action of thyroxamine on metamorphosing tadpoles, Roth (2).—Spawning and ecology, Rühmekorf (1).—Action of light, Rühmekorf (3).—Larval temperature selection, Rühmekorf (2).—Action of sodium floride on metamorphical selection of sodium floride on metamorphical selection. phosis, Saadi-Nazim.—In marshy areas of Dernford Fen, Sharp.—Blood pressure and pressure pulses in arterial arches, Simons.—Ovulation experiments, Taurand.—Tadpoles their electromotive forces and their significance, Viaud & Dreyfus.—Control of gonadotroph activity, Vivien & Schott.—Occytes concentrations of iodine and thyroid hormones, Volpent, Michel & Roche.—Acetylcholine contracture of muscle, Zeleny & Kozák.—R. t. temporaria, effects of tryptaflavin on mitosis in tadpoles, Meyer. arterial arches, Simons .- Ovulation experiments. Spawning site observations, Warwick.

Rana tigrina, cytoplasmic inclusions in ageing neurones, Malhotra.—Lipids in cogenesis, Nath, Gupta & Kaur.—R. tigrina crassa, on cannibalistic habits, Morgan-Davies (3).—R. tigrina pantherina, from Thailand, Taylor & Elbel.

Rana (Ptychadena) tournieri, from Mt. Nimba; range; photo., Guibé & Lamotte (4).

Rana venulosa Laurenti, 1768, specific name as published in the above combination to be placed on the Official Index of Specific Names, (specific name of type sp. of Phrynohyas Fitzinger, 1843), Opinion 520.

Rana virgatipes, on occurrence in Maryland and Delaware, Conant (1).—From Worcester Co., Maryland, Reed (2).

Rana viridis, reactions caused by stimulation of the horizontal semi-circular canal, Gribenski.

Trichobatrachus robustus, importance of hair as a possible means of discrimination by the sexes, Mertens. (1).—Habits and habitat and note on function of "hairs"; photo., Perret (3).

#### HYBRIDS

Rana esculenta  $\mathcal{Q} \times Bufo\ viridis\ \mathcal{J}$ , Hertwig, Weiss & Ziemann.

Rana pipiens  $\mathcal{Q} \times R$ . sylvatica  $\mathcal{S}$ , Hertwig, Weiss & Ziemann.

Rana pipiens × R. sylvatica, transfer of haploid nuclei, Moore (3).

phiman on ter,

the

97

T08,

ion,

t on

ion,

ohy,

rom

logy

fect

Co.,

lar-

doptive ozoa egg itch tion, e in ony-

ave, concanss & d K

ated ining ki. tory inger zyme on in liumnicrotaine

reloping of ation, and artin. ng & cyto(1). re (3). coased

nt in dated spinal rami, ck. tions, right.

ounds midt, elimb noma sions, legree ett.—

nitary

[

Bo

So

Ari

mo

of

Ple

bel

S. i

sho

## RHACOPHORIDAE

Afrizalus dorsalis, A. fulvovittatus fulvovittatus, from Mt. Nimba; photos.; habitats; larvae desed. and figd. A. wiedholzi, from Mt. Nimba; note on habitat and range; photo., Guibé & Lamotte (4).—A. dorsalis regularis, from Boma Terr., Lower Congo, Laurent (2).

Chiromantis petersi kelleri, breeding and development; photos., Cherchi (2).—C. rufescens, from Mt. Nimba; photo. Arboreal form from humid forest. Larva desed. and figd., Guibé & Lamotte (4).—C. rufescens, C. xerampelina, on nesting habits, figs., Verbaak.

Hylambates (Phlyctimantis) leonardi, from Mt. Nimba; photo.; habitat; larva desed. and figd., Guibé & Lamotte (4).

Hyperolius, Key to forms occurring in Mt. Nimba region. Descriptions, photographs, variation, Lauregion. Descriptions, photographs, variation, Laurent (1).—H. acutivostris, photo., Laurent (1).—[H.] Rappia aylmeri, E. G. Boulenger—H. fusciventris, Laurent (1).—H. baumanni—H. picturatus, Laurent (1).—H. chlorosteus, of French Guinea. May include H. pleurotaenius (non Boulenger), Laurent (1).—H. c. concolor, of Mt. Nimba region. Includes guineensis, depressus and togoensis. Compared with H. picturatus; photo., Laurent (1).—H. depressus—H. c. concolor, Laurent (1).—H. festivus—H. picturatus, Laurent (1).—H. fuscivus—H. Mt. Nimba. Incarrent (1).—H. fuscivus—H. f Laurent (1).—H. f. fusciventries, of Mt. Nimba. Includes Rappia aylmeri, H. oeseri, H. trifasciatus. Discussion on affinities with H. burtoni. Photo., Laurent (1).—H. granulatus, of Mt. Nimba; discussion of the contraction cussion on relationship with oxyrhynchus, Laurent (1).

—H. guineensis = H. c. concolor, Laurent (1).

H. guitulatus, of Yanlé, Mt. Nimba. Includes
H. reticulatus, H. lagoensis and H. liberiensis.
Photo., Laurent (1).—H. lagoensis = H. guttulatus, Laurent (1) .- H. lamottei, larval development and metamorphosis fully descd. and figd., Arnoult & Lamotte.—H. lamottes sp. nov. p. 292, photo., from between Zouguépo and Serengbara, Mt. Nimba. Related to H. quinquevittatus, Laurent (1). — H. liberiensis = H. guttulatus, Laurent (1).—H. marmoratus, from Inyanga district, S. Rhodesia; note on colour variation, Fitzsimons.—Suggestion that the Ponds reed frog is a colour-variant, Wager (1).— Ponds reed frog is a colour-variant, Wager (1).—
H. m. quarrei, from Kasai, Belgian Congo, Laurent (2).
H. nasutus, from Inyanga district, S. Rhodesia,
Fitzsimons.—H. oeseri=H. fusciventris, Laurent (1).—
H. picturatus, of Mt. Nimba. Discussion on status
and relationships; includes H. baumanni and
H. festivus. Compared with H. c. concolor. Photo.,
Laurent (1).—H. polli, from Kasai, Belgian Congo,
Laurent (2).—H. reticulatus = H. guttulatus, Laurent
(1).—H. rosaceus, of Mt. Nimba region. Note on
colour and coology: photo. Laurent (1).—H. (1).—H. rosaceus, of Mt. Nimba region. Note on colour and ecology; photo., Laurent (1).—H. togoensis = H. c. concolor, Laurent (1).—H. verrucosus, compared with the painted reed frog. H. marmoratus, Wager (1).—H. viridiflavus nimbae subsp. nov. p. 290, photo., from Yanlé, Mt. Nimba, Laurent (1).—H. zonatus, larval development and metamorphosis desed. and figd., Arnoult & Lamotte.—H. zonatus, sp. nov. p. 276, photo., Yanlé, Mt. Nimba. Distinguished from H. acutirostris and H. pulcher, Laurent (1).

Kassina, Species listed. K. argyreivittis, K. somalica, K. benueana considered synonyms of senegalensis. K. angeli, K. owamboensis considered races of senegalensis. K. poweri considered a synonym of Hylambates maculatus. Species and subspecies recognised:—
cassinoides, maculata, cochranae, decorata, maculosa, maculifer, obscura, abyssinica, kounhiensis, pulchra, wittei, mertensi, kuvangensis, w. weali, w. fitzsimonsi, w. quinquevitata, modesta, poweri, thabanchuensis. Three distinct species occur in Cameroun:— s. senegalensis, cassinoides and decorata; note on biotope. K. decorata desed., notes on colour pattern variation, vomerine teeth, biotope and call; photos.; larva desed., mouth parts figd., Perret (2).—K. arygreivittis ruandae, from Ituri, Laurent (2).—K. ? decorata from Mt. Nimba; photo.; larva desed. and figd., Guibé & Lamotte (4).

Leptopelis, species recognised from French Cameroons:— notatus, ocellatus, brevirostris, calcaratus, palmatus, boulengeri, aubryi, milleoni, anchetae, viridis, Perret (1).—L. aubryi, L. viridis, from Mt. Nimba; photo.; characters; hand figd., Guibé & Lamotte (4).—L. brevirostris, of Fr. Cameroons. Note on colouration and reproduction. Photo., Perret (1).—L. notatus, includes L. tessmanni and probably also L. haugi. Notes on colour pattern variations, habitat in Fr. Cameroons, voice and reproduction; photos., Perret (1).—L. ocellatus, includes Nyctibates laevis and L. acuticeps. Note on colouration. Photo., Perret (1).

Philautus schmardanus, general note; fig., Morgan-Davies (3).

Phlyctimantis leonardi, from Boma Terr., Lower Congo; distinguished from verrucosus, Laurent (2).—See also under Hylambates.

Polypedetes, see under Rhacophorus.

Rhacophorus, larvae fractionation of radio iodine, Koshida (2).—R. appendiculatus appendiculatus first record from Negros Island, Alcala (2).—Polypedates [Rhacophorus] buergeri, notes on hatching glands, Yanai.—R. cruciger, R. leucomystax maculatus, range of colour variation and descriptions; photos, Silva.—R. c. cruciger, reproduction, metamorphosis and ecology desed. and figd., Morgan-Davies (1).—R. c. cruciger, R. c. eques, general notes; photo. of mominate form and description of nest and spawn, Morgan-Davies (3).—R. japonicus, of Amami group, Loo Choo Is., Koba.—R. l. leucomystax, of Thailand; photo., Taylor & Elbel.—R. microtympanum, life history notes, Morgan-Davies (3).—R. schlegelii, tadpoles abnormal gonadial differentiation due to gonad trophine injections, Iwasawa (2).—R. schlegelii amamiensis, of Amami group, Loo Choo Is., Koba.—R. schlegelii arborea metabolites in the perivitelline fluid, Arima.—Observations on movements, Kato (1).—Breeding habits, Kato (2).—Iodine accumulation in pigmented tissues, Koshida (1).—R. s. schlegeli, from Honshu, Japan, Alexander & Diener.

## RHINODERMATIDAE

Rhinoderma darwini, distribution, Capurro (1).

#### SCAPHIOPODIDAE

Scaphiopus. Absence of isolation mechanisms (call and genetic incompatibility) between S. holbrooki and S. hurteri, Blair (5).—S. bombifrone, figured with sonagram p. 186, Bogert.—Parasitism of tadpoles by Saprolegnia, Bragg, A. N. & W. N.—Of S. Dakota, Woodbury.—S. couchi, figured with sonagram p. 186,

a,

a,

0.

ı, B

m &

18, 1e, [t.

te

l). so

l). n-

10, rst tes ds, ge nd of m, ip, ife lii, to ileba. ine 1). in om

and ith by ota, Bogert.—From Gómez Farias, Tamaulipas, Martin.—S. hammondi figured with sonagram p. 186, Bogert.—[Scaphiopus] Spea h. hammondi, from Pima Co., Arizona, Williams & Chraplimy.—S. hammondi intermontanus, of Glen Canyon, Tanner.—S. holbrooki, of S. Florida; variation, habitat, call, breeding, food, Duellman & Schwartz.—† Scaphiopus holbrooki, from Pleistocene of Saber tooth cave, Florida, Holman (1).—S. holbrooki, formulae for body measurement relationships, Pearson (2).—S. h. holbrooki, reproductive behaviour, Hansen.—Population ecology, Pearson (1).—Distribution in the U.S.A., skull figured p. 314 f. 2a and relation to S. h. husteri discussed, Wasserman.—S. holbrooki husteri, on mass movement of tadpoles and daytime breeding, Bragg (1).—On geographic relations of Ozarkian amphibians, Dowling.—Distribution in the U.S.A.; cross breeding tests indicate that it should be regarded as a subspecies of S. h. holbrooki skull figured p. 314, Wasserman.—S. h. husteri × S. h. holbrooki indicates that husteri should be regarded as a subsp. of holbrooki, hybrid skull figured p. 314

## †SUPER ORDER

#### STEGOCEPHALI

†As yet undetermined fragmentary Labrinthodont plates—first ever recorded from China, Huene (1).

## **†ORDER EMBOLOMERI**

†Gyracanthus spine from the mid-carboniferous of Nova Scotia, Romer.

†Pholiderpeton (†) bretonensis sp. nov. p. 6, specific diagnosis valueless. Mid Carboniferous, Nova Scotia, Romar.

†Sagenodonus crushed quadrate found in Nova Scotia, mid Carboniferous deposits, Romer.

## ORDER RHACHITOMI


†Rhachitomi from the Triassic of India, Huene (2).

†Neorhachitomi from the Triassic of Central India, Huene (2).

## ORDER STEREOSPONDYLI

†Stereospondyli from the Triassic of Panchet and Mangli, India, Huene (2).

†Laidleria gen. nov. p. 68, text-figs. 16-19, of Trematosauridae, L. gracilis sp. nov. p. 68, Lower Triassic (Cynognathus-zone) Elucweewe in the Engeobo district, Eastern Cape Province. Kitching.



## THE ZOOLOGICAL RECORD

THE object of The Zoological Record is to give, by means of an annual volume, complete lists of the works and publications relating to zoology in all its branches that have appeared during the year preceding the issue of the volume, together with a subject and systematic index.

The complete volume can be obtained from the Society at the price of £10 0s. 0d. per copy.

The following volumes are available:-

## In Reprint Edition

Vols. 1-30 (1864-1893). Obtainable from Messrs. Butterworths, 4-5, Bell Yard, Temple Bar, London, W.C.2.

Complete set of the first 25 volumes, bound in cloth, price £221 0s. 0d.

Complete set of 30 volumes, in paper covers, price £256 10s. 0d.

Separate volumes, in paper covers, vols. 1-20 £8 0s. 0d. each; vols. 21-25 £8 18s. 6d. each; vols. 26-30 £10 10s. 0d. each.

The "Insecta" section, obtainable separately, paper covers. Price complete set Vols. 1-30 £157 5s. 0d.; single volumes £5 5s. 0d. each.

## In Original Edition

Vols. 65-71 (1928-1934) at £5 each. Vols. 85-91 (1948-1954) at £6 each. Vols. 92-95 (1955-1958) at £8 each.

For particulars of the availability and prices of separate Sections, application should be made to the Scientific Director, The Zoological Society of London, Regent's Park, London, N.W.1.

It should be noted that the separate Section Insecta can only be obtained from the Commonwealth Institute of Entomology, 56 Queen's Gate, London, S.W.7.

Nomenciator Zoologicus.—Lists the names of all the genera and subgenera in zoology from the 10th Edition of Linnaeus 1758 to the end of 1945, with a bibliographical reference to the original description of each. Price 17½ guineas, postage extra. Price of separate volumes: Vols. 1-4 (1758-1935), Vol. 5 (1936-1945), 3½ guineas each, postage extra.

# SEPARATE SECTIONS OF THE ZOOLOGICAL RECORD

In addition to the complete bound volume, the separate Sections may be obtained singly, bound in printed paper covers.

The prices of the Sections from Volume 96 are:-

										,
1.	Comprehensive	Zool	logy		***	•••			s. 4	d. 0
2.	Protozoa	•••	•••	***	•••	***			13	0
3.	Porifera	•••	•••	•••	•••		***	***	3	0
4.	Coelenterata	•••	•••	***	***		•••	•••	5	0
5.	Echinodermata	•••	•••	•••			•••	•••	4	0
6.	Vermes	•••	•••	•••	•••	***	•••	• • •	12	0
7.	Brachiopoda	•••	•••	•••	•••	***	•••	•••	4	0
8.	Bryozoa	•••	•••	•••	•••	•••		***	3	0
9.	Mollusca		•••	***		•••	•••	•••	16	0
10.	Crustacea	•••	•••	•••	•••		•••	•••	8	0
11.	Trilobita	•••	•••	•••	•••	•••		***	8	0
12.	Arachnida	•••	•••	•••	•••	•••			14	0
13.	*Insecta	•••	• • •		•••		•••	•••	60	0
14.	Protochordata	•••	•••	•••	•••		•••	***	3	0
15.	Pisces	***	• • •	•••	•••	•••	•••		11	0
16.	Amphibia	•••	•••	•••	•••		•••		12	0
17.	Reptilia	•••	•••	•••	•••	•••	•••	***	12	0
18.	Aves	•••	•••		•••	•••	***	•••	12	0
19.	Mammalia	•••	•••		•••	•••	•••	***	12	0
20.	List of New G	enera	and S	ubgene	ra	***		•••	4	0

\*Obtainable only from the Commonwealth Institute of Entomology, 56 Queen's Gate, London, S.W.7.

